CAB INTERNATIONAL
MYCOLOGICAL INSTITUTE
LIBRARY
IMI \ Books / SEA

HERB.

IMPERIAL BUREAU
16 JAN. 1929
OF MYCOLOGY

THE NORTH AMERICAN CUP-FUNGI



PHILLIPSIA CHARDONIANA SEAVER

The North American Cup-fungi

(Operculates)

BY

FRED JAY SEAVER, M.S., Ph.D.

CURATOR IN THE NEW YORK BOTANICAL GARDEN
AND
EDITOR OF MYCOLOGIA

ILLUSTRATED

NEW YORK
PUBLISHED BY THE AUTHOR
1928

Copyright, 1928
By Fred Jay Seaver

[Issued December 30]

LANCASTER PRESS, INC.
LANCASTER, PA.

To

THOMAS HUSTON MACBRIDE

President Emeritus, State University of Iowa

TABLE OF CONTENTS

PAGE

Preface 1–2

General features of the group 3–35

The following subjects are briefly discussed: 1. Phylogeny; 2. Gross morphology; 3. Origin of the apothecium; 4. Ascospore formation; 5. Alternation of generations; 6. Heterothallism and homothallism; 7. Spore discharge; 8. Heliotropism or phototropism; 9. Dehiscence of the ascus; 10. Significance of the mode of dehiscence; 11. Eccentricity of the ascostome; 12. Asexual reproduction; 13. Ecology; 14. Pyrophilous forms; 15. Coprophilous forms; 16. Spore germination; 17. Viability of the spores; 18. Mycophagy; 19. Classification; 20. Geographical distribution; 21. Nomenclature.

Signs 36

Taxonomic treatment 37–256

Bibliographical note 256

Explanation of plates 257–265

Index to recognized genera 267

Index to genera and species 267–284

PREFACE

The present work is the outgrowth of studies begun under Professors T. H. Macbride and B. Shimek, while a graduate student in the State University of Iowa, where three years were spent as a research Scholar and later Fellow in botany, and has been continued as a special problem, intermittently, up to the present time.

Official connection with The New York Botanical Garden, since the autumn of 1908, has afforded unusual opportunity to collect in the eastern states and the islands of the American tropics. Two expeditions have been made to the Bermuda Islands; the first in company with Dr. N. L. Britton and the late Stewardson Brown of Philadelphia, and the second with Professor H. H. Whetzel of Cornell University; one to the Island of Trinidad, just off the South American Coast, again in company with Dr. Britton; and one to Porto Rico and the American Virgin Islands, St. Thomas and St. Croix, as guest of the Porto Rican government. All of these trips have furnished valuable data for the work which is now being presented.

No private herbarium has been maintained but all of the collections made by the author have been merged with those of The New York Botanical Garden, which collection is very rich in American and European material. In the course of the work, collecting has been carried on from North Dakota to Colorado and New Mexico and east to New York and New England. No time has been spent in the far west. These collections have been supplemented by material sent, not only from every part of North America, but from nearly every part of the world.

Early in his work on Discomycetes the author came into contact, by correspondence, with Dr. E. J. Durand, then of Cornell University, and at that time an enthusiastic and, by all means, the foremost student of Discomycetes in America. Unfortunately his work was cut short by death before he had published any considerable part of his results on the operculate cup-fungi. It was the author's privilege to correspond and to exchange specimens with him up to the time of his death. His

extensive collection, containing much valuable information, is now at Cornell University.

Since there is no monograph of the operculate cup-fungi in America it seems not out of place to publish at this time a summary of our knowledge of the group which has accumulated after many years of more or less intermittent observation and study. In fact the author has been urged by mycologists to do so. If this work is successful it may be followed by a larger and more extensive one with a larger number of illustrations. Looking to this, criticisms and difference of opinion will be welcomed, especially where accompanied by observations, photographs and actual specimens.

One of the chief difficulties to be encountered in the study of this group of fungi is that, while some of the species are very common, others are found only once or twice in a lifetime, so that it is difficult to make a detailed study of such forms over an extended period. Others are so small that they are never seen unless one is looking for them. In such cases the amount of material which is available for study is very limited. Some apparently valid species have been collected, described, and often never seen again. Perhaps this adds interest and is all the more reason for the publication of such a monograph as this at the present time. It should be borne in mind, however, that there is still an inexhaustible field of study for any student who is sufficiently interested to follow it up.

The writer wishes to express his thanks to all those who have in any way contributed to the completion of the work to this point. To attempt to acknowledge them all by name would only result in embarrassing omissions and perhaps require more space than warranted.

Your coöperation is solicited in making this work a success and thereby paving the way for something better. If it does nothing more than to stimulate a renewed interest in this group of fungi the author will feel that his efforts have not been in vain.

NEW YORK,
MARCH, 1928

GENERAL FEATURES OF THE GROUP

1. Phylogeny

As the higher land plants are thought to have originated from lower aquatic forms, so the Discomycetes or cup-fungi, as well as the other higher fungi, which might be referred to as the "terrestrial thallophytes," are also believed to have been derived from aquatics. We would naturally look for the ancestors of these chlorophylless plants among the chlorophyll-bearing forms, the algae.

While most mycologists will concede that the fungi have evolved from the algae, directly or indirectly, there seems to be a difference of opinion as to whether their course has been a direct or an indirect one. On this point there are two theories, one that the ascomycetes have come directly from the red algae and the other that they have come indirectly through the Phycomycetes or lower fungi.

Those who hold the former view base their argument on the similarity which they seem to see between the sex organs and other structures of the two groups, especially the resemblance of the trichogyne of the ascomycetes and of the red algae. Those who oppose this view contend that they can see no analogy or homology between the unicellular trichogyne of the red algae and the multicellular trichogyne of the ascomycetes. Even in *Pyronema* where the trichogyne is itself unicellular the wall at the base of the trichogyne is a little barrier which has caused a great deal of discussion.

These and other views advanced are, however, only hypothetical and since the purpose of the present work is to treat the Discomycetes as *they are* rather than to philosophize on *how they might have come to be*, we will proceed with the task in hand and leave the question of their origin for those who have given more thought to this phase of the work. For a full discussion of the subject see Atkinson on the "Phylogeny of the Ascomycetes" (Ann. Missouri Bot. Gard. **2**: 315–376).

2. Gross Morphology

Like all of the higher fungi, the Discomycetes or cup-fungi present two distinct phases in their life histories, the growing

or vegetative and the fruiting or reproductive. The vegetative stage consists usually of a mass of loose hyphae which penetrate often for a great distance into the substratum in search of food which must be obtained in the organic form since these lowly plants have no chlorophyll. Occasionally as in the genus *Wynnea* a **sclerotium** is formed. From the mycelium or sclerotium the fruiting body is produced usually after a longer or shorter period of vegetative growth, the length of the period varying with the

FIG. 1. Stalked apothecia of *Paxina sulcata* in various stages of growth. Photograph by W. R. Fisher.

species. We often speak of the fungi as sporadic because they seem to appear at irregular intervals without rule or reason. The vegetative stage, however, is often persistent year after year in the same place where the substratum is constant, only the fruiting bodies being sporadic and even these occur with much more regularity than is usually supposed.

The fruiting stage of the cup-fungus is known as the **ascophore,** ascocarp, or ascoma and usually consists of a sessile or stalked disc or cup (FIG. 1) or in the Elvelaceae* of a distinct

*Various spellings have been employed for the generic name on which this family is based as follows: *Elvela*, *Elvella*, *Helvela*, and *Helvella*. The

stem and **pileus.** The disc or cup is referred to as the **apothecium,** while the pileus is simply a modified form of the apothecium. The apothecium consists of two primary parts, the fertile layer on the upper side of the disc or lining the cup and known as the **hymenium** and the sterile portion below or partially enclosing the hymenium, the **hypothecium,** the upper portion of which is sometimes referred to as the **subhymenial layer.** The outer part of the hypothecium is often modified into a more or less well-defined layer, the **excipulum.** The extreme outer part of the excipulum may be modified still further into a thin membrane, the **ectal layer.**

Fig. 2. Apothecia of *Bulgaria globosa.*

The apothecium is usually rather thin, only a few millimeters in thickness and often quite brittle. The thickness and consistency vary much in the different species, some being very soft while others are tough and leathery or gelatinous. The extreme is probably encountered in *Bulgaria globosa* (Fig. 2). In this species the apothecium is at first subglobose, tough and gelatinous, resembling in appearance a rubber ball. When mature the hymenium appears as a circular depression on the upper side. This species seems to be very rare in this country. It has been collected in Canada by Mr. J. W. Gussow, to whom I am indebted for the accompanying photograph.

original spelling has been adopted in this work and citations are made in this form regardless of the spelling used by the various authors.

The tissue of the hypothecium consists of an interwoven mass of mycelium known as **prosenchyma,** or the mycelium may be transformed into a paranchyma-like tissue, the **pseudo-parenchyma.** The one grades imperceptibly into the other. The cells of the excipulum may give rise to rigid or flexuous hairs which clothe the exterior of the cup.

The hymenium is composed of a palisade of sterile, club-shaped or filiform bodies, the **paraphyses** which are often pigmented giving color to the hymenium (see frontispiece), and larger cylindrical, club-shaped or ovoid bodies known as the **asci** in which the spores are produced in definite numbers in the following ratio: 2–4–8–16–32–64–128, etc. The variation in the form and markings of the spores will be discussed in a later chapter. Occasionally the tips of the paraphyses form a rather distinct layer, the **epithecium.**

3. Origin of the Apothecium

The formation of the ascophore is known to be preceded in a number of species by a definite sexual process so that it has come to be regarded as the sexual method of reproduction although the process has been demonstrated in a comparatively small number of species and in some of these it is not certain that the organs, though present, are actually functional. Whether sex organs exist in all species of the group but have not yet been found or whether they have been lost in many is a question which can be answered only after long and careful research.

One of the first species in which a sexual process was observed by the early workers, DeBary and Tulasne, is *Pyronema omphalodes*, probably because the sex organs are so very large and distinct in this particular form and so easily obtained, since the fungus probably occurs in every part of the world. At any rate, it has been found by the writer in every place that he has collected, both in continental North America as well as in the most obscure islands of the sea. All that is necessary for its growth is a burned area with sufficient moisture to insure growth. Detailed suggestions regarding the artificial growth of this fungus will be discussed in a later chapter.

The sex organs in *Pyronema* as described by DeBary consist of clusters or **archicarps** and **antheridia** produced side by side and later arranged in pairs. The archicarp consists of a large

globular cell or **oögonium** surmounted by a tubular outgrowth, the **trichogyne**, and attached to the mycelium by means of **basal stalk cells.** The antheridia are produced in a similar manner and are of about the same length as the oögonia but much more slender. The trichogyne at first grows upright but later curves about and partially surrounds the top of the antheridium, finally fusing with it. A wall is formed at the base of the trichogyne and about the same time the wall breaks down at the point of fusion of the trichogyne with the antheridium.



Fig. 3. Sex organs in *Pyronema* from DeBary. a, cluster of antheridia and oögonia. b, ascogonium producing ascogenous hyphae, with antheridium still attached.

After the fusion of the trichogyne with the antheridium the oögonium, which now becomes the **ascogonium,** swells up and produces outgrowths in a dozen places or more, which outgrowths become the **ascogenous hyphae** and eventually give rise to the asci. This is the process as described by DeBary (Fig. 3), but it remained for later workers to add the cytological details.

Both the oögonium and the antheridium are multinucleate as well as the trichogyne. The nuclei of the latter later degenerate. The nuclei from the antheridium gradually pass into the trichogyne. Although there is some difference of opinion on this point, it is generally conceded that the wall at the base of the trichogyne now breaks down, permitting the male nuclei to pass

into the oögonium, after which the wall is rebuilt, cutting off the oögonium once more as a single large cell. The male and female nuclei now pair up within the oögonium and eventually fuse, although again opinions differ as to whether they fuse in the oögonium or merely pass in pairs through the ascogenous hyphae, increasing by conjugate division, finally completing their fusion in the ascus. The latter view seems to be the most generally accepted one.

The vegetative cells at the base of the archicarps and the antheridia now put out a growth of sterile mycelium which surrounds the ascogenous hyphae, the two forming a compact mass which constitutes the apothecium.



FIG. 4. Sex organs of *Ascobolus carbonarius*. a, conidiophore with germinating conidium. b, antheridial conidiophore. c, coiled archicarp with trichogyne entwined about the antheridial conidium. d, ascogenous hyphae developing from enlarged cells of the archicarp.

As shown by DeBary and other early workers the archicarp in *Ascobolus furfuraceus* differs from that of *Pyronema* in that it consists of a chain of cells of variable size, one of which near the center functions as an oögonium and eventually produces the ascogenous hyphae.

While working as a student in The New York Botanical Garden, B. O. Dodge, at the suggestion of the writer, took up a study of the Ascobolaceae, which study has added much to our knowledge of the morphology of this group of cup-fungi. Perhaps the most interesting of his discoveries is the method of reproduction of *Ascobolus carbonarius* which, unlike most of the

members of this genus, grows on burnt ground and up to that time had not been studied morphologically by any American student. In fact the species is so obscure that it had been rarely collected. The reproductive process as described by Dodge is as follows: The mycelium of this species produces a large number of **conidia.** Some of these conidia while still attached to the mycelium germinate and give rise immediately to an archicarp which consists of a coil of swollen cells (Fig. 4) twenty to forty in number, growing larger then smaller in size as they extend outward, finally tapering off into a multicellular trichogyne which entwines itself about an antheridial conidium similar in appearance to the one from which the archicarp



Fig. 5. Sex organs in *Ascobolus magnificus*. a, antheridial branch. b, archicarp branch. c, ascogonium showing the beginning of the ascogenous hyphae. Figs. 4 and 5, after Dodge.

originated. While these are regarded as sex organs, and probably justifiably so, there is no definite proof that the multicellular trichogyne actually fuses with the antheridial conidium and no cytological details to show that these structures function as sex organs. About three cells of the archicarp nearest the stalk cell enlarge, the second cell of which produces the ascogenous hyphae and is regarded as the ascogonium.

Dodge has also made some interesting observations on *Ascobolus magnificus*, a species which he himself has described. This species was kept under observation for a long period in the laboratory. According to his report, the apothecia originate from club-shaped bodies which are produced in numbers throughout the colony, often in pairs. The bodies are at first one-

celled, later two- or three-celled. At this point one ceases to grow and apparently becomes the antheridium. The other continues to grow until it becomes a multicellular archicarp, the end of which tapers into a long slender, multicellular trichogyne which coils itself about and finally fuses with the antheridium (FIG. 5). One of the cells enlarges and becomes the ascogonium. This is similar in all essential details to that recorded for other species of *Ascobolus*. In this particular case the sex organs are so large that they can be dissected out with fine needles.

While definite sex organs have been observed in a number of species of operculate cup-fungi and the cytological details worked out in several, a great many are still to be investigated before we can draw any general conclusions as to the extent to which these organs are present and actually function.

4. Ascospore Formation

After the pairing or fusing of the male and female nuclei in the oögonium the ascogenous hyphae develop from the



FIG. 6. Beginning of ascus formation. a, basal cell. b, loop cell. c, tip cell.

ascogonium, the paired or fused nuclei passing into the ascogenous hyphae. Just what transpires between the time the nuclei pass into the ascogenous hyphae and the beginning of the formation of the asci from their ultimate branches it is difficult to say since the ascogenous hyphae intertwine in such a manner that it is almost impossible to follow any one through the entire process. The ultimate branch of the ascogenous hypha forms a loop consisting of three cells which might be designated as **basal** or **stalk cell, loop** or **dome cell,** and **tip** or **ultimate cell** (FIG. 6).

The basal and tip cells each contain one nucleus while the loop cell contains two. The two nuclei fuse and the loop cell elongates into the ascus. The original nucleus of the ascus which has resulted from the fusion just mentioned divides three or more times, usually giving rise to as many nuclei as there are to be spores in the ascus although some may degenerate or several be included in a single spore. After arranging themselves in the proper position in the ascus, each nucleus surrounds itself with a part of the cytoplasm of the ascus and becomes a spore, surrounded with a rather firm wall, the firmness of the wall varying much in different species.

The development of the spores by the process of free cell formation has been described in detail by R. A. Harper in his studies on *Pyronema* (Ann. Bot. **14**: 363–368). At the pole of each nucleus, after the final division of the fusion nucleus in the ascus, is an aster of fibers which persist and arrange themselves in the form of a wheel about the beak of the nucleus and finally cut out a nearly spherical mass of cytoplasm which encloses the nucleus and, with the development of the wall, becomes the young ascospore. A part of the cytoplasm of the young ascus is used in the formation of the spore and the remainder, which is known as the **epiplasm,** serves to nourish the young ascospore and to furnish material for the sculpturing of the wall, in those species which have sculptured spores.

All ascospores are at first smooth and many remain permanently so. In a number of species of the present group, however, the walls of the mature spores have elaborate markings of a very definite character. These seem to be formed by an accretion of epiplasm about the spore-wall. The process is well illustrated in *Sphaerosoma echinulatum* of the writer (which in this work is referred to *Boudiera*). After the spores are formed in this species, a definite sheath of epiplasm four to five microns thick appears about the spore and is known as the **exospore.** The sculpturing first appears on the surface of the spore-wall and gradually grows outward until it reaches the outer boundary of the exospore. Even after the sculpturing is completed a definite line persists which marks the outer limits of the original sheath. In other species such as *Lamprospora Crec'hqueraultii* the sculpturing seems to come about as a gradual growth after the spore is formed. So far as our observations have gone the spores of the operculate cup-fungi are permanently one-celled,

although the spores of the inoperculate forms are often multicellular.

5. Alternation of Generations

Assuming that the sex organs do function in the cup-fungi and taking as an illustration the best known form, *Pyronema omphalodes*, it is interesting to trace the alternation of generations as we know it in the fungi. The sporophytic generation would begin with the pairing of the nuclei in the ascogonium and continue through the ascogenous hyphae up to the reduction division in the ascus. The gametophytic generation, initiated at the conclusion of reduction division, is continued through the ascospores, mycelium, and up to the formation and fusion of the trichogyne with the antheridium, and the pairing of the nuclei in the oögonium.

6. Heterothallism and Homothallism

While heterothallism has been observed and established in the Phycomycetes and some of the higher Basidiomycetes, it is only recently that it has been demonstrated in the ascomycetes, including some of the species of operculate Discomycetes. Dodge states that in *Ascobolus magnificus* he was unable to obtain sex organs from one spore culture while they were abundantly formed on the mixing of two strains. In this case the sex organs were not produced in a definite line where the strains come together but the mycelia mixed so that they were produced throughout the culture but more abundantly where the two strains came first into contact.

Under the direction of Dodge, Edwin Betts has recently demonstrated heterothallism in *Ascobolus carbonarius* (Am. Jour. Bot. **13**: 427–432. 1926). While it was found impossible to produce ascocarps from single spore cultures, they could be produced by the intermingling of the mycelia from different spores, even those from the same ascus. From this it was concluded that sex factors are segregated at the time of the ascospore formation in the ascus and that probably four of the spores are of one sex and four of the other.

It has also been shown that in some species of ascomycetes which have only four spores in the ascus, two nuclei, one of each sex, may be included in the same spore. When such spores are germinated they produce a homothallic or monoecious mycelium, while a spore originally containing only a single nucleus will

produce a heterothallic, dioecious, or unsexual mycelium. While these facts have been demonstrated in one species of the ascomycetes, further investigation is necessary in order to determine whether this is true of the operculate Discomycetes. A number of species of this group are known to have usually, if not always, four spores instead of eight. Unfortunately these forms are not often collected in sufficient numbers for cytological and cultural study. This phase of the subject also offers excellent opportunity for research.

7. Spore Discharge

Once the spores are formed, the next step, so far as the fungus is concerned, is to "broadcast" them so that they

Fig. 7. Puffing of the spores in one of the inoperculate cup-fungi, *Sclerotinia* Photograph by W. R. Fisher.

may best serve the purpose for which they were designed, that of the reproduction and dissemination of the species. Probably there is no group of fungi in which the apparatus for spore dissemination is more highly developed than in the present one. Formed in a closed sac or ascus, usually of cylindrical form, they are held there for a time, apparently under high tension. Finally when conditions are just right, the ascus opens in an

orderly fashion and the spores are literally shot out. Just why so many asci discharge their spores in unison, it is difficult to state except that all naturally respond in a similar manner to the same stimulus. Just what this stimulus is no one has been able to decide with certainty. The contraction of the ascus after rupture would have a tendency to force the spores out. It was at one time thought that the Ascobolaceae ejected their asci but this was a mistake.

In the larger cup-fungi so many asci discharge their spores at once that it gives rise to a faintly audible puff and is apparent to the unaided eye as a cloud of smoke (FIG. 7). This phenomenon is well known to all students of fungi but it remained for Mr. W. R. Fisher of Cornell University to photograph the process and to him I am indebted for the accompanying illustration which has been previously published in Phytopathology. A. H. R. Buller believes that puffing does not take place under natural conditions. It is certainly frequently observed in the field although it may be due to unusual disturbance of some kind.

The force with which the spores are discharged has been indicated by Buller in his " Researches on Fungi " for *Ascobolus immersus*. His experiments showed that the maximum height of projection was 35 cm., the violence of the discharge probably not being exceeded by any other ascomycete.

8. Heliotropism or Phototropism

While the vegetative stage of the cup-fungi is able to grow in the absence of light, being chlorophylless, the fruiting stage invariably seeks the light and is positively heliotropic or phototropic. This is especially apparent in the asci themselves. At maturity, the asci often protrude half their length or more above the surface of the hymenium. This character was at first thought to be restricted to the Ascobolaceae and in fact was the distinguishing character of the family, as at that time known. It is, however, now known to be common to many if not all of the operculate cup-fungi although much more apparent in some than others. It is especially apparent in *Ascobolus immersus* because of the large size of the spores and asci and the fact that the dark color of the latter renders them more conspicuous by reason of contrast. When grown in the laboratory near the window the asci will be found to bend at a strong angle toward the light.

This heliotropic response of the ascus is most apparent in those species which have comparatively large spores. It is a prominent character in many species of the genus *Lamprospora* for, in spite of the small size of the plants of this genus, the asci and spores are relatively large, while in some of the large cups of the genus *Peziza* it is much less apparent because of the small size of the asci and spores.

If the asci protrude far above the surface of the hymenium the bending in response to light occurs far down on the ascus, while in those that protrude less strongly the bending is nearer the apex, interfering with the bilateral symmetry of the ascus.

9. Dehiscence of the Ascus

There are in the Discomycetes several ways in which the ascus may rupture in order to permit the spores to escape. On



Fig. 8. Methods of dehiscence of the ascus. a, operculate. b, bilabiate. c, inoperculate.

the bases of the dehiscence of the ascus, they may be classified as operculate and inoperculate forms. By the term operculate we mean the presence of a lid which is quite circular in form and of variable size by means of which the ascus opens (Fig. 8, a). The operculum almost invariably remains attached at one side after the discharge of the spores and often one or two spores may be left behind after the process is completed. The opening formed by the rupture of the ascus will be here designated as the **ascostome.**

The inoperculate method of dehiscence consists of a simple rupture which leaves the margin of the ascostome toothed and ragged after the spores have been ejected (Fig. 8, c) but with no sign of an operculum or lid.

A third type of dehiscence is known among a few of the cup-fungi and is referred to as the bilabiate method (FIG. 8, b). This consists of the rupture of the ascus by means of a transverse slit across the top so that the ascostome gaps open like a fish's mouth after the expulsion of the spores. This is characteristic of the genus *Streptotheca* and a few allied forms. Roughly this is considered as a modification of the operculate method and those forms which show this type of dehiscence are classed with the Operculates since otherwise they show more affinities with this group than with the Inoperculates.

10. Significance of the Mode of Dehiscence

The operculum was first observed by H. M. Crouan in some species of *Ascobolus* but he had not seen it in the larger Pezizaceae. It was at first regarded as one of the characters of *Ascobolus* and some species in which the operculum had been noted were transferred to *Ascobolus* for this reason. This is no longer looked upon as a generic character and species originally placed in *Ascobolus* are now treated as *Ascophanus*, *Humarina* and *Lamprospora*, the genus *Ascobolus* being used in a more restricted sense.

The inoperculate method of dehiscence was also observed by Crouan but no great importance was attached to it at that time. Later M. E. Boudier found the asci of many of the larger Pezizaceae to be operculate and he first suggested dividing the Discomycetes into two groups, the operculate and the inoperculate forms, and proceeded to show that the presence or absence of the operculum in the ascus was a character of fundamental importance. It is so regarded in this work and, so far as the writer knows, by all modern students of cup-fungi.

As pointed out by Boudier this seems to be a very natural division and there is little difficulty in determining the method of dehiscence although few workers at that time took the trouble to mention it. Boudier also made, in the opinion of the writer, a very important observation when he mentioned that in most cases the operculum was horizontal or at right angles to the long axis of the ascus while in a few species it was oblique.

That the separation of the cup-fungi into the Operculates (Operculatae) and Inoperculates (Inoperculatae) is a natural division and morphologically sound is indicated by the fact that associated with this character are certain other morpho-

logical " ear-marks " which run true to form so that it is not actually necessary to see the operculum in order to know, with reasonable certainty, that a cup-fungus belongs to the operculate group. These characters are as follows:

The spores of the Operculates are comparatively large, never septate, often warted or sculptured, and ranging from globose to broad-ellipsoid, more rarely fusiform but never very slender or filiform. The plants of this section usually occur on soil, dung, well-rotted wood and only rarely on hard woody stems.

The spores of the Inoperculates on the other hand are comparatively small, usually smooth, narrow and often very long and slender or filiform and often septate or showing a strong tendency to become so. The plants of this group are seldom terrestrial (except the Geoglossaceae) but usually occur on hard stems, wood or leaves. While the spores of this group are in a very few cases globose, they are, in such forms, very much smaller than the globose spores of the Operculates. There is also a decided difference in the type of hairs clothing the outside of the apothecia, when such are present.

Some collectors object to using this character as a basis of division because of the erroneous idea that it is a difficult one to observe. This is not the case, but even if it were, it is scarcely a valid excuse for failing to recognize so fundamental a difference.

11. Eccentricity of the Ascostome

As mentioned above, the operculum is usually situated at right angles to the long axis of the ascus while in some species it is oblique. In the course of this work the writer has noted this latter character in a number of species of tropical Discomycetes, especially those belonging to the genera *Phillipsia*, *Cookeina* and *Wynnea*. It has also been observed in *Plectania hiemalis*, a temperate form but one which shows a general relationship with the tropical forms just mentioned. It may occur in many others but these are the forms in which it has been especially noted. Perhaps too little importance has been attached to this character but, up to the present time, it has merely been accepted by the writer as one of the many morphological characters of certain species without attempting to explain the reasons for its existence.

In connection with his excellent researches on fungi Dr. Buller has offered a very interesting theory in an attempt to

account for the eccentricity of the ascostome. He believes that the phenomenon is heliotropic and that it will vary in the same species with the direction of the light. Following this theory, those species with deep cups in which the asci stand at an angle of 45 degrees, the operculum would be developed on the upper side of the ascus in order that the spores might be directed upwards instead of being shot across to the other side of the cup.

So far as the observations of the writer are concerned, there appears to be no relation between the depth of the cup and the eccentricity of the operculum and ascostome, since the latter character has been found to be very constant in species belonging to certain genera without regard to the form of the cup. A fine example of this is afforded in *Phillipsia Chardoniana* illustrated



FIG. 9. Eccentricity of the ascostomes in *Phillipsia Chardoniana*, shown on the frontispiece.

in the frontispiece of this work. In this species the apothecia are as flat as a pancake yet, so far as observed, the ascostomes are always eccentric or, at least, they are conspicuously and predominantly so (FIG. 9).

Another species in which the eccentric ascostome has been especially observed is *Cookeina Colensoi*, a shallow cup which occurs on wood in the tropics. So far as observed, the ascostome in this species is always located decidedly on one side and is unusually small compared with the size of the spores so that there must be a good deal of stretching when the spores are discharged through it (FIG. 10).

Having noted this character in various species of *Phillipsia* and *Cookeina* and recalling that both of these genera have much in common with the genus *Wynnea*, species of this latter genus were examined and the same conditions found to exist (FIG. 11). In all of these cases, however, it was necessary to work with

dried material, which is certainly much more unsatisfactory than fresh material would have been.

While no exhaustive observations have been carried on, it can be stated that the eccentricity of the ascostome is a character common, if not universal, in species of *Phillipsia*, *Cookeina* and

Fig. 10. Asci and spores of *Cookeina Colensoi* showing eccentric ascostomes and the comparatively large size of the spores.

Wynnea, which genera are essentially tropical although one species of *Wynnea* has been reported from the north. It has also been observed in *Plectania hiemalis*, a northern species but one which resembles *Cookeina* in many respects. It seems to

Fig. 11. Asci and spores of *Wynnea americana* showing eccentric ascostomes and comparatively large size of the spores.

make no difference whether the apothecium is plane, concave, regular or otherwise. This character appears also to be accompanied by certain other morphological characters. The spores are unusually large, fusoid, unequal-sided and in nearly every case marked with the peculiar striations consisting of light and

dark bands which have been mentioned so frequently by the writer in connection with his studies of tropical ascomycetes. Whether we consider the eccentricity of the ascostome as a morphological character or otherwise there is certainly a striking similarity in all of the species in which it has been observed by the writer, and we are inclined to regard this as one of the " earmarks " of certain genera and groups of operculate Discomycetes. Just how frequently it occurs in other genera than those indicated cannot be stated.

It still remains that if the asci in those species with flat apothecia have eccentrically placed ascostomes, it may be due to the fact that the light rays reached the plant in an oblique manner and that the ascostomes were adjusted accordingly. It is impossible to follow out this line of argument since the forms in which this observation has been made are tropical and we have had no opportunity to experiment with them in a growing condition. Even this theory would not explain the fact that the ascostomes are always eccentric in some species while never so in others, both of which have grown out-of-doors and been subjected to the same light influences. Also it has been noted that where the ascostome is eccentric the operculum usually remains attached at the lower margin of the ascostome where it would impede the passage of the spore in that direction.

Our observations, up to the present time, do not seem to fit in with Buller's theory, and we feel that further evidence is necessary before this can be accepted as an explanation of this interesting phenomenon in the species mentioned above. A careful study of those forms should be made in the field.

In all of the species in which the eccentricity of the ascostome has been noted there is a great discrepancy between the size of the ascostome and the spore which has passed through it (FIGS. 9–11). Since the ascostome is intact after the spores have been ejected from the ascus, there is only one conclusion—that the ascostome has stretched and contracted during the process of spore ejection. Another observation worthy of mention is that the spores in all of these forms are strongly fusoid. From these facts we might easily conclude that the small size of the ascostome, as compared with the spore, is no accident but is in some way concerned with the process of spore ejection. When the spore has passed half-way through the ascostome which is at that time stretched to the limit, the contraction of the ascostome

would have a tendency to accelerate the force with which the spore is ejected or "snapped out" of the ascus just as a seed of a citrous fruit may be snapped from between the fingers by bringing them quickly and tightly together. There must be a contraction between each spore ejection, the ascus operating like a machine-gun.

Another point that might be mentioned here is the fact that in all of these forms the spores are not only fusoid, but unequal-sided. Since they do not always lie in the same position in the ascus, this would have a tendency to throw the spores in different directions, which would be a decided advantage, as these particular forms have large, heavy spores as compared with other species of the group and would not be so easily blown about by the wind. While these latter facts seem to have no particular bearing on the eccentricity of the ascostome they are mentioned here because they have been noted in connection with that character.

12. Asexual Reproduction

While asexual spores have been reported for a few species of operculate cup-fungi, such as *Peziza vesiculosa* and *Peziza repanda*, on the whole this type of reproduction is conspicuous by its absence in many of the species of this group, or, at least, has not been observed. Plants of this section are in marked contrast to some of those of the Inoperculates such as *Sclerotinia* in which the reproduction by means of asexual spores has been developed to a very high degree. So far as *Sclerotinia* is concerned, this may be an adaptation to the parasitic habits of the plants of the genus, the conidia functioning as repeating spores for the rapid dissemination of the fungus while its host is in a susceptible stage.

The botryoide type of asexual spores has been reported by Dodge for one species of the present group, *Patella abundans* (Fig. 12). This is a surprise since this type of conidia is characteristic of one section of the species of *Sclerotinia*, a genus which in every other character is absolutely unrelated to *Patella*. Perhaps careful search will reveal asexual spores in cases where they have not been suspected up to the present time.

Since several of the operculate cup-fungi in which conidia have been found are heterothallic, it has been suggested that in this group the conidia are especially adapted to facilitate the bringing together of the plus and minus strains, thereby in-

suring sexual reproduction. It is easy to conceive that they might serve this purpose.

While asexual reproduction by means of conidia is comparatively rare in the present group, so far as our present knowledge goes, some are readily reproduced from spawn. This is especially true of the coprophilous forms. The dried substratum with the spawn can be kept dormant for long periods, while growth is resumed almost immediately on being restored to moist conditions.



FIG. 12. *Botrytis* type of asexual spores reported by Dodge for *Patella abundans*.

13. Ecology

With a few possible exceptions, the plants of this group are all saprophytic, although it is not easy to determine absolutely whether a fungus is parasitic or saprophytic without extended investigation. A few species of operculate cup-fungi which appear on dead foliage may have been parasitic on the living host, coming to fruit only after its death. *Pithya Cupressi* on the foliage of certain evergreens might be cited as one of this type. Also *Humarina axillaris* grows in the axils of the leaves of a moss and *Humarina ithacaensis* on the thallus of a liverwort. The great majority, however, are strictly saprophytic, growing on all kinds of decaying material, especially vegetable humus.

Even saprophytes are often very selective in their habits or tastes and grow only on certain types of decaying material. Several species of *Humarina* grow only on the humus of decaying

mosses and some species of *Lamprospora* show the same tendency. Others grow on decaying wood, often showing a preference for certain kinds of wood, while still others occur on burnt soil or on dung, being designated as pyrophilous and coprophilous forms respectively. Since the latter forms are of more than usual interest they will be treated more in detail.

14. Pyrophilous Cup-fungi

There are quite a number of operculate cup-fungi which occur in nature only on burnt places, especially where piles of



Fig. 13. Left, flask of heated soil with abundant growth of *Pyronema*. Right, same soil unheated but otherwise treated in the same manner produces no growth.

wood or brush have been burned. The most common of these is the well-known *Pyronema*, first encountered by the writer during his early work in Iowa. Without stopping to investigate the matter it was simply assumed that this fungus occurred on burnt places because it preferred ashes to grow on as other

species preferred wood which had been rotted rather than burned.

While working at The New York Botanical Garden as a student in 1906, the attention of the writer was called to a pink fungus overrunning the soil in the greenhouse which had been sterilized with steam. On examination, the fungus proved to be *Pyronema*. This exploded the previous assumption that the



FIG. 14. Culture of *Pyronema* on heated soil, suitable for class study, showing all stages from the clustered sex organs to mature apothecia.

fungus occurred on burnt places because of the ashes, for there were no ashes in the steam-sterilized soil. As a result of these observations experiments were undertaken which finally proved that *Pyronema* could be grown on any rich soil which had been heated to a high temperature either by dry heat or steam and later cooled, wet, and inoculated with the spores of the fungus.

In fact, nature usually supplies the spores provided we comply with the other requirements, but to make certain we usually supply them. Control experiments show that *Pyronema* cannot be grown on unheated soil, even though the same soil is used and all the other requirements are met (Fig. 13). It must be borne in mind, however, that the spores in this case do not require heating. In fact it is doubtful if they could withstand the process. It is only the soil that should be heated.

Later experiments conducted by the writer in coöperation with Dr. E. D. Clark (Mycologia 2: 109–124. 1910), formerly of the Department of Biological Chemistry of Columbia University, proved that the question of growth of *Pyronema* on heated soil was purely a food problem. The heating renders insoluble organic matter soluble, which material serves as an excellent medium for the growth of many fungi as well as *Pyronema*. The growth of *Pyronema* on steamed soil had been noted in Europe but, so far as we are aware, no one who knew the fungus had ever observed it in America. As full data on this subject have been published in preliminary papers it seems unnecessary to repeat them here.

Whether other species of pyrophilous cup-fungi grow on burnt soil for the same reason has not been demonstrated but it is suspected that this factor may influence them to a greater or lesser degree. Betts (l.c.) mentioned the fact that he could not produce sex organs in *Ascobolus carbonarius* on agar but that they developed readily when germinated spores were transferred to sterilized soil, indicating that the sterilized soil offered the better medium. *Pyronema*, however, fruits readily on agar.

15. Coprophilous Cup-fungi

This term is applied to those species which grow exclusively on the dung of animals, and some show a decided preference for the dung of certain kinds of animals. *Ascodesmis porcina* has been grown only on pig dung, first on material from Porto Rico and later on the same kind of dung from New Jersey. It is probable that such a species would grow on the dung of closely related animals but up to date it has been found only on the dung of the one kind. Others are not so discriminating in their tastes but thrive on almost any kind of excrement. That coprophilous forms can grow on other substrata is evident from the fact that they are produced in cultures on agar.

From the fact that the spores of the species of many coprophilous fungi do not germinate readily, it has been assumed that it is not only possible but necessary that they pass through the alimentary canal of the animal before they could be induced to germinate. Many experiments have been conducted to show that this is the case and there is little question that this process does stimulate germination.

The distribution of the spores in such forms is easily accounted for. Growing as they do on dung in pastures, the spores which are more or less sticky on being discharged from the ascus easily adhere to blades of grass and other forms of vegetation. The grass is eaten by cows or other animals and the spores gain entrance to the alimentary canal and, on being released, are ready for action. It is not unusual to find spores of coprophilous fungi on leaves of living plants in the field. The writer once spent considerable time trying to identify a supposedly parasitic fungus from Bermuda on blades of grass only to find later that they were the spore caps from *Pilobolus* which adhered so closely that they appeared to have grown there.

16. Spore Germination

The spores of many of the species of cup-fungi germinate readily and without any special treatment. In some of the species, however, as indicated above, the spores do not germinate by the ordinary method. This has already been referred to in the coprophilous forms, especially species of *Ascobolus*. The members of this genus are not all coprophilous, some occurring on wood and one species, *Ascobolus carbonarius*, on burnt ground and still others on naked soil.

On the assumption that the spores of the coprophilous species must pass through the body of the animal in order to stimulate them to grow, attempts have been made to simulate these conditions by artificial means and favorable results have been obtained by this method.

Dodge in his researches on Ascobolaceae, referred to above, tried many experiments and found that spores of several species of *Ascobolus*, including *Ascobolus carbonarius*, could be made to germinate in culture by heating them to a temperature of 50 to 70 degrees centigrade. The heat stimulus seemed to be a favorable substitute for the chemical stimulus of the alimentary canal. From this we might assume that *Ascobolus carbonarius*

has been evolved from a coprophilous species, the spores of which having been planted on the ground, stimulated by the heat of the fire, and supplied with a suitable nutrient medium released by the heating of the soil, grow and become adapted to this habitat. It differs from *Pyronema* which it resembles in habitat by the fact that the spores of the latter do not require the heat stimulus to induce germination.

17. Viability of the Spores

The length of time the spores of the cup-fungi will keep their viability probably varies greatly with different species. The writer has not carried on any extensive observations along this line but the one species *Pyronema omphalodes* has been experimented with and the spores found to keep their viability for nearly three years in an ordinary herbarium packet in the herbarium. How much longer they might have retained their power to germinate has not been determined.

This fact ought to be of especial interest to those who are interested in keeping this fungus on hand for cultivation in class use.

18. Mycophagy

The majority of the operculate cup-fungi are too small to be worthy of consideration as articles of diet. Many of the larger species are recommended by mycophagists and are used whenever they can be found in sufficient quantity. None, so far as the writer knows, are decidedly poisonous although one species has been under suspicion. This is the so-called *Gyromitra esculenta* (Fig. 15) which the writer here regards as merely a gyrose form of *Elvela infula*, at least so far as our American forms are concerned. It is interesting to note that this form which was originally designated as "esculent" should be the first one to be brought into disrepute. Of course our American form may be different from the European which goes under the same name. Some mycophagists claim to have eaten it without the slightest discomfort but, if used at all, one should proceed very cautiously.

Probably the most valuable species from the culinary point of view is *Morchella esculenta*, more commonly known as the "common morel," the "spring mushroom" or the "honeycomb fungus." In the northern states this species occurs almost exclusively in the month of May, hence the name spring mush-

room. Of the twenty-three specimens in our collections, nineteen were collected in May while the remainder which were obtained in April are usually from the southern states although they may occur in the north during the latter part of April if the season is an early one.

The morel is often found in abundance. Underwood (Trans. Ind. Hort. Soc. 1893: 63. 1894) reports having gathered eighty-two fruiting bodies around an old stump in his back yard in Indiana, and in an apple orchard, twice as many. He refers to the morel as "a storehouse of nitrogenous food as luscious



FIG. 15. *Gyromitra esculenta* gyrose form of *Elvela infula*. Photograph by P. J. Anderson.

as an oyster and for an equal weight containing even more nutritious matter." In the same article he states: "It is safe to say that the total natural production of morels in the state of Indiana during each spring is not less than 50,000 bushels, and may be easily double that. These, in a fair market, are worth $200,000 and in the best markets would command a much higher price. With the exception of a few plants eaten here and there by a family, this wealth of food, so much esteemed in foreign countries by peasant and wealthy epicure alike, is shamefully allowed to go to waste. The peasants of Germany call the fungi "manna of the poor" while the ancient Greeks justly regarded them as "food of the gods." Any bright boy

could make large wages collecting and selling these plants, and with little effort the people would learn to esteem them as they do oysters now. There is no reason why they should not be as commonly exposed for sale by grocers as they are in foreign countries."

Since the fruiting season of the morel is short, they could be collected and dried for future use, thus making use of the excess crop. In Europe they are often strung and hung in festoons in the houses, much the same method as used in drying apples in early days.

All of the species of *Morchella* are edible, not one of them having even been under suspicion. The following directions have been given for preparing and cooking: "Having washed and cleansed them from earth, which is apt to collect between the plaits or hollows of the plant, dry thoroughly in a napkin and put them in a saucepan with pepper, salt and parsley, adding or not a piece of ham. Stew for an hour, pouring in occasionally a little broth to prevent burning. When sufficiently done, bind with the yolks of two or three eggs and serve on buttered toast." These directions may be varied to suit the taste. They may be cleaned, stewed and fried in butter, the stewing being essential, since they are rather inclined to toughness.

19. Classification

A classification of any group of plants should serve two purposes: first, to offer a convenient arrangement of the species treated and, second, to show, so far as possible, the natural relationship existing between the various members of the groups and subgroups. The principles of classification have always been a bone of contention with the morphologist, as the idealist, on the one hand, who is clamoring for a classification based on fundamentals only, and the taxonomist on the other, who is compelled to use so-called superficial characters which may or may not show natural relationship.

Some morphologists even contend that there is no use trying to work out a real classification of any group of fungi until we know the life histories of all of the species composing the group, especially the details of sexual reproduction, assuming that it exists. Of course at the present time this is only a "pipe dream" and while waiting for the morphologist to work out his ideals the taxonomist will be compelled to use the characters which

present themselves in order to secure a convenient arrangement and trust that, to some extent at least, they may also be an index to natural relationship.

But what do we mean by superficial and fundamental characters? Superficial characters are those which may be easily detected by the casual student who does not have access to a fully equipped laboratory or even compound microscope and must rely largely upon a hand-lens or the naked eye in making his determinations of the species. The presence or absence of the operculum in the ascus may be regarded as a fundamental character but, as has already been shown, this rather obscure character is accompanied by several other more superficial ones which will enable the student to place the species in the proper group without actually seeing the operculum. It is not possible in every case to identify a species in the present group absolutely without the use of a compound microscope, but in many cases they may be placed in the proper section or in the genus and occasionally even referred to the species.

Superficial characters are often a sure index to more fundamental ones and there is no reason why they should not be used by the taxonomist, although there may be a great difference of opinion as to which should or should not be employed. The protrusion of the ascus was at one time used but at the present time is considered of no value as a taxonomic character. Whether the hymenium is exposed from the first (gymnocarpous) or is at first enclosed by the apothecium (angiocarpous) has been made a good deal of in the past but it does not now seem to have any particular value.

No attempt has been made here to revolutionize the classification of the cup-fungi, although it has been somewhat modified to suit the author's own ideas. No doubt a much better one may be devised in the future, but in the meantime the one employed here will serve as a medium through which our ideas regarding species of cup-fungi occurring in North America can be transmitted to the student. If this can be done it will have served its purpose.

The term cup-fungi or Discomycetes is used to include those ascomycetes in which the hymenium is freely exposed at maturity. On this character, the cup-fungi merge into the Hysteriales and Phacidiales although generally the term is restricted to the Pezizales. Following Boudier's suggestion, the entire group is

then divided into the Operculates and Inoperculates. Only the former are considered in this work and probably the operculate cup-fungi are about as well defined as any section of the fungi. The Elvelaceae and Geoglossaceae cannot be considered in the same section, as has been done by Lindau, since the one family is made up of operculate species and the other of inoperculate. Only two families of operculate cup-fungi are here recognized, the Pezizaceae and Elvelaceae.

The family Pezizaceae has then been divided into eight tribes, in the segregation of which spore characters have been permitted to play a part. As the details on which these tribes are based are treated fully in the body of the work which follows, it seems unnecessary to discuss them here.

20. Geographical Distribution

Like other higher fungi, many of the cup-fungi are cosmopolitan or widely distributed over the surface of the earth, apparently having little regard for climate, altitude or other factors which have such a powerful influence on the distribution of the higher plants. Some, however, seem to be restricted to tropical regions but here the climatic influence may be only indirect. As already indicated, most of the operculate cup-fungi are saprophytic, growing on decaying wood and humus resulting from decaying plant remains. As the climate determines the nature of the phanerogamic flora, so it determines the type of humus found in any given region. If a saprophyte chances to be a selective feeder, its occurrence will be governed accordingly, and might be thus indirectly influenced by climate.

Where a saprophyte is not too selective in its habits and the substratum is essentially the same it seems to thrive equally well in tropical and temperate regions. While collecting in the Rocky Mountains the writer was impressed with the number of species found at high altitude there which were identical with those collected at low altitude around New York City. On the other hand, forms which were naturally confined to certain kinds of coniferous wood or branches were restricted to those regions where this particular conifer grew. The altitude seemed to be only an indirect factor in determining the distribution of the fungus. Species of *Lamprospora* were found among mosses at snow-line in the mountains and a few hundred feet above sea level in New York, where the substrata were apparently similar.

Again in Trinidad, which is almost under the tropical sun, while some species were found which were restricted to the tropics, the writer was frequently surprised to find what appeared to be *Peziza badia* of the north. While some mycologists might regard this as a geographical species, no morphological difference could be detected. This was especially common in Trinidad on clay pits where clay had been burned for the surfacing of the roads.

Perhaps the most interesting genus from the standpoint of distribution is *Lamprospora*, which might be regarded as the writer's "pet" genus since he has made a special effort to collect and study the species of this particular form. Because of their small size, the plants of this genus are usually overlooked by mycologists. Having had great difficulty in determining the species because of inadequate descriptions, the writer in 1914 put out a preliminary monograph calling attention to some of the peculiar markings of the spores.

Following this, Dr. Ethel McLennan and Miss Isabel Cookson (Proc. Roy. Soc. Victoria **38**: 70–76. 1926) have reported from Australia several species described as new in our preliminary monograph. These are as follows: *Lamprospora tuberculata; Lamprospora tuberculatella; Lamprospora Maireana;* and *Lamprospora areolata* Seaver, var. *australis*. Of these *Lamprospora Maireana* is also known from North Africa as well as North America. A number of other species of the genus are found in Europe, America and Australia and will probably be found to be world wide. It is difficult to draw general conclusions because so little attention is given to these obscure forms.

During the two collecting excursions to the Bermuda Islands, the cup-fungus most commonly collected was a European species, *Lamprospora Planchonis*, which has never been found on the mainland of North America, or at least has never been reported. This is one of the larger species of the genus and would not have been overlooked if it occurred here as abundantly as it does in Bermuda. From the standpoint of geographical distribution, there seems to be no more reason for its occurrence in Bermuda than in continental North America. When we recall that the soil of Bermuda is made up of disintegrated coral limestone, the nature of the substratum might have something to do with its occurrence there. Under just what conditions it occurs in Europe has not been learned.

Probably the ease with which the spores of the cup-fungus, as well as fungi in general, can be blown about by the wind has much to do with their dissemination over the surface of the earth. From the observations made, it would seem that where the substrata are essentially identical the most of the species of this group will grow in any part of the world, without regard to climate. It is possible, however, that some of the strictly tropical species may be directly influenced by climate.

21. Nomenclature

In general the writer has attempted in the following treatment to follow the rules of the American Code of Nomenclature, which is based on the principle of priority. While the application of the rules of any code will result in some temporary eruptions in a group of this size, we feel that the application of the rule of priority at least tends toward stability. The few radical changes which have been made in the present work in order to conform to rules have been discussed in a preliminary paper, but some of them perhaps should be dealt with a little more in detail here.

There are those in the mycological world who are avowedly opposed to the strict application of any code of nomenclature on the ground that such application would result in change, as of course it does in some cases. This small group of mycologists proposes to follow "usage," without regard to priority, on the plea that we would thereby avoid change of names. Now if the application of the principle ot so-called usage would accomplish what is claimed for it, there might be some excuse for following these suggestions. But the term "usage" is merely "a snare and a delusion" as can be shown by an attempt to apply this principle in the cup-fungi.

Since there is no extensive monograph of the operculate cup-fungi in America, there is naturally no well-established usage in this country. Whether we follow usage or priority, we must look largely to Europe for precedent. There have been several outstanding figures in the study of Discomycetes of Europe, to whom we may look for inspiration and information, of a taxonomic nature, in the study of this group of fungi. Among these are William Phillips, M. C. Cooke, and George Massee of England; H. Rehm and G. Lindau of Germany; C. C. Gillet and M. E. Boudier of France; and P. A. Saccardo of Italy. While all of these have contributed much to our knowledge of the taxon-

omy of this group of fungi, in the light of our present knowledge, none of them can be accepted as an infallible guide.

Although Boudier suggested in 1879 the presence or absence of the operculum in the ascus as a fundamental basis on which to separate the Discomycetes into two large sections, his contemporaries in other European countries seem to have overlooked or disregarded the suggestion. Both Lindau of Germany and Saccardo of Italy adopted classifications of their own, each different, and each grouping operculate and inoperculate species promiscuously in the same family and often in the same genus, entirely disregarding this most fundamental of morphological characters.

The effect of this diversity of treatment on nomenclature might best be illustrated by taking as an example the genus *Lamprospora*, as treated in this work. The name *Lamprospora* was proposed for this genus by De-Notaris in 1864. In 1869, Fuckel introduced the name *Crouania*, which, however, was untenable, since it had previously been used for another plant genus. In 1889, Saccardo suggested the name *Barlaea*, but later finding that it was untenable changed it to *Barlaeina*. Rehm followed Saccardo's suggestion in the use of *Barlaea* but he, also, later found that it was untenable and proposed the use of the generic name *Detonia*. Lindau, recognizing the untenability of the name *Barlaea*, proposed the substitution of *Plicariella*. Boudier continued to use the name *Lamprospora*.

This was the situation which confronted the writer in 1914, when he undertook a preliminary monograph of this genus for North America. At least three names were in common use in Europe: *Lamprospora* by Boudier and his followers; *Barlaea* or *Barlaeina* by Saccardo and his adherents; and *Plicariella* by Lindau. Rehm's suggestion to substitute *Detonia* for the genus had not gained any wide following. On the basis of usage, what would one do under these circumstances? The writer followed what appealed to him as being the most logical course and adopted the name *Lamprospora* on the ground of priority as well as usage. Since that time, this name has come into quite general use throughout the world. Yet, notwithstanding this fact, advocates of usage in this country have actually recommended the overturn of the genus *Lamprospora*, substituting for it *Detonia* because Rehm proposed it, although he himself never followed the suggestion. If this were done we would have had

twenty-three new combinations in this genus in the present work, instead of two, by the use of the name *Lamprospora*. Thus we see that the advocates of usage, which really means nothing, often defeat the very end for which we are all working.

Some have attempted to adopt Saccardo's Sylloge Fungorum as a basis for usage. Although this is a most valuable index, it can scarcely be so used in the present group, for, as already pointed out, Saccardo has adopted a system of classification which leaves entirely out of consideration the most important morphological character, that of the presence or absence of the operculum in the ascus. The genus *Bulgaria* as treated by Saccardo contains both operculate and inoperculate species, which according to present-day students of Discomycetes belong in different sections of the cup-fungi. Not only are operculate and inoperculate species placed in the same genus but the inoperculate are in this case treated first and have come to be regarded as typical, although the genus was originally established on an operculate species.

After considering the above and other suggestions of the advocates of usage, their propositions have been discarded as absolutely impracticable and not at all conducive to the ends which we all have in view, stability of nomenclature. Even the most radical change made in the present work, the substitution of *Patella* for *Lachnea*, could not have been avoided by the most ardent advocates of usage, for even they would not tolerate the use of the same name for two different plant genera.

The writer, however, is merely stating his position in the matter and has no desire to force his views upon anyone else. Those who have been in the habit of calling *Patella scutellata* a *Lachnea* will doubtless continue to do so whether the name is tenable or not. The writer, himself, will probably continue to do the same by force of habit. The tenable name has been adopted in this work, believing that it is just as easy for the new student of the younger generation to learn the right name as the wrong one, assuming that the one adopted here is right. By right we mean in accordance with the principles laid down by some well-formulated code. So far as the writer is aware, the name *Lachnea* would not be recognized as tenable under any existing code.

In following the rules of the American Code of Nomenclature, the writer, therefore, feels that he has adopted the course which will bring about the least possible confusion.

SIGNS

μ = the Greek mu, which indicates micromillimeters, one of which equals .001 of a millimeter or 1/25,400 of an inch.

mm. = a millimeter or 1000 μ.

cm. = a centimeter or 10 mm.

§ = subgenus.

– = dash between extreme measurements.

× = used between numbers representing length and breadth.

Order **PEZIZALES**

Ascophores consisting of discoid or cup-shaped apothecia, or more rarely clavate, columnar, or pileate; apothecia devoid of associated algal cells (except in the Patellariaceae, which family may contain either algicolous or non-algicolous species), free or more rarely seated on a subiculum or springing from a sclerotium, ranging in size from a fraction of a millimeter to several centimeters, variously colored, the hymenium concave, plane, or convex, and circular or subcircular in form, more rarely elongate or star-shaped, either enclosed by the apothecium or the excipulum when young or free from the first, very poorly developed or partially to entirely enclosing the hymenium when young, usually expanding at maturity leaving the hymenium freely exposed or, in a few cases, remaining closed until pierced or ruptured by the maturing asci, the tissue of the hypothecium either composed of loosely interwoven hyphae, prosenchymatous, or giving rise to a parenchyma-like tissue, pseudoparenchyma, or pseudoparenchymatous below and filamentous above; pileus bell-shaped, saddle-shaped, but never cup-shaped, even or irregularly convoluted or corrugated, surmounted by the hymenium; substance fleshy, leathery, cartilaginous, or corneous; asci ovoid to cylindric, two- to many-spored, operculate or inoperculate, rarely bilabiate; spores globose, ellipsoid, fusiform or filiform, one- to many-celled, hyaline or variously colored, yellowish, violet, brown or more rarely olivaceous, smooth or variously sculptured, echinulate, verrucose, tuberculate, reticulate, ringed or marked with irregular ridges; paraphyses filiform to clavate, simple or branched, variously colored.

Asci operculate (opening by a circular lid at the apex), or more rarely by a transverse slit which gives to the open ascus a bilabiate appearance; in the latter the ascus is surrounded just below its apex by a thickened ring or collar. — Section 1. Operculates.

Ascophores consisting of cup-shaped or discoid apothecia; apothecia sessile or more rarely stipitate. — Family 1. Pezizaceae.

Ascophores pileate, subglobose or columnar (never cup-shaped or discoid); pileus saddle-shaped or bell-shaped, always supported by a distinct stem. — Family 2. ELVELACEAE.

Asci inoperculate (opening by a simple pore at the apex). — Section 2. INOPERCULATES.*

* Not treated in this work.

Family 1. PEZIZACEAE

Apothecia cup-shaped to discoid or more rarely convex, sessile or stipitate, variously colored, externally naked or clothed with hairs; substance fleshy, waxy, leathery, cartilaginous or corneous; hairs varying from a soft tomentum to stiff bristles, hyaline or colored; asci cylindric to ovoid, operculate or more rarely opening by a transverse slit at the apex giving the open ascus a bilabiate appearance, occasionally marked by a thickened ring or collar near the apex, two- to many-spored; spores globose, ellipsoid or fusiform, hyaline or colored, the color ranging from yellowish to brown, violet, or more rarely greenish, smooth or variously sculptured; paraphyses present, filiform to clavate, hyaline or variously colored.

Spores globose, hyaline to pale-brown. — Tribe 1. SPHAEROSPOREAE.

Spores ellipsoid to fusoid or fusiform, more rarely subglobose.

Spores becoming violet, later brown to blackish. — Tribe 2. ASCOBOLEAE.

Spores permanently hyaline or at most pale-brown.

Spores becoming reticulate at maturity, and often apiculate. — Tribe 3. ALEURIEAE.

Spores smooth or sculptured but never reticulate.

Apothecia small, rarely exceeding 1 cm. in diameter and often less than 1 mm., sessile or subsessile.

Apothecia naked, or at least not conspicuously setose or hairy. — Tribe 4. HUMARIEAE.

Apothecia conspicuously setose or hairy. — Tribe 5. LACHNEEAE.

Apothecia medium to large, usually exceeding 1 cm. in diameter when mature.

Apothecia strongly unequal sided or split on one side. — Tribe 6. OTIDEEAE.

Apothecia normally symmetrical in form.

Apothecia usually stipitate and densely hairy or tomentose; substance usually tough and leathery. Tribe 7. SARCOSCYPHEAE.

Apothecia sessile or short-stipitate, not densely hairy or tomentose; substance usually fleshy and brittle. Tribe 8. PEZIZEAE.

Tribe 1. SPHAEROSPOREAE. Apothecia fleshy to cartilaginous, sessile or stipitate, naked or hairy; hairs either bristly or flexuous, hyaline or colored; asci usually 8-spored but often with a part of the spores undeveloped, often strongly protruding, operculate; spores globose or rarely very slightly elongated, smooth or sculptured, hyaline or faintly colored, yellowish to pale-brown or blackish; paraphyses filiform to clavate.

Plants growing on humus or living plants, not restricted to the dung of animals.
- Apothecia clothed with well-developed bristles or flexuous hairs.
  - Apothecia red, brown or greenish; substance soft and fleshy or waxy. 1. SPHAEROSPORA.
  - Apothecia black or brownish-black; substance tough or cartilaginous. 2. PSEUDOPLECTANIA.
- Apothecia naked, not clothed with well-developed hairs.
  - Apothecia subglobose, externally entirely covered by the hymenium; substance approaching cartilaginous. 3. SPHAEROSOMA.
  - Apothecia discoid to convex-hemispheric with the hymenium restricted to the upper surface and sides; substance soft and fleshy or waxy.
    - Plants growing on the ground or on humus.
      - Hymenium strongly convex, approaching *Sphaerosoma* in form. 4. BOUDIERA.
      - Hymenium concave, plane or slightly convex. 5. LAMPROSPORA.
    - Plants growing on living or recently killed foliage and branches of coniferous trees; substance tough and *Ciboria*-like. 6. PITHYA.

Plants normally growing only on the dung of animals.
- Spores becoming pale-brown as they mature. 7. ASCODESMIS.
- Spores remaining permanently hyaline. 8. CUBONIA.

Tribe 2. ASCOBOLEAE. Apothecia usually small, often less than 1 mm. or rarely exceeding 1 cm. in diameter, usually

yellowish or greenish at least when young, very soft and fleshy, the paraphyses and asci often surrounded by a mucilaginous, yellowish or greenish slime, never conspicuously hairy, occurring on dung or more rarely on soil, etc.; asci broad-clavate, operculate, strongly protruding at maturity and very conspicuous on account of their large size and the dark-colored spores which they contain; spores becoming some shade of violet or purple as they mature, later brown or occasionally blackish and opaque, ellipsoid or rarely almost globose, smooth or sculptured, spore-sculpturing taking the form of warts or reticulations, the reticulations consisting of ridges or crevices; paraphyses variable.

Spores free in the ascus. 9. ASCOBOLUS.
Spores united in a ball in the ascus. 10. SACCOBOLUS.

Tribe 3. ALEURIEAE. Apothecia small to medium or large, usually bright-orange or red, more rarely dark-brown, sessile or stipitate, growing on soil or humus, naked or hairy; hairs soft and flexuous or bristly, hyaline or brown; asci cylindric or subcylindric, operculate, 8-spored; spores hyaline to pale smoky-brown, at maturity sculptured; spore-sculpturing consisting of ridges which usually give rise to very regular net-like reticulations; paraphyses usually strongly thickened above and colored.

Apothecia not clothed with colored hairs.
Hymenium bright-orange; spores hyaline. 11. ALEURIA.
Hymenium dark-brown; spores smoky-brown. 12. ALEURINA.
Apothecia clothed with colored hairs. 13. MELASTIZA.

Tribe 4. HUMARIEAE. Apothecia small, not usually exceeding 1 cm. in diameter and often less than 1 mm., soft and fleshy or rarely slightly tough, scutellate to discoid or occasionally with the hymenium strongly convex, usually sessile, occurring on the dung of animals, soil, or debris; asci cylindric to clavate or ovoid, in a few cases marked with a ring or collar near the apex, operculate or bilabiate, 4- to many-spored; spores ellipsoid, hyaline or subhyaline, smooth, or minutely sculptured; paraphyses variable in size and color.

Apothecia closely attached to the substratum to the extreme margin, usually occurring on wood. 14. PSILOPEZIA.
Apothecia attached to the substratum by the center only, occurring on soil, humus or the dung of animals.
Vegetative mycelium superficial, forming a conspicuous web on burnt places. 15. PYRONEMA.

Vegetative mycelium immersed in the substratum, not usually restricted to burnt places.
Asci 8-spored.
Apothecia fleshy, very soft.
Plants coprophilous. 16. ASCOPHANUS.
Plants humicolous. 17. HUMARINA.
Apothecia tough, *Ombrophila*-like. 18. PSEUDOMBROPHILA.
Asci more than 8-spored.
Asci bilabiate or opening by a transverse slit at the apex. 19. STREPTOTHECA.
Asci operculate or opening by a circular lid at the apex.
Asci and spores small; spores not exceeding 15 μ long. 20. RYPAROBIUS.
Asci and spores large; spores 35–38 μ long. 21. THECOTHEUS.

Tribe 5. LACHNEEAE. Apothecia rarely exceeding 1 cm. in diameter, scutellate to discoid, sessile, or rarely stipitate, externally clothed with hyaline or colored hairs; asci operculate, 8-spored but often with a part of the spores undeveloped; spores smooth or sculptured, hyaline to pale-yellowish; paraphyses variable.

Apothecia partially to entirely buried in the ground. 22. SEPULTARIA.
Apothecia superficial on the substratum.
Asci with an external thickened ring near the apex; plants occurring on the dead foliage of cedar. 23. PSEUDOPITHYELLA.
Asci without external thickened ring; plants occurring on soil, wood, humus, or the dung of animals.
Apothecia densely clothed with brick-red hairs which give to the whole plant a red appearance. 24. PERROTIA.
Apothecia clothed with hyaline or brown hairs which appear white or black to the naked eye.
Hair simple (without cross-walls). 25. LASIOBOLUS.
Hairs septate, usually erect and bristle-like. 26. PATELLA.

Tribe 6. OTIDEEAE. Apothecia unsymmetrical, either elongated or split on one side to the base, occurring singly or springing in clusters from a sclerotium, rather tough and subcoriaceous, externally smooth, pruinose, or occasionally tomentose; asci cylindric, operculate, 8-spored; spores smooth but often marked

longitudinally by light and dark bands; paraphyses straight or hooked at their apices and usually enlarged above.

Apothecia springing in clusters from a sclerotium which is buried in the ground. 27. WYNNEA.
Apothecia usually isolated and not springing from a sclerotium.
  Spores striate with light and dark bands. 28. PHILLIPSIA.
  Spores not striate. 29. SCODELLINA.

Tribe 7. SARCOSCYPHEAE. Apothecia stipitate, or rarely sessile, tough and inclined to coriaceous (not shrinking much in drying), bright-colored or brown to blackish, growing attached to buried or partially buried sticks (except the genus *Paxina*, the plants of which genus grow on the ground), externally densely tomentose or bristly; stem very variable in length, its length often depending on the depth at which the substratum is buried, occasionally wanting, slender or more often thick, even or corrugated; asci operculate, 4–8-spored; spores smooth, but often marked with longitudinal light and dark bands as in *Wynnea* and *Phillipsia*, or minutely sculptured; paraphyses usually slender but variable.

Apothecia bright-colored, yellow to scarlet within; mycelium white.
  Spores striate and hairs fasciculate; plants tropical. 30. COOKEINA.
  Spores not striate and hairs not fasciculate; plants temperate. 31. PLECTANIA.
Apothecia dark-colored, brown to blackish; mycelium usually dark-colored.
  Apothecia with a thick gelatinous hypothecium. 32. BULGARIA.
  Apothecia without thick gelatinous hypothecium.
    Apothecium opening with a star-shaped aperture or splitting *Geaster*-like; growing attached to sticks. 33. URNULA.
    Apothecium not opening with a star-shaped aperture; growing on the ground. 34. PAXINA.

Tribe 8. PEZIZEAE. Apothecia sessile or with an inconspicuous stem, externally smooth or nearly so, occasionally slightly tomentose, varying in size from one to several centimeters in diameter; hymenium concave, plane or convex, bright-colored or more commonly brownish or blackish; asci cylindric or subcylindric, operculate, 8-spored; spores ellipsoid to fusoid, hyaline or rarely faintly colored, smooth or sculptured; paraphyses very variable, usually filiform.

Apothecia superficial.
  Apothecia medium-sized, shaped like an acorn-cup with a short slender stem. 35. GEOPYXIS.
  Apothecia medium to large, sessile or with a stout stem-like base, often irregularly contorted.
    Spores apiculate.
      Apothecia attached to the substratum by numerous root-like processes. 36. RHIZINA.
      Apothecia not attached by numerous root-like processes. 37. DISCINA.
    Spores not apiculate. 38. PEZIZA.
Apothecia entirely buried in the ground, at least when young. 39. SARCOSPHAERA.

1. SPHAEROSPORA Sacc. Syll. Fung. 8: 188. 1889.

Peziza § *Sphaerospora* Sacc. Michelia 1: 594. 1879.
Sphaerosporula Kuntze, Rev. Gen. Pl. 3³: 530. 1898.

Apothecia becoming scutellate or occasionally discoid, externally clothed with hairs; hymenium plane, slightly concave or rarely convex, usually red or brown, more rarely greenish or pale-yellow; hairs bristly or flexuous, hyaline or colored; asci cylindric to clavate, 8-spored; spores globose, hyaline or colored, smooth or becoming sculptured; spore-sculpturing consisting of warts or spines; paraphyses slender below, clavate above.

Type species, *Peziza trechispora* Berk. & Br.

Spores becoming sculptured at maturity.
  Spore-sculpturing consisting of warts.
    Apothecia bright-red; hairs rigid. 1. *S. trechispora.*
    Apothecia greenish; hairs flexuous. 2. *S. flavovirens.*
  Spore-sculpturing consisting of spines. 3. *S. perplexa.*
Spores remaining permanently smooth.
  Apothecia reddish-brown.
    Apothecia small, 1–6 mm. in diameter, forming congested masses on burned places. 4. *S. brunnea.*
    Apothecia large, reaching 1 cm. in diameter, scattered, on sandy soil in woods. 5. *S. hinnulea.*
  Apothecia pale-yellow. 6. *S. monilifera.*

1. Sphaerospora trechispora (Berk. & Br.) Sacc. Syll. Fung. 8: 188. 1889.

Peziza trechispora Berk. & Br. Ann. Mag. Nat. Hist. **18**: 77. 1846.
Lachnea trechispora Gill. Champ. Fr. Discom. 77. 1879.
Peziza asperior Nyl. Not. Fauna Fl. Fenn. **10**: 21. 1869.
Leucoloma asperior Rehm, Ber. Naturh. Ver. Augsburg **26**: 6. 1881.

Sphaerospora asperior Sacc. Syll. Fung. **8**: 188. 1889.
Sphaerosporula asperior Kuntze, Rev. Gen. Pl. 3^3: 530. 1898.
Sphaerosporula trechispora Kuntze, Rev. Gen. Pl. 3^3: 530. 1898.
Ciliaria asperior Boud. Hist. Class. Discom. Eu. 62. 1907.
Ciliaria trechispora Boud. Hist. Class. Discom. Eu. 62. 1907.

Apothecia gregarious or scattered, at first subglobose, expanding and becoming scutellate to discoid, reaching a diameter of 1 cm.; the margin slightly elevated, externally clothed with rigid hairs which project and form a conspicuous fringe about the margins of the apothecia, giving them a dark-brown color; hymenium bright-red, plane or very slightly concave, smooth; the excipular cells irregularly polygonal, the walls pale-brown; hairs sharp-pointed, reddish-brown with transmitted light, 2–5-septate, variable in length but reaching an extreme of 300–400 μ and a diameter of 20–26 μ at the broadest point a little above the base, contracted toward the point of attachment and often forked at the base; asci cylindric or subcylindric above, gradually tapering below into a stem-like base, reaching a length of 200–250 μ and a diameter of 25–30 μ; spores 1-seriate, at first smooth and containing one large oil-drop, finally becoming sculptured, reaching a diameter of 20–26 μ; spore-sculpturing consisting of rounded wart-like tubercles about 2 μ in diameter; paraphyses clavate, septate, reaching a diameter of 10 μ at their apices, reddish.

On rich soil, especially among mosses.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to Colorado; also in Europe.

ILLUSTRATIONS: Boud. Ic. Myc. *pl. 375, 378;* Cooke, Mycographia *pl. 33, f. 129;* Gill. Champ. Fr. Discom. *pl. 64;* Phill. Brit. Discom. *pl. 7, f. 41;* Grevillea **3**: *pl. 33, f. 103;* Jour. Linn. Soc. **31**: *pl. 16, f. 20, 21;* Rab. Krypt.-Fl. 1^2: 1029, *f. 1–4;* Ann. Myc. **4**: 208, *f. 28;* 209, *f. 29.*

2. **Sphaerospora flavovirens** (Fuckel) Sacc. Syll. Fung. **8**: 189. 1889.

Plicaria flavovirens Fuckel, Symb. Myc. Nacht. **2**: 64. 1873.
Peziza flavovirens Cooke, Mycographia 68. 1876.
Phaeopezia elaeodes Clements, Bot. Surv. Nebr. **4**: 16. 1896.
Sphaerosporula flavovirens Kuntze, Rev. Gen. Pl. 3^3: 530. 1898.
Sphaerospora Durandi Rehm, Ann. Myc. **2**: 36. 1904.

Apothecia scattered, at first shallow cup-shaped, becoming scutellate to discoid, reaching a diameter of 5 mm. to 2 cm.,

externally pale-brown, slightly verrucose and tomentose; hymenium concave, plane or slightly convex, olivaceous; hairs flexuous, septate, pale-brown; asci cylindric or subcylindric, reaching a length of 200–300 μ and a diameter of 15–18 μ; spores 1-seriate, containing one large oil-drop, at first smooth, becoming rough, reaching a diameter of 12–16 μ; spore-sculpturing consisting of minute warts or indistinct reticulations; paraphyses filiform below, enlarged above, septate, reaching a diameter of 5–6 μ at their apices.

On damp sand-banks.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to Nebraska; also in Europe.

ILLUSTRATION: Cooke, Mycographia *pl. 31, f. 122.*

3. **Sphaerospora perplexa** Seaver, sp. nov.

Apothecia gregarious, sessile, when young closely nestling in the substratum, at first closed and globose, expanding and becoming discoid, reaching a diameter of 1–3 mm.; externally clothed about the margin with fascicles of adpressed, pale-brown hairs which often form fifteen or twenty tufts about the border of the disc; hymenium plane, bright-orange and slightly darker than the outside of the apothecium; hairs flexuous, club-shaped, sparingly septate and often constricted at the septa, smooth or very slightly roughened, reaching a diameter of 21–33 μ at their apices; asci cylindric or subcylindric above, rather abruptly tapering below into a short stem-like base, reaching a length of 350 μ and a diameter of 23 μ; spores 1-seriate, densely granular within, becoming sculptured, reaching a diameter of 21–23 μ; spore-sculpturing assuming the form of sharp spines (resembling those of *Lamprospora Crec'hqueraultii*); paraphyses stout, reaching a diameter of 7–8 μ.

On bare soil in woods.

TYPE LOCALITY: Woods near Yonkers, New York.

DISTRIBUTION: Known only from the type locality.

Specimens recently sent from Sweden by Dr. John Axel Nannfeldt appear to conform with the above, which would extend the range of distribution if such should prove to be the case.

This may be only an abnormal form of *Lamprospora Crec'hqueraultii*, the development of hairs being due to unusual conditions. Except for the hairs it is identical with that species.

4. **Sphaerospora brunnea** (Alb. & Schw.) Massee, Brit. Fungus-Fl. **4**: 295. 1895.

Peziza brunnea Alb. & Schw. Consp. Fung. 317. 1805.
?Peziza schizospora Phill. Grevillea **3**: 31. Hyponym. 1874.
?Peziza sphaeroplea Berk. & Curt.; Cooke, Bull. Buffalo Soc. Nat. Sci. **2**: 289. 1874.
Peziza confusa Cooke, Bull. Buffalo Soc. Nat. Sci. **2**: 291. 1875.
Lachnea confusa Phill. Grevillea **18**: 83. 1889.
Sphaerospora confusa Sacc. Syll. Fung. **8**: 190. 1889.
?Barlaea schizospora Sacc. Syll. Fung. **8**: 116. 1889.
Scutellinia brunnea Kuntze, Rev. Gen. Pl. **2**: 869. 1891.
Ciliaria confusa Boud. Hist. Class. Discom. Eu. 62. 1907.
Sphaerosporula confusa Kuntze, Rev. Gen. Pl. 3^3: 530. 1898.

Apothecia gregarious or more often densely crowded, forming congested masses several cm. in diameter, at first subglobose, expanding and becoming scutellate or discoid, reaching a diameter of 5–6 mm., externally brownish, clothed with short hairs; hymenium plane or slightly concave, pale- to dark-brown; the excipular cells irregularly polygonal, more or less elongated near the margin, pale-brown, often reaching a diameter of 25–30 μ; hairs scarcely projecting above the margin of the apothecium, rigid, rather sharp-pointed, subhyaline or pale-brown, 1–4-septate, reaching an extreme length of 200 μ and a diameter of 6–8 μ, the young hairs often very irregularly crooked or bent; asci cylindric or subcylindric above with a long slender stem-like base, reaching a length of 160–200 μ and a diameter of 20 μ; spores 1-seriate, containing one large oil-drop, hyaline, smooth or very minutely roughened, reaching a diameter of 15–18 μ; paraphyses clavate, reaching a diameter of 6–7 μ at their apices, filled with brown granules.

On burnt ground and charcoal beds.

TYPE LOCALITY: Europe.

DISTRIBUTION: Massachusetts to Iowa and Alabama; also in Europe.

ILLUSTRATIONS: Boud. Ic. Myc. *pl. 379;* Cooke, Mycographia *pl. (20, f. 80?); 32, f. 124;* Alb. & Schw. Consp. Fung. *pl. 9, f. 8.*

5. **Sphaerospora hinnulea** (Berk. & Br.) Massee, Brit. Fungus-Fl. **4**: 294. 1895.

Peziza hinnulea Berk. & Br. Ann. Mag. Nat. Hist. IV. **7**: 433. 1873.
Peziza scutelloides Ellis, Bull. Torrey Club **9**: 18. 1882.
Lachnea hinnulea Phill. Brit. Discom. 219. 1887.
Barlaea hinnulea Sacc. Syll. Fung. **8**: 117. 1889.
Sphaerospora scutelloides Sacc. Syll. Fung. **8**: 188. 1889.
Sphaerosporula scutelloides Kuntze, Rev. Gen. Pl. 3^3: 530. 1898.

Apothecia gregarious or scattered, never densely crowded, at first subglobose, expanding and becoming scutellate to discoid, reaching a diameter of 5 mm. to 1 cm., externally clothed with rigid hairs which project above the margin, giving it a dark-brown color; hymenium slightly concave or plane, reddish-brown; the excipular cells irregularly polygonal, reaching a diameter of 30–35 μ, the cell walls pale-brown; hairs straight or irregularly bent, rather sharp-pointed or obtuse, pale-brown, 3–5-septate, reaching an extreme length of 200 μ and a diameter of 10 μ at the base; asci cylindric or subcylindric above with a slender stem-like base reaching a length of 200 μ and a diameter of 20 μ; spores at first irregularly 2-seriate, separating and becoming 1-seriate, containing one large oil-drop and often several smaller ones, smooth, reaching a diameter of 15 μ; paraphyses strongly enlarged above, septate, brown.

On sandy soil in woods, often among mosses.

TYPE LOCALITY: Europe and Connecticut.

DISTRIBUTION: New York to Iowa; also in Europe.

ILLUSTRATIONS: Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 5, f. 1;* Cooke, Mycographia *pl. 13, f. 52.*

EXSICCATI: N. Am. Fungi *838, 2142.*

6. **Sphaerospora monilifera** (Berk. & Curt.) Seaver, comb. nov.

Peziza monilifera Berk. & Curt.; Berk. Jour. Linn. Soc. **10**: 367. 1868.
Pyronemella monilifera Sacc. Syll. Fung. **8**: 194. 1889.

Apothecia becoming subdiscoid, reaching a diameter of 1 mm., externally pale-yellow, clothed with delicate hairs; hymenium pale-yellow, plane or nearly so; hairs septate, hyaline or faintly colored; asci cylindric; spores smooth, hyaline, reaching a diameter of 15–16 μ; paraphyses slightly enlarged above.

On the ground.

TYPE LOCALITY: Cuba.

DISTRIBUTION: Known only from the type locality.

ILLUSTRATION: Cooke, Mycographia *pl. 40, f. 158.*

2. PSEUDOPLECTANIA Fuckel, Symb. Myc. 324. 1869.

Caloscypha Boud. Bull. Soc. Myc. Fr. **1**: 103. 1885.
Otidella Sacc. Syll. Fung. **8**: 99. 1889.
Melascypha Boud. Hist. Class. Discom. Eu. 56. 1907.

Apothecia gregarious or scattered, sessile or stipitate, medium-sized, usually exceeding 1 cm. in diameter, fleshy to cartilaginous,

externally clothed with short, slender, flexuous, often coiled or twisted hairs, giving to the outside of the apothecium a tomentose appearance, blackish to orange or greenish; hymenium concave, orange or blackish; asci cylindric or subcylindric, 8-spored; spores perfectly globose, smooth, hyaline or faintly colored; paraphyses very slender, straight or curved at their apices.

Type species, *Peziza nigrella* Pers.

Apothecia entirely black or brownish-black.
- Long-stipitate, sparingly clothed with straight or slightly flexuous hairs. 1. *P. vogesiaca.*
- Short-stipitate or sessile, densely clothed with coiled hairs. 2. *P. nigrella.*

Apothecia orange or occasionally with a greenish tint about the outer margin. 3. *P. fulgens.*

1. **Pseudoplectania vogesiaca** (Pers.) Seaver, Mycologia **5**: 300. 1913. (PLATE 1, FIG. 1.)

?Peziza fuscocana Alb. & Schw. Consp. Fung. 312. 1805.
Peziza vogesiaca Pers.; Moug. & Nest. Stirpes Crypt. *584.* 1818.
Peziza melania Pers. Myc. Eu. **1**: 239. 1822.
Peziza melaena Fries, Syst. Myc. **2**: 60. 1822.
Peziza spongiosa Peck, Bot. Gaz. **5**: 35. 1880.
Bulgaria spongiosa Peck, Bot. Gaz. **6**: 241. 1881.
Pseudoplectania melaena Sacc. Syll. Fung. **8**: 165. 1889.
Pulparia spongiosa Sacc. Syll. Fung. **8**: 612. 1889.
?Otidella fuscocana Schröt. Krypt.-Fl. Schles. **3²**: 49. 1893.
Melascypha melaena Boud. Hist. Class. Discom. Eu. 56. 1907.

Apothecia gregarious or scattered, stipitate, cup-shaped or occasionally nearly plane, the margin often wavy, reaching a diameter of 2–3 cm. and a depth of 1–2 cm., externally very sparingly tomentose, becoming strongly wrinkled when dry, black or blackish; substance subcartilaginous; hymenium dark olivaceous-brown, concave or nearly plane; stem very variable in length but occasionally reaching 2–3 cm. and about 3 mm. thick, black, attached at the base by a dense mass of coarse, dark-brown hairs; tomentum consisting of short, brown, flexuous, sparingly septate hairs; asci cylindric above, with a very long stem-like base, reaching a length of 200-275 μ and a diameter of 16–18 μ; spores 1-seriate, containing one large oil-drop, becoming very pale-brown, reaching a diameter of 12–14 μ; paraphyses slender, slightly enlarged above and coiled or hooked at their apices, reaching a diameter of 3–4 μ.

1. PSEUDOPLECTANIA VOGESIACA
2. PSEUDOPLECTANIA NIGRELLA

On decaying wood in coniferous woods, especially among *Sphagnum*.

TYPE LOCALITY: Europe.

DISTRIBUTION: Vermont to Washington; also in Europe.

ILLUSTRATIONS: Boud. Ic. Myc. *pl. 343;* Cooke, Mycographia *pl. 49, f. 193;* Rab. Krypt.-Fl. 1^3: 1030, *f. 1;* Mycologia **5**: *pl. 109* (upper figure) and *pl. 10;* Gill. Champ. Fr. Discom. (suites).

2. **Pseudoplectania nigrella** (Pers.) Fuckel, Symb. Myc. 324. 1869. (PLATE 1, FIG. 2.)

Elvela hemisphaerica Wulf. Beob. Nat. Ges. Nat. Fr. Berl. 2^1: 141. 1787.
Not *Peziza hemisphaerica* Weber. 1780.
Peziza nigrella Pers. Syn. Fung. 648. 1801.
Lachnea nigrella Gill. Champ. Fr. Discom. 78. 1874.
?Peziza stygia Berk. & Curt. Grevillea **3**: 153. 1875.
Plectania nigrella Karst. Act. Fauna Fl. Fenn. **2**: 119. 1885.
?Pseudoplectania stygia Sacc. Syll. Fung. **8**: 166. 1889.
Otidella nigrella Schröt. Krypt.-Fl. Schles. 3^2: 48. 1893.
Sphaerospora nigrella Massee, Brit. Fungus-Fl. **4**: 296. 1895.

Apothecia gregarious or occasionally closely crowded, sessile or substipitate, at first closed and subglobose, expanding and becoming shallow cup-shaped to discoid, margin often wavy and slightly incurved, externally clothed with very fine hairs, black or blackish, reaching a diameter of 5 mm. to 1.5 cm.; hymenium concave or nearly plane, smooth, brownish-black; hairs very long but usually coiled and twisted, giving to the exterior of the apothecium a slightly woolly appearance, reaching a diameter of 4–6 μ and of nearly uniform thickness throughout their entire length, sparingly septate, pale-brown; asci cylindric or subcylindric with a long stem-like base, reaching a length of 300–325 μ and a diameter of 15 μ; spores reaching a diameter of 12–14 μ, usually containing one large oil-drop and several smaller ones; paraphyses enlarged at their apices and filled with brown coloring matter, reaching a diameter of 4 μ.

On decaying wood in coniferous woods, especially among *Sphagnum*.

TYPE LOCALITY: Europe.

DISTRIBUTION: New Jersey to Manitoba, Wisconsin, Alabama, and Jamaica; also in Europe and Australia.

ILLUSTRATIONS: Boud. Ic. Myc. *pl. 344;* Cooke, Mycographia *pl. 31, f. 120;* Gill. Champ. Fr. Discom. *pl. 65;* E. & P. Nat. Pfl. 1^1: 179, *f. 146, A–B;* Rab. Krypt.-Fl. 1^3: 1030, *f. 2;* Mycologia **5**:

pl. 109 (lower figure); Krombh. Abbild. *pl. 16, f. a, b;* Proc. Roy. Soc. Victoria II. **38**: *pl. 4, f. 4–6; pl. 6, f. 2.*

3. **Pseudoplectania fulgens** (Pers.) Fuckel, Symb. Myc. 324. 1869.

Peziza fulgens Pers. Myc. Eu. **1**: 241. 1822.
Peziza cyanoderma DeBary in Rab. Fungi Eu. *516.* 1863.
Aleuria fulgens Gill. Champ. Fr. Discom. 41. 1879.
Otidella fulgens Sacc. Syll. Fung. **8**: 99. 1889.
Barlaea fulgens Rehm in Rab. Krypt.-Fl. 1^3: 930. 1896.
Plicariella fulgens Lindau in E. & P. Nat. Pfl. 1^1: 180. 1897.
Detonia fulgens Durand, Bull. Torrey Club **29**: 459. 1902.
Caloscypha fulgens Boud. Hist. Class. Discom. Eu. 54. 1907.

Apothecia scattered or more rarely crowded, regular or irregular in form, occasionally unequal-sided or split *Otidea*-like, substipitate and attached to the substratum by a dense mass of coarse mycelium which penetrates into the humus binding it together, reaching a diameter of 5 mm. to 2.5 cm., externally pale-orange, becoming greenish or olivaceous, especially about the margin, sparingly clothed with poorly developed hairs; hymenium orange, brighter than the outside of the apothecium; hairs short, clavate, consisting of a few loosely united cells; asci cylindric above, gradually tapering below into a long stem-like base, reaching a length of 100 μ and a diameter of 10–12 μ; spores at first 2-seriate, separating and finally becoming 1-seriate, reaching a diameter of 6–8 μ, hyaline; paraphyses slender, scarcely enlarged above, filled with orange granules.

On soil in coniferous woods.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to Montana and California; also in Europe.

ILLUSTRATIONS: Boud. Ic. Myc. *pl. 319;* Cooke, Mycographia *pl. 53, f. 209;* E. & P. Nat. Pfl. 1^1: 179, *f. 146, C, D;* Gill. Champ. Fr. Discom. *pl. 38;* Pat. Tab. Fung. *f. 377.*

The bright-orange color of this species is in marked contrast to that of the other two species recorded here both of which are black. The outside of the apothecium is often overcast with an olive tint. It is a beautiful species but only rarely collected.

3. **SPHAEROSOMA** Klotzsch; Dietrich, Fl. Boruss. *Pl. 467.* 1839.

Sphaerozosma Corda, Ic. Fung. **6**: 52. 1854.
Ruhlandiella P. Henn. Hedwigia **42**: (23). 1903.

Apothecia subglobose, sessile or substipitate; hymenium covering all of the exposed surface of the apothecium, brownish; substance fleshy or slightly cartilaginous; asci broad-cylindric to subclavate, 8-spored; spores globose, at first smooth, becoming sculptured, slightly colored; paraphyses rather slender.

Type species, *Sphaerosoma fuscescens* Klotzsch.

1. **Sphaerosoma hesperium** (Setchell) Seaver, comb. nov. (PLATE 2, FIG. 3.)

?Sphaerosoma fuscescens Klotzsch; Dietrich, Fl. Boruss. *Pl. 467.* 1839.
?Ruhlandiella berolinensis P. Henn. Hedwigia **42**: (22). 1903.
Ruhlandiella hesperia Setchell, Univ. Calif. Pub. **4**: 116. 1910.

Apothecia subglobose or reniform, short-stipitate or attached by a small tuft of mycelium near the center of the under side, reaching a diameter of 2–6 mm.; hymenium covering all of the exposed surface of the apothecium, pale-brown; asci broad-cylindric or clavate above, narrowed below, reaching a length of 250–300 μ and a diameter of 28–30 μ; spores at first 2-seriate or irregularly crowded near the end of the ascus, gradually separating and becoming 1-seriate, at first smooth, becoming sculptured, reaching a diameter of 20–27 μ; spore-sculpturing taking the form of reticulations, the reticulations 2–3 μ deep as indicated by the projections about the periphery of the spore, the ridges often appearing as spines; paraphyses filiform, curved at the ends, enlarged and cemented together, forming a rather firm epithecium.

On the ground under leaf-mould.

TYPE LOCALITY: Berkeley, California.

DISTRIBUTION: California; also in (Europe?).

ILLUSTRATION: Univ. Calif. Publ. Bot. **4**: *pl. 15, f. 1–4.*

4. **BOUDIERA** Cooke, Grevillea **6**: 76. 1877.

Apothecia sessile, at first subglobose, becoming expanded and subdiscoid, or convex-hemispheric and *Sphaerosoma*-like, rather small, reaching a diameter of 2–5 mm.; hymenium occupying the upper surface only or covering the entire exposed surface and

in vertical section forming a complete semi-circle; asci very broad, 8-spored; spores globose, becoming sculptured; spore-sculpturing consisting of reticulations or spines, pale-yellowish; paraphyses very stout and much thickened at their apices.

Type species, *Boudiera areolata* Cooke & Phill.

Apothecia subdiscoid. 1. *B. areolata.*
Apothecia convex-hemispheric. 2. *B. echinulata.*

1. **Boudiera areolata** Cooke & Phill.; Cooke, Grevillea **6**: 76. 1877. (PLATE 2, FIG. 1.)

Plicaria tracheia Rehm, Ascom. *451.* Hyponym. 1878.
Barlaea areolata Massee, Brit. Fungus-Fl. **4**: 398. 1895.

Apothecia scattered, sessile, globose when young, becoming expanded and subdiscoid, reaching a diameter of 2–3 mm., externally smooth, whitish; hymenium occupying the upper surface, convex, dark-brown; asci subcylindric to clavate, tapering below into a long stem-like base, strongly protruding at maturity and appearing in dried specimens as minute, whitish granules over the surface of the hymenium; spores at first 2-seriate or irregularly bunched near the apex of the ascus, gradually separating, finally becoming 1-seriate, at first smooth and surrounded by a hyaline band, gradually becoming sculptured, reaching a diameter of 30–35 μ including the spines; spore-sculpturing assuming the form of reticulations which are often more or less indistinct or broken, at maturity echino-reticulate, the spines reaching a length of 4–5 μ; paraphyses stout, septate, brown.

On damp soil in swampy places.

TYPE LOCALITY: North Wales.

DISTRIBUTION: Maine; also in Europe.

ILLUSTRATIONS: Grevillea **6**: *pl. 97, f. 14, 15;* E. & P. Nat. Pfl. 1^1: 192, *f. 154, A;* Boud. Ic. Myc. *pl. 417;* Phill. Brit. Discom. *pl. 9, f. 53;* Rab. Krypt.-Fl. 1^3: 1110, *f. 1–5;* Mycologia **6**: *pl. 123, f. 5–6.*

2. **Boudiera echinulata** Seaver, comb. nov. (PLATE 2, FIG. 2.)

Sphaerosoma echinulatum Seaver, Jour. Myc. **11**: 2. 1905.

Apothecia scattered, gregarious or occasionally crowded, at first almost globose and regular in outline, becoming expanded and finally convex-hemispheric, the lower surface nearly plane, whitish and attached to the soil by very delicate hyphae, very easily detached, reaching a diameter of 5–8 mm.; hymenium



1. BOUDIERA AREOLATA
2. BOUDIERA ECHINULATA
3. SPHAEROSOMA FUSCESCENS

beginning by a dark-brown spot in the center of the upper surface and expanding until it covers all of the exposed surface of the apothecium, convex-hemispheric, even or becoming convolute and umbilicate, reddish-brown to brown, and velvety in appearance due to the stout paraphyses which extend far beyond the asci; asci clavate above, tapering below, never protruding so far as noted, reaching a length of 300–350 μ and a diameter of 40–50 μ; spores at first irregularly bunched near the end of the ascus, gradually spreading, finally becoming 1-seriate, at first smooth, filled with oil-drops and surrounded by a broad, hyaline band, gradually becoming sculptured, reaching a diameter of 25 μ excluding the sculpturing, or 35 μ including the sculpturing; spore-sculpturing becoming more pronounced with age, often assuming the form of reticulations, later strongly echinulate, the spines reaching a length of 4–5 μ and a diameter of 2–2.5 μ at their base, straight or bent at their apices and often with a trace of ridges connecting their bases; paraphyses very stout, septate, brownish, reaching a diameter of 12–15 μ at their apices.

On damp clayey soil in swampy place in the margin of woods.

TYPE LOCALITY: Iowa City, Iowa.

DISTRIBUTION: Iowa; also in Europe.

ILLUSTRATIONS: Jour. Myc. **11**: *pl. 75;* Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 6;* Mycologia **6**: *pl. 123, f. 7–10;* Bull. Acad. Sci. Cracovie **1909**: 88, *f. 3*, and 89, *f. 4.*

EXSICCATI: Rehm, Ascom. *1601.*

This species was collected in abundance in a ravine near Iowa City when the writer first undertook the study of the Discomycetes in 1904, and although it was later reported from central Europe, it has never again been encountered in America. The species differs only slightly from the preceding, which was collected once in Maine by Dr. Roland Thaxter. This is the extent of our knowledge of this genus in America.

The writer suspects that *Sphaerosoma alveolatum* McLennan & Cookson (Proc. Roy. Soc. Victoria II. **35**: 153. 1923) is closely related to, if not identical with, the above.

DOUBTFUL SPECIES

Boudiera marginata Phill. & Hark. Bull. Calif. Acad. Sci. **1**: 25. 1884. This species is reported on rabbit's dung from California and is said to be

just as striking as *B. areolata*. The spores are described as globose, 20 μ in diameter, rufous and areolate. No specimen has been seen, but the description is suggestive of *Ascodesmis*.

5. **LAMPROSPORA** De-Not. Comm. Critt. Ital. **1**: 388. 1864.

Crouania Fuckel, Symb. Myc. 320. 1869. Not *Crouania* Agardh. 1842.
Plicaria Fuckel, Symb. Myc. 325. 1869.
Barlaea Sacc. Syll. Fung. **8**: 111. 1889. Not *Barlaea* Reichenb. f 1877.
Detonia Sacc. Syll. Fung. **8**: 105. 1889.
Phaeopezia Sacc. Bot. Cent. **18**: 218. 1884 (in part).
Curreyella Massee, Brit. Fungus-Fl. **4**: 401. 1895.
Plicariella Lindau in E. & P. Nat. Pfl. **1**[1]: 179. 1897.
Barlaeina Sacc. & Sydow in Sacc. Syll. Fung. **14**: 30. 1899.
Pulvinula Boud. Hist. Class. Discom. Eu. 69. 1907.

Apothecia small to medium-sized, shallow cup-shaped or more often scutellate or discoid, externally smooth or verrucose but never clothed with well-developed hairs; hymenium concave, plane or convex, usually bright-colored, some shade of red orange or yellow, more rarely pallid or dark-colored, purple or brownish-black, often roughened by the protruding asci; asci usually clavate or subcylindric, 8-spored; spores globose or subglobose, at first smooth, becoming sculptured or remaining permanently smooth, hyaline or faintly colored; spore-sculpturing very variable, consisting of spines, minute warts, coarse tubercles or ridges, the ridges often assuming the form of reticulations, or in one case forming rings about the spore; paraphyses filiform or clavate, straight or curved.

Type species, *Ascobolus miniatus* Crouan.

Apothecia small, not exceeding 5 mm. in diameter (usually 1 or 2 mm.); spores hyaline or subhyaline.
Spores sculptured (reticulate, spinulose, verrucose, tuberculate, or annulate).
Spore-sculpturing consisting of ridges.
Ridges giving rise to reticulations (net-like markings over the surface).
Reticulations shallow, barely roughening the surface of the spore.
Spores at maturity 15–22 μ in diameter, ridges of reticulations about 1 μ thick.
Apothecia reaching a diameter of 3–5 mm.; reticulations regular. 1. *L. Crouani.*
Apothecia not exceeding 1–2 mm. in diameter; reticulations very irregular. 2. *L. laetirubra.*

Spores at maturity 12–15 μ in diameter, reticulations very delicate, scarcely more than lines over the surface of the spore. 3. *L. dictydiola.*

Reticulations deep, extending 2 μ or more beyond the periphery of the spore and appearing as a broad band about its surface. 4. *L. areolata.*

Ridges not giving rise to reticulations.

Ridges usually curved and extending in various directions as in some species of *Ascobolus*. 5. *L. ascoboloides.*

Ridges giving rise to two distinct rings about the spore. 6. *L. annulata.*

Spore-sculpturing consisting of spines, warts, or tubercles.

Spores covered with spines.

Apothecia not exceeding 1 mm. in diameter; spines short, blunt. 7. *L. spinulosa.*

Apothecia at maturity 1–5 mm. in diameter; spines long and very sharp.

Spores (including spines) 20–25 μ in diameter; spines slender. 8. *L. Crec'hqueraultii*

Spores (including spines) 30 μ or more in diameter; spines stout.

Spines reaching an extreme length of 10 μ, with a diameter of 3 to 4 μ at the base. 9. *L. macrantha.*

Spines conic, about 2–3 μ in diameter and rarely exceeding the same in length. 10. *L. brevispinosa.*

Spore-sculpturing consisting of warts or tubercles.

Tubercles large, twelve or rarely fourteen or fifteen about the circumference of the spore.

Individual tubercles smooth, not sculptured. 11. *L. tuberculata.*

Individual tubercles covered with minute sculpturing, giving them a translucent appearance. 12. *L. Maireana.*

Tubercles small, about twenty or more in the circumference of the spore.

Spores subglobose, on bark among moss. 13. *L. Wrightii.*

Spores perfectly globose, on soil.

Apothecia pale-orange. 14. *L. tuberculatella.*

Apothecia violaceous. 15. *L. amethystina.*

Spores smooth.
Apothecia pale-orange.
About 3 or 4 mm. in diameter at maturity, crowded, on burnt ground. 16. *L. carbonaria.*
About 1 mm. or less in diameter, scattered, on damp soil. 17. *L. haemastigma.*
Apothecia bright-red.
Several mm. in diameter, on damp soil.
Spores 15–20 μ in diameter. 18. *L. Constellatio.*
Spores 10 μ in diameter. 19. *L. wisconsinensis.*
Less than 1 mm. in diameter; spores 8–9 μ in diameter, on foliage of *Sequoia.* 20. *L. gemmea.*
Apothecia pallid or creamy. 21. *L. discoidea.*
Apothecia salmon-colored. 22. *L. salmonicolor.*
Apothecia large, usually 5 mm. or more in diameter; spores often faintly colored.
Apothecia dark-colored, purple or brown to blackish.
Brown or blackish, occurring on burnt ground.
Spores rough.
Apothecia externally rough; spores about 16 μ in diameter.
Spore-sculpturing consisting of rather coarse warts. 23. *L. trachycarpa.*
Spore-sculpturing consisting of minute papillae or pits. 24. *L. Detonia.*
Apothecia externally smooth; spores about 9 μ in diameter. 25. *L. nigrans.*
Spores smooth. 26. *L. leiocarpa.*
Apothecia dark-purple, occurring on damp soil. 27. *L. Planchonis.*
Apothecia bright-colored, orange.
Spores sculptured.
Spore-sculpturing consisting of irregular lumps. 28. *L. lobata.*
Spore-sculpturing consisting of minute warts. 29. *L. exasperata.*
Spores smooth. 30. *L. polytrichina.*

1. **Lamprospora Crouani** (Cooke) Seaver, Mycologia **6**: 8. 1914. (PLATE 3, FIG. 2; 45, FIG. 2, 3.)

Ascobolus miniatus Crouan, Ann. Sci. Nat. IV. **10**: 197. 1858. Not *A. miniatus* Preuss, Linnaea **24**: 147. 1851.
Ascobolus Crouani Cooke, Jour. Bot. **2**: 151. 1864.
Peziza Crouani Cooke, Grevillea **3**: 31. 1874.
Crouania miniata Fuckel, Symb. Myc. 320. 1869.
Lamprospora miniata De-Not. Comm. Critt. Ital. **1**: 388. 1864.
Aleuria Crouani Gill. Champ. Fr. Discom. 50. 1879.

Barlaea miniata Sacc. Syll. Fung. **8**: 111. 1889.
Plicariella miniata Lindau in E. & P. Nat. Pfl. **1**[1]: 180. 1897.
Detonia miniata Rehm; Dodge, Trans. Wisc. Acad. **17**: 1037. 1914.

Apothecia gregarious or crowded, at first globose, expanding and becoming subdiscoid, externally pale-orange with a whitish fringe-like border, reaching a diameter of 3–5 mm., but often not exceeding 1 mm.; hymenium becoming plane or a little concave, bright reddish-orange; excipulum composed of a palisade of parallel or rather loosely interwoven, closely septate, and strongly swollen mycelium, the loose ends of which project above, giving rise to the whitish border, the palisade mycelial threads clavate, reaching a diameter of 10–12 μ, often delicately roughened; asci cylindric or subcylindric above, reaching a length of 350 μ and a diameter of 20–26 μ, tapering below into a pedicel 7–8 μ in diameter; spores 1-seriate, at first smooth, hyaline, containing one oil-drop which nearly fills the spores, finally becoming sculptured and reaching a diameter of 18–22 μ including sculpturing; spore-sculpturing taking the form of reticulations, the meshes of the reticulations ranging from 2–4 or rarely 5–6 μ in diameter, 3–6-sided, often giving rise to almost perfect hexagonal figures, the ridges of the reticulations less than 1 μ thick and shallow as indicated by the projections about the periphery of the spore; paraphyses stout, about 5 or 6 μ in diameter below, enlarged above and reaching a diameter of 9 μ at their apices, filled with minute orange granules.

On damp soil, usually among mosses.

TYPE LOCALITY: Brest, France.

DISTRIBUTION: New York to Colorado and Bermuda; also in Europe and Australia.

ILLUSTRATIONS: Ann. Sci. Nat. IV. **10**: *pl. 13, f. 44–47;* Cooke, Mycographia *pl. 5, f. 17;* Gill. Champ. Fr. Discom. *pl. 52, f. 2;* Mycologia **6**: *pl. 114, f. 2;* Massee, Brit. Fungus-Fl. **4**: 290, *f. 1–4;* Ann. Mag. Nat. Hist. III. **15**: *pl. 16, f. 26;* Phill. Brit. Discom. *pl. 4, f. 19;* Trans. Linn. Soc. **24**: *pl. 51, f. 19;* Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 12, f. 1;* Pat. Tab. Fung. *f. 96;* Proc. Roy. Soc. Victoria II. **38**: *pl. 4, f. 1;* Jour. Bot. **2**: 151, *f. 3.*

A well-known, widely distributed and rather commonly collected species, small but rather conspicuous by reason of its bright color. It was commonly collected in Bermuda during our recent expedition there.

2. **Lamprospora laetirubra** (Cooke) Lagarde, Ann. Myc. **4**: 213 1906.

Peziza laetirubra Cooke, Grevillea **3**: 31. 1874; Mycographia 14. 1875.

Apothecia scattered or thickly gregarious, at first globose or subglobose, expanding and becoming patellate, entirely bright-orange, reaching a diameter of 1–2 mm.; hymenium strongly convex, becoming pitted and rough with age after the spores have been discharged; asci cylindric or subcylindric; spores perfectly globose, containing one large oil-drop which nearly fills the spore, reaching a diameter of 15–18 μ, finally becoming delicately sculptured; spore-sculpturing taking the form of very coarse and irregular reticulations; paraphyses slender, enlarged above, filled with orange granules.

On damp soil in gardens which have been overrun with mosses; also in pastures and meadows.

TYPE LOCALITY: Europe.

DISTRIBUTION: Iowa and Indiana; also in Europe.

ILLUSTRATIONS: Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 12, f. 2* (as *Barlaea cinnabarina* (Fuckel) Sacc.); Ann. Myc. **4**: 213, *f. 33;* Grevillea **3**: *pl. 30, f. 56.*

According to Fuckel *Crouania cinnabarina* with which the present species is thought by some to be synonymous has reticulate spores. Examination of a specimen from Fungi Rhenani No. *2481* shows the spores to be smooth. While this may have been an immature specimen it leaves the identity of Fuckel's plant in doubt. It may be the same as the one here discussed. Our specimens are characterized by the very coarse, loose reticulations. These plants have been frequently collected in Iowa.

3. **Lamprospora dictydiola** Boud. Hist. Class. Discom. Eu. 68. 1907.

Barlaeina dictydiola Sacc. & Trott. in Sacc. Syll. Fung. **22**: 621. 1913.

Apothecia scattered or gregarious but never crowded, at first subglobose, expanding and becoming subdiscoid, reaching a diameter of 1 mm. (in the living plants examined); hymenium becoming plane or nearly so, bordered by a delicate, ragged, fringe-like margin, roughened by the protruding asci, finally becoming more or less pitted apparently from the collapsing of the old asci, orange, a little darker than the outside of the apothecium; asci cylindric or subcylindric, reaching a length of

230–300 μ and a diameter of 18–20 μ; spores 1-seriate, at first smooth and containing one large oil-drop, finally becoming delicately sculptured, reaching a diameter of 12–15 μ, hyaline; spore-sculpturing taking the form of reticulations, the meshes of the reticulations not exceeding 1 μ in diameter, the ridges appearing as delicate lines (when examined with a one-sixth objective); paraphyses nearly straight or a little curved, enlarged above, reaching a diameter of 5 μ at their apices.

On charcoal beds which have been overrun with mosses.

TYPE LOCALITY: France.

DISTRIBUTION: New York; also in Europe.

ILLUSTRATIONS: Boud. Ic. Myc. *pl. 403;* Mycologia **6**: *pl. 114, f. 4.*

4. **Lamprospora areolata** Seaver, Mycologia **4**: 48. 1912. (PLATE 3, FIG. 3.)

Apothecia gregarious but not crowded, at first globose, expanding and becoming subdiscoid, reaching a diameter of 0.5 to 1 mm., orange to bright-red; hymenium becoming plane or slightly convex, roughened by the protruding asci, of about the same color as the outside of the apothecium; asci cylindric or subcylindric, gradually tapering below, reaching a length of 350 μ and a diameter of 15–22 μ; spores 1-seriate, at first smooth and containing one large oil-drop, finally becoming sculptured, reaching a diameter of 18–20 μ, including sculpturing, hyaline; spore-sculpturing taking the form of deep reticulations, the meshes of the reticulations reaching a diameter of 3–5 μ, 3–6-sided, often forming almost perfect hexagonal figures or with the sides of unequal length, 2–3 μ deep as indicated by the projecting ridges about the periphery of the spore, the ridges scarcely 1 μ thick; paraphyses strongly thickened above.

On damp soil among mosses and algae.

TYPE LOCALITY: Yonkers, New York.

DISTRIBUTION: New York and Connecticut.

ILLUSTRATIONS: Mycologia **4**: *pl. 57, f. 5–8;* also **6**: *pl. 114, f. 3.* (Cf. Proc. Roy. Soc. Victoria II. **38**: *pl. 4, f. 2a, b.*)

Lamprospora areolata Seaver, var. *australis* McLennan & Cookson has identical spore characters but differs mainly in the larger size of the apothecia and in the character of the margin.

5. **Lamprospora ascoboloides** Seaver, Mycologia 6: 10. 1914. (PLATE 3, FIG. 7.)

Apothecia gregarious, at first globose and closed, nestling in little depressions in the substratum, gradually expanding and, becoming discoid or subdiscoid, externally slightly floccose, usually not exceeding 1 mm. in diameter, pale-orange; hymenium at first concave, becoming plane and finally convex, roughened by the protruding asci, the entire hymenium collapsing in dried specimens and becoming concave; asci cylindric-clavate, tapering rather abruptly below into a stem-like base, reaching a length of 175–225 μ and a diameter of 18–23 μ; spores at first smooth and containing one or two oil-drops, finally becoming sculptured, reaching a diameter of 15–18 μ, hyaline; spore-sculpturing taking the form of ridge-like markings, the ridges straight or curved and extending in almost any direction, occasionally being parallel or anastomosing and giving rise to a few imperfect reticulations but never perfectly reticulate, about 2 μ thick, resembling those of some species of *Ascobolus;* paraphyses clavate, reaching a diameter of 5–6 μ, at their apices, densely filled with minute granules.

On the ground among mosses in meadows.

TYPE LOCALITY: Portland, Connecticut.

DISTRIBUTION: Connecticut, New York and Virginia.

ILLUSTRATION: Mycologia 6: *pl. 114, f. 8.*

6. **Lamprospora annulata** Seaver, Mycologia 6: 11. 1914. (PLATE 3, FIG. 8.)

Apothecia gregarious but not crowded, at first subglobose and nestling in little cavities in the substratum, expanding and becoming discoid or subdiscoid, pale-orange, reaching a diameter of 0.5–1 mm.; hymenium becoming plane or nearly plane and more or less pitted or roughened, a little darker than the outside of the apothecium; asci cylindric or subcylindric above, rather abruptly attenuated below into a short, much contorted pedicel, reaching a length of 200 μ and a diameter of about 20 μ, at first almost filled with the spores, later stretching and the lower part becoming almost equal in length to the spore-bearing part; spores 1-seriate from the first, smooth, when young containing a number of small oil-drops, finally with one large oil-drop, reaching a diameter of 16–18 μ, hyaline, becoming minutely verrucose and marked about the proximal and distal sides by

two rings; spore-rings at first small, increasing in size until they reach a thickness of 3 or 4 μ, giving the appearance of four circles where the rings pass about the periphery of the spore, the two rings usually parallel with each other but occasionally shifted out of their normal position and assuming a subspiral position, the whole spore at maturity having a short-cylindric appearance, reaching a diameter of 16–18 μ; paraphyses thickened above and filled with granules, reaching a diameter of 5 μ at their apices.

On soil among mosses and algae.

TYPE LOCALITY: Portland, Connecticut.

DISTRIBUTION: New York and Connecticut.

ILLUSTRATION: Mycologia 6: *pl. 114, f. 9.*

This species, which is very distinct in its spore-markings, has been collected several times by the writer in New York and Connecticut but so far as known has never been found by anyone else.

7. **Lamprospora spinulosa** Seaver, Mycologia 6: 11. 1914. (PLATE 3, FIG. 4; 45, FIG. 5, 6.)

Apothecia closely gregarious but not usually crowded, at first subglobose, expanding and becoming short-cylindric or discoid, their bases closely nestling in little cavities in the substratum, reaching a diameter of 1 mm., pale-orange; hymenium plane or nearly so, roughened by the asci which protrude often half their length above its surface, collapsing after discharging their spores and leaving the hymenium pitted, surrounded by a fringe-like border; excipulum composed of a palisade of strongly swollen, rather closely septate mycelial threads which extend above the hymenium in irregular clusters giving rise to the fringe-like border, the ends of the peridial mycelium reaching a diameter of 17–18 μ, minutely granular within and delicately roughened without; asci cylindric above, tapering below into a short, slender, stem-like base, reaching a length of 200 μ and a diameter of 18–20 μ, stretching before discharging their spores and then reaching a length of 300 μ and a diameter of 26 μ; spores at first smooth, finally becoming sculptured, reaching a diameter of 18–20 μ, containing one oil-drop which nearly fills the spore, hyaline; spore-sculpturing taking the form of minute blunt spines 1 μ thick and 2 μ long; paraphyses septate, gradually enlarged above, reaching a diameter of 7–8 μ, minutely granular within and dilutely colored, orange.

On the ground among mosses.

TYPE LOCALITY: The New York Botanical Garden.

DISTRIBUTION: New York City and New Jersey.

ILLUSTRATION: Mycologia **6**: *pl. 114, f. 5.*

8 **Lamprospora Crec'hqueraultii** (Crouan) Boud. Hist. Class. Discom. Eu. 69. 1907. (PLATE 3, FIG. 1; 45, FIG. 8.)

Ascobolus Crec'hqueraultii Crouan, Ann. Sci. Nat. IV. **10**: 194. 1858.
Peziza modesta Karst. Not. Fauna Fl. Fenn. **10**: 122. 1869.
Peziza echinosperma Peck, Ann. Rep. N. Y. State Mus. **24**: 95. 1872.
Peziza auriflava Cooke, Mycographia 16. 1875.
Aleuria auriflava Gill. Champ. Fr. Discom. 50. 1879.
Mollisia Crec'hqueraultii Gill. Champ. Fr. Discom. 118. 1882.
Cromania asperella Rehm, Hedwigia **24**: 226. 1885.
Humaria Crec'hqueraultii Quél. Ench. Fung. 288. 1886.
Barlaea Crec'hqueraultii Sacc. Syll. Fung. **8**: 113. 1889.
Barlaea asperella Sacc. Syll. Fung. **8**: 113. 1889.
Barlaea modesta Sacc. Syll. Fung. **8**: 113. 1889.
Humaria echinosperma Sacc. Syll. Fung. **8**: 130. 1889.
Pilacriella modesta Lindau in E. & P. Nat. Pfl. 1^1: 180. 1897.
Barlaeina Crec'hqueraultii Sacc. & Trott. in Sacc. Syll. Fung. **22**: 621. 1913.

Apothecia gregarious, becoming discoid to scutellate, very pale-orange, externally smooth, reaching a diameter of 2–5 mm.; hymenium becoming plane or more often convex, the margin indistinct, pale-orange fading to a dirty-yellow in dried plants; excipulum composed of large roundish cells below and giving rise to a palisade of mycelium above, the loose ends of which are club-shaped, reaching a diameter of 17–18 μ; asci cylindric or sub-cylindric above, reaching a length of 300–325 μ and a diameter of 27 μ, often protruding above the surface of the hymenium; spores 1-seriate, at first smooth, becoming sculptured, reaching a diameter of 20–25 μ including sculpturing, hyaline; spore-sculpturing taking the form of very sharp spines, the spines broad at the base, reaching a length of 2–3 μ, in dried specimens often bent or adpressed but regaining their normal form when wet; paraphyses rather stout, reaching a diameter of 3–4 μ below, enlarged above and often reaching a diameter of 10 μ at their apices.

On clayey soil.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to Delaware, West Virginia and Colorado; also in Europe and Australia.

ILLUSTRATIONS: Ann. Rep. N. Y. State Mus. **24**: *pl. 3, f.*

10–13; Ann. Sci. Nat. IV. **10**: *pl. 13, f. 12–15;* Boud. Ic. Myc. *pl. 404;* Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 12, f. 4;* Cooke, Mycographia *pl. 6, f. 22, 23;* Mycologia **6**: *pl. 114, f. 1;* Proc. Roy. Soc. Victoria II. **38**: *pl. 5, f. 4.*

EXSICCATI: N. Am. Fungi *840* (as *Peziza echinosperma* Peck); *841* (as *Peziza modesta* Karst.); Clements, Crypt. Form. Colo. *115* (as *Detonia modesta*).

One of the most common and widely distributed species of the genus.

9. **Lamprospora macrantha** (Boud.) Seaver, sp. nov.

Lamprospora Crec'hqueraultii var. *macrantha* Boud. Hist. Class. Discom. Eu. 69. 1907.

Apothecia gregarious but not usually crowded, subglobose, becoming short-cylindric, finally discoid, externally pale-yellow, reaching a diameter of 1–2 mm.; hymenium becoming plane or very slightly concave, with a delicate fringe-like border, pale-orange; excipulum giving rise to a palisade of mycelium above, which forms the fringe about the hymenium, the ends of the palisade mycelium clavate, reaching a diameter of 25–30 μ; asci cylindric above, gradually tapering below into a stem-like base, reaching a length of 300–325 μ and a diameter of 30–35 μ; spores 1-seriate from the first, smooth and densely filled with oil-drops, becoming sculptured apparently by the accumulation of protoplasmic material about the outside of the spore, reaching a diameter of 20 μ excluding spines and 30–35 μ including spines, hyaline; spore-sculpturing at first indistinct, gradually assuming the form of sharp spines which give the spore a beautiful crystalline appearance, the spines reaching a length of 10 μ and a diameter of 3–4 μ at the base; paraphyses strongly enlarged above, reaching a diameter of 7–10 μ at their apices.

On clayey soil in wet places.

TYPE LOCALITY: France.

DISTRIBUTION: New York; also in Europe.

ILLUSTRATION: Boud. Ic. Myc. *pl. 405.*

10. **Lamprospora brevispinosa** Seaver, sp. nov.

Apothecia gregarious, at first globose and closely nestling in little depressions in the substratum, expanding and becoming discoid or subdiscoid, externally orange, reaching a diameter of 1 mm.; hymenium delicately roughened by the protruding asci,

a little brighter than the outside of the apothecium; excipulum composed of a palisade of mycelium, the loose ends of which give rise to a fringe-like border, the ends enlarged, attaining a diameter of 20–25 μ at their extremities; asci broad-cylindric to clavate, tapering rather abruptly below into a stem-like base, reaching a length of 300–325 μ and a diameter of 30–35 μ; spores 1-seriate, at first smooth, densely filled with orange granules, becoming sculptured, reaching a diameter of 25 μ excluding sculpturing or 30–32 μ including sculpturing, hyaline; spore-sculpturing gradually assuming the form of sharp spines, 2–3 μ in diameter at the base and of about the same length, the spines decidedly conic in form; paraphyses stout, containing several large oil-drops, reaching a diameter of 12 μ at their apices.

On moist soil among mosses and algae by roadside.

TYPE LOCALITY: Springside, New Jersey, June 9, 1914.

DISTRIBUTION: Known only from the type locality.

While this and the preceding might be regarded as mere forms of *Lamprospora Crec'hqueraultii*, one of the most commonly collected species of the genus, there is such a wide and constant variation in the size and form of the spines with which the spores are decorated that it seems best to segregate the two latter under the names suggested.

11. **Lamprospora tuberculata** Seaver, Mycologia 4: 47. 1912. (PLATE 3, FIG. 6.)

Apothecia gregarious in small clusters, not crowded but rarely with two or three in close contact, at first globose or subglobose and almost buried in the sandy soil in which they grow, gradually expanding and reaching a diameter of 0.5–1 mm.; superficial but with the base still nestling in the substratum; hymenium plane or nearly so, orange, a little darker than the outside of the apothecium, roughened by the protruding asci which appear as minute white spines; excipulum composed of a palisade of swollen, closely septate mycelium, the loose ends of which give rise to a delicate fringe-like border, the ends of the peridial mycelium about 10 μ in diameter; asci cylindric or subcylindric, tapering below into a stem-like base, reaching a length of 275–300 μ and a diameter of 20–30 μ; spores 1-seriate, at first smooth and usually containing one large oil-drop, gradually becoming sculptured, reaching a diameter of 18–20 μ, hyaline; spore-sculpturing assuming the form of tubercles covering the



LAMPROSPORA

surface of the spore and appearing about its periphery like great lumps, often projecting further in one place than another, giving the spore an irregular outline, the tubercles about 3–4 μ in diameter, 12 or 14, rarely 15, appearing about the periphery of the spore; paraphyses enlarged above, attaining a diameter of 3–4 μ below and a diameter of 6–8 μ at their apices, filled with minute pale-orange granules.

On the soil in open places among mosses and algae.

TYPE LOCALITY: Woods near Yonkers, New York.

DISTRIBUTION: New York to Virginia; also in Australia.

ILLUSTRATIONS: Mycologia **4**: *pl. 57, f. 1–4;* also **6**: *pl. 114, f. 7;* Proc. Roy. Soc. Victoria II. **38**: *pl. 5, f. 1.*

McLennan and Cookson believe that the Tasmanian species described by L. Rodway (Proc. Roy. Soc. Tasmania 1924: 112. 1925) under the name of *Barlaea verrucosa* is identical with our species.

12. **Lamprospora Maireana** Seaver, Mycologia **6**: 14. 1914.

Apothecia gregarious, at first globose, expanding and reaching a diameter of about 2 mm., pale-orange; hymenium plane or slightly concave, of about the same color as the outside of the apothecium; asci cylindric or subcylindric, gradually tapering below into a stem-like base, reaching a length of 300–325 μ and a diameter of 30 μ; spores 1-seriate, becoming sculptured, reaching a diameter of 23 μ, hyaline or subhyaline; spore-sculpturing taking the form of tubercles which are as large as 3–5 μ in diameter, appearing about the periphery of the spore as scallops, the tubercles bearing secondary sculpturing which gives to each tubercle a minutely verrucose appearance and to the whole spore a translucent effect; paraphyses strongly thickened at their apices, reaching a diameter of 8 μ.

On the ground among mosses and algae.

TYPE LOCALITY: Algiers, North Africa.

DISTRIBUTION: New York; also in North Africa and Australia.

ILLUSTRATIONS: Mycologia **6**: *pl. 114, f. 13;* Proc. Roy. Soc. Victoria II. **38**: *pl. 5, f. 3.*

EXSICCATI: Maire, Myc. Bor. Africana *22* (as *Lamprospora tuberculata* Seaver).

13. **Lamprospora Wrightii** (Berk. & Curt.) Seaver, Mycologia **6**: 15. 1914. (PLATE 3, FIG. 9.)

Peziza Wrightii Berk. & Curt.; Berk. & Br. Ann. Mag. Nat. Hist. III. **15**: 444. 1865.

Barlaea Wrightii Sacc. Syll. Fung. **8**: 112. 1889.

Humaria Wrightii Boud. Hist. Class. Discom. Eu. 68. 1907.

Apothecia gregarious or scattered, at first globose, expanding and becoming subdiscoid, reaching a diameter of 1–2 mm., externally pale-yellow and minutely roughened; hymenium plane or slightly concave, surrounded by a delicate fringe-like border, a little darker than the outside of the apothecium; asci cylindric or slightly clavate above, rather abruptly tapering below into a stem-like base, reaching a length of 175–200 μ and a diameter of 17–18 μ; spores usually 1-seriate, or occasionally irregularly crowded, globose or very slightly ellipsoid, at first smooth and containing one to several oil-drops, finally becoming sculptured, reaching a diameter of 15–17 μ, hyaline; spore-sculpturing taking the form of small warts which are usually rather widely scattered over the surface of the spore; paraphyses strongly enlarged above, filled with granules.

On the bark of trees among mosses.

TYPE LOCALITY: Bodelwyddan, Flintshire, Wales.

DISTRIBUTION: Alabama, Texas, Cuba, and reported from Nebraska; also in Europe.

ILLUSTRATIONS: Ann. Mag. Nat. Hist. III. **15**: *pl. 15, f. 16;* Boud. Ic. Myc. *pl. 399;* Cooke, Mycographia *pl. 3, f. 18;* Mycologia **6**: *pl. 114, f. 11.*

14. **Lamprospora tuberculatella** Seaver, Mycologia **6**: 15. 1914. (PLATE 3, FIG. 5; 45, FIG. 9, 10.)

Apothecia gregarious but never crowded, often five or six plants in the space of 1 cm., at first subglobose, expanding and becoming short-cylindric, finally discoid, pale-orange, reaching a diameter of 0.3–0.5 mm. or rarely as large as 1 mm.; hymenium becoming plane or nearly so, a little darker than the outside of the apothecium, roughened by the protruding asci which appear as minute white spines, surrounded by a delicate fringe-like border; asci cylindric or subcylindric above, tapering rather abruptly into a stem-like base, reaching a length of 225–250 μ and a diameter of 25 μ; spores 1-seriate, at first smooth and containing one large oil-drop, reaching a diameter of 18–20 μ, hyaline, finally becoming sculptured; spore-sculpturing taking the

form of tubercles, the number appearing about the periphery of the spore ranging from 20–25 in the circumference of the spore; paraphyses gradually enlarged upwards and filled with orange granules, reaching a diameter of 8–9 μ at their apices.

On soil among mosses.

TYPE LOCALITY: Yonkers, New York.

DISTRIBUTION: New York and Maine; also in Australia.

ILLUSTRATIONS: Mycologia **6**: *pl. 114, f. 6;* Proc. Roy. Soc. Victoria II. **38**: *pl. 5, f. 2.*

15. **Lamprospora amethystina** (Quél.) Seaver, Mycologia **6**: 16. 1914.

Humaria Persoonii amethystina Quél. Assoc. Fr. Av. Sci. Comp. Rendu **14**[2]: 451. 1886.
Barlaea amethystina Sacc. Syll. Fung. **8**: 116. 1889.

Apothecia gregarious, at first subglobose, expanding and becoming discoid, externally pale, whitish or purplish, reaching a diameter of 2 mm.; hymenium becoming plane or a little concave, purplish, a little darker than the outside of the apothecium, bordered by a whitish fringe-like margin; asci cylindric or subcylindric above, gradually tapering below; spores 1-seriate, at first smooth, hyaline, reaching a diameter of 12 μ, becoming sculptured; spore-sculpturing consisting of small warts similar in size and general appearance to those of *L. tuberculatella;* paraphyses slender, slightly enlarged at their apices.

On the ground among mosses.

TYPE LOCALITY: Jura, France.

DISTRIBUTION: Iowa; also in Europe.

ILLUSTRATION: Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 12, f. 3.*

16. **Lamprospora carbonaria** (Fuckel) Seaver, Mycologia **6**: 16. 1914.

Crouania carbonaria Fuckel, Symb. Myc. Nachtr. **2**: 64. 1873.
Peziza sanguinaria Cooke, Grevillea **3**: 31. 1874.
Barlaea carbonaria Sacc. Syll. Fung. **8**: 112. 1889.
?Lamprospora carbonicola Boud. Hist. Class. Discom. Eu. 68. 1907.
Pulvinula carbonaria Boud. Hist. Class. Discom. Eu. 70. 1907.
Barlaeina carbonicola Sacc. & Trav.; Sacc. & Trott. in Sacc. Syll. Fung. **22**: 622. 1913.

Apothecia gregarious or crowded, at first globose, becoming expanded and scutellate to discoid, the margin even or wavy,

pale-orange, reaching a diameter of 1–4 mm.; hymenium becoming plane or slightly concave, roughened by the protruding asci, a little darker than the outside of the apothecium; asci cylindric or subcylindric, reaching a length of 225 μ and a diameter of 18–20 μ; spores smooth, hyaline, reaching a diameter of 15–18 μ, containing one oil-drop which nearly fills the spore; paraphyses filiform or slightly enlarged at their apices, extending far beyond the young asci and strongly curved or hooked, reaching a diameter of 3–4 μ at their apices.

On soil among mosses in places which have been burned but subsequently overgrown with mosses.

TYPE LOCALITY: Oestrich and Budenheim woods, Germany.

DISTRIBUTION: New York and Connecticut to Montana; also in Europe.

ILLUSTRATION: Boud. Ic. Myc. *pl. 401* (as *L. carbonicola*).

17. **Lamprospora haemastigma** (Hedw.) Seaver, Mycologia **6**: 17. 1914. (PLATE 45, FIG. 1.)

?Peziza convexella Karst. Not. Fauna Fl. Fenn. **10**: 123. 1869.
Humaria haemastigma Quél. Ench. Fung. 289. 1886.
Pyronema haemastigma Sacc. Syll. Fung. **8**: 108. 1889. (excl. descr.)
?Octospora haemastigma Hedw. Descr. **2**: 17. 1788.
?Barlaea convexella Sacc. Syll. Fung. **8**: 114. 1889.
Pulvinula haemastigma Boud. Hist. Class. Discom. Eu. 70. 1907.
Detonia convexella Dodge, Trans. Wisc. Acad. **17**: 1037. 1914.

Apothecia rather thickly gregarious but not usually crowded, at first subglobose, expanding and becoming discoid, pale-yellow to red, becoming brighter in dried specimens, reaching a diameter of 1 mm.; hymenium plane or slightly concave, similar in color to the outside of the apothecium; asci cylindric or subcylindric above, tapering below into a stem-like base, reaching a length of 300 μ and a diameter of 20–23 μ; spores 1-seriate, smooth, usually containing one to several oil-drops, reaching a diameter of 20 μ, hyaline; paraphyses very slender, straight or often strongly curved at their apices, scarcely thickened above, reaching a diameter of 2 μ, densely filled with yellow granules.

On damp soil among mosses.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to Wisconsin; also in Europe.

ILLUSTRATIONS: Hedw. Descr. **2**: *pl. 5, f. 1–5;* Boud. Ic. Myc. *pl. 406*.

18. **Lamprospora Constellatio** (Berk. & Br.) Seaver, Mycologia **6**: 18. 1914. (PLATE 45, FIG. 7.)

?Crouania humosa Fuckel, Symb. Myc. 320. 1869. (excl. syn.)
Peziza Constellatio Berk. & Br. Ann. Mag. Nat. Hist. IV. **17**: 142. 1876.
Leucoloma Constellatio Rehm, Ber. Naturh. Ver. Augsburg **26**: 5. 1881.
Pulvinula Constellatio Boud. Bull. Soc. Myc. Fr. **1**: 107. 1885.
Aleuria Constellatio Gill. Champ. Fr. Discom. 207. 1888.
Barlaea Constellatio minuta Clements, Bot. Surv. Nebr. **4**: 10. 1896.
Plicariella Constellatio Lindau in E. & P. Nat. Pfl. 1^1: 180. 1897.
Barlaeina Constellatio minuta Sacc. & D. Sacc. in Sacc. Syll. Fung. **14**: 749. 1899.
Barlaeina Constellatio Sacc. & Sydow in Sacc. Syll. Fung. **14**: 749. 1899.
Detonia Constellatio Rehm; Dodge, Trans. Wisc. Acad. **17**: 1037. 1914.

Apothecia gregarious or scattered, at first globose, expanding and becoming scutellate to discoid, pale-orange, reaching a diameter of 1–5 mm.; hymenium bright-red, the color becoming brighter in dried specimens, often almost scarlet; asci cylindric or subcylindric above, tapering into a long stem-like base, reaching a length of 250–300 μ and a diameter of 20 μ; spores 1-seriate, smooth, hyaline, reaching a diameter of 15–20 μ, containing one large oil-drop which is often surrounded by numerous smaller ones; paraphyses filiform, only slightly thickened at their apices and very much curved and hooked, septate, filled with red granules.

On bare ground, especially on rich soil in woods or meadows.

TYPE LOCALITY: Addington, Kent, Great Britain.

DISTRIBUTION: New Jersey to Ontario, Colorado, California and Jamaica; also in Europe.

ILLUSTRATIONS: Boud. Ic. Myc. *pl. 407;* Cooke, Mycographia *pl. 21, f. 81;* Pat. Tab. Fung. *f. 372.*

EXSICCATI: N. Am. Fungi *2036.*

19. **Lamprospora wisconsinensis** Seaver, nom. nov.

Barlaea laeterubra Rehm, Ann. Myc. **3**: 516. 1905. Not *Peziza laetirubra* Cooke.
Barlaeina laeterubra Sacc. & Trott. in Sacc. Syll. Fung. **22**: 622. 1913.
Detonia laeterubra Dodge, Trans. Wisc. Acad. **17**: 1037. 1914.

Apothecia scattered, at first globose or subglobose, expanding and becoming discoid or subdiscoid, pale-yellow, scarcely exceeding 1 mm. in diameter; hymenium plane or nearly so, similar in color to the outside of the apothecium; asci cylindric or subcylindric above, tapering below into a stem-like base, reaching a length of 150–175 μ and a diameter of 12–14 μ; spores at

first partially 2-seriate or irregularly crowded, finally becoming 1-seriate, smooth, granular within but containing no conspicuous oil-drops, hyaline, never exceeding 10 μ in diameter and often smaller; paraphyses filiform, 1.5–2 μ in diameter, slightly thickened above and strongly curved at their apices, sparingly branched, filled with hyaline granules.

On the ground and on burnt ground which has been overrun with mosses.

TYPE LOCALITY: Cemetery woods, Wisconsin.

DISTRIBUTION: New York and Wisconsin.

20. **Lamprospora gemmea** (Phill.) Seaver, Mycologia **6**: 18. 1914.

Peziza gemmea Phill. Grevillea **7**: 21. 1878.
Barlaea gemmea Sacc. Syll. Fung. **8**: 112. 1889.

Apothecia gregarious, sessile, subturbinate, finally expanding and becoming more or less discoid, externally pale-reddish, reaching a diameter of 0.5 mm.; hymenium plane or slightly concave, bright-red; asci cylindric or subcylindric, gradually tapering below; spores 1-seriate, smooth, hyaline, reaching a diameter of 8 μ, granular within; paraphyses filiform, very slender, scarcely enlarged above, branched, straight or more or less curved at their apices, not exceeding 2 μ in diameter.

On decaying foliage of *Sequoia sempervirens*.

TYPE LOCALITY: California.

DISTRIBUTION: Known only from the type locality.

ILLUSTRATION: Cooke, Mycographia *pl. 111, f. 398*.

21. **Lamprospora discoidea** (P. Henn. & E. Nym.) Seaver, Mycologia **6**: 19. 1914. (PLATE 45, FIG. 4.)

Barlaea discoidea P. Henn. & E. Nym. Monsunia **1**: 33. 1900.
Barlaeina discoidea Sacc. & Sydow in Sacc. Syll. Fung. **16**: 710. 1902.

Apothecia scattered, gregarious or occasionally closely congested, sessile, at first subglobose, finally expanding and becoming discoid, pale-whitish, reaching a diameter of 1–2 mm.; hymenium plane or convex, with the margin slightly elevated, roughened by the protruding asci, whitish or grayish with a tinge of yellow or cream; excipulum composed of a palisade of mycelium, the loose ends of which are clavate; asci subcylindric above, tapering below into a slender stem-like base which is usually forked, reaching a length of 200–250 μ and a diameter of 20 μ; spores

1-seriate, smooth, rather thick-walled, reaching a diameter of 15–20 μ, hyaline; paraphyses slender below, reaching a diameter of 3 μ, enlarged upwards to 5 μ, occasionally branched, straight or strongly curved or hooked at their apices.

On bare sandy soil or among mosses.

TYPE LOCALITY: Java.

DISTRIBUTION: New York; also in Europe.

22. **Lamprospora salmonicolor** Seaver, Mycologia **17**: 47. 1925.

Apothecia discoid, with the hymenium plane or slightly convex, reaching a diameter of 2 mm., pale salmon-colored; asci cylindric or subcylindric, reaching a length of 275 μ and a diameter of 20–24 μ; spores globose, filled with minute granules, smooth, 20 μ in diameter when mature; paraphyses clavate.

On bare ground.

TYPE LOCALITY: El Yunque, Porto Rico.

DISTRIBUTION: Known only from the type locality.

23. **Lamprospora trachycarpa** (Curr.) Seaver, Mycologia **6**: 19. 1914.

Peziza trachycarpa Curr. Trans. Linn. Soc. **24**: 493. 1864.
?Bulgaria carbonaria Fuckel, Fungi Rhen. *1137*. 1865.
?Peziza atrospora Fuckel, Fungi Rhen. *1224*. 1865.
?Plicaria carbonaria Fuckel, Symb. Myc. 326. 1869.
?Plicaria ferruginea Fuckel, Symb. Myc. 326. 1869.
Peziza scabrosa Cooke, Mycographia 170. 1877.
Discina trachycarpa Karst. Act. Fauna Fl. Fenn. **2**: 113. 1885.
Aleuria trachycarpa Gill. Champ. Fr. Discom. 207. 1886.
Detonia trachycarpa Sacc. Syll. Fung. **8**: 105. 1889.
Phaeopezia scabrosa Sacc. Syll. Fung. **8**: 472. 1889.
?Phaeopezia atrospora Sacc. Syll. Fung. **8**: 472. 1889.
Curreyella trachycarpa Massee, Brit. Fungus-Fl. **4**: 401. 1895.
Plicariella trachycarpa Rehm in Rab. Krypt.-Fl. 1^3: 996. 1896.

Apothecia gregarious or densely crowded, often forming continuous masses extending over a space of several cm., at first subglobose, expanding and becoming shallow cup-shaped with the margin elevated and incurved, or more rarely scutellate and closely adhering to the substratum, regular in form or cochleate and becoming very irregular from mutual pressure, externally paler than the hymenium and usually rough and scabrous, reaching a diameter of from a few mm. to 2 cm.; hymenium smooth or convolute, dark reddish-brown or slightly olivaceous, becoming black when old or in dried specimens; asci cylindric or

subcylindric above, reaching a length of 250–300 μ and a diameter of 14–16 μ or rarely 18 μ; spores 1-seriate, at first smooth, becoming sculptured, reaching a diameter of 15–18 μ, becoming pale-yellow or smoky; spore-sculpturing taking the form of minute warts or short, interrupted ridges; paraphyses about 3–4 μ in diameter below, sparingly branched, enlarged at their apices where they reach a diameter of 8 μ, adhering together, yellowish-brown.

On burnt ground and charcoal beds.

TYPE LOCALITY: Ascot Heath, Great Britain.

DISTRIBUTION: New Hampshire to Colorado; also in Europe.

ILLUSTRATIONS: Boud. Ic. Myc. *pl. 306;* Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 14, f. 1;* Cooke, Mycographia *pl. 67, f. 257;* Trans. Linn. Soc. **24**: *pl. 51, f. 3, 5;* Mycologia **6**: *pl. 114, f. 10.*

One of the larger species of the genus and rather commonly collected on burnt places.

24. **Lamprospora Detonia** Brenckle, Mycologia **8**: 318. 1916.

Apothecia scattered, scutellate, with the margins elevated, slightly incurved and free from the substratum, externally brown and verrucose, quite regular in form, becoming convolute in drying, reaching a diameter of 5–12 mm.; hymenium smooth, concave to plane, dark-brown to black with a green tint; asci cylindric or subcylindric, reaching a length of 300 μ and a diameter of 15 μ; spores 1-seriate, crowded in the upper third of the ascus, at first smooth, becoming sculptured, containing one large oil-drop, becoming brown and opaque so as to appear black under the microscope, reaching a diameter of 12–15 μ; spore-sculpturing consisting of minute warts or pits which are much finer than those of *L. trachycarpa;* paraphyses slender, yellowish, the apex thickened and colored.

Among mosses on shaded ground in woods.

TYPE LOCALITY: Cheyenne River near Anselm, North Dakota.

DISTRIBUTION: Known only from the type locality.

25. **Lamprospora nigrans** (Morgan) Seaver, Mycologia **6**: 20. 1914.

Peziza nigrans Morgan, Jour. Cinn. Soc. Nat. Hist. **18**: 43. 1895.
Detonia nigrans Sacc. & Sydow in Sacc. Syll. Fung. **14**: 747. 1899.

Apothecia at first roundish, expanding and becoming cup-shaped, finally scutellate and often more or less irregular, ex-

ternally smoky-pallid and smooth, attached to the soil by slender fibers, reaching a diameter of 1–2 cm.; hymenium plane or concave, becoming black or blackish; asci cylindric or subcylindric, reaching a length of 150 μ and a diameter of 10 μ; spores 1-seriate, at first smooth, becoming sculptured and smoky-brown, reaching a diameter of 9 μ; spore-sculpturing consisting of rather coarse warts, the warts rounded, rather conspicuous and about 2 μ in diameter; paraphyses slender, thickened above, yellowish-brown.

On burnt ground.

TYPE LOCALITY: Preston, Ohio.

DISTRIBUTION: Known only from the type locality.

ILLUSTRATION: Jour. Cinn. Soc. Nat. Hist. **18**: *pl. 3, f. 17.*

26. **Lamprospora leiocarpa** (Curr.) Seaver, Mycologia **6**: 21. 1914.

Peziza leiocarpa Curr. Trans. Linn. Soc. **24**: 493. 1864.
Plicaria foveata Fuckel, Symb. Myc. 326. 1869.
Detonia leiocarpa Sacc. Syll. Fung. **8**: 105. 1889.
Detonia foveata Sacc. Syll. Fung. **8**: 105. 1889.
Aleuria leiocarpa Gill. Champ. Fr. Discom. (suites). 1890.
Curreyella foveata Massee, Brit. Fung.-Fl. **4**: 402. 1895.
Plicariella leiocarpa Rehm in Rab. Krypt.-Fl. 1^3: 994. 1896.

Apothecia gregarious, scattered or crowded, at first globose and closed, expanding and becoming shallow cup-shaped, regular in form or cochleate and occasionally incised and *Otidea*-like, at length scutellate or subdiscoid, externally reddish-brown to brownish-black with a slight tinge of olive, often strongly roughened, reaching a diameter of 2–3 cm.; hymenium smooth and even or undulated and lobed, similar in color to the outside of the apothecium but a little darker; asci cylindric or subcylindric above, tapering below into a stem-like base, reaching a length of 350 μ and a diameter of 15–18 μ; spores at first irregularly 2-seriate, finally separating and becoming 1-seriate, hyaline or slightly yellowish, rather thick-walled, smooth, reaching a diameter of 10–12 μ, usually containing one large central oil-drop; paraphyses slender, rather abruptly enlarged above, densely granular within, reaching a diameter of 7 μ, showing a tendency to adhere together in fascicles.

On burnt ground.

TYPE LOCALITY: Ascot Heath, Great Britain.

DISTRIBUTION: California, Washington, Idaho, Montana; also in Europe.

ILLUSTRATIONS: Trans. Linn. Soc. **24**: *pl. 51, f. 4, 6;* Boud. Ic. Myc. *pl. 304;* Gill. Champ. Fr. Discom. (suites).

27. **Lamprospora Planchonis** (Dun.) Seaver, Mycologia **6**: 21. 1914.

Plicaria Planchonis Dun.; Boud. Bull. Soc. Myc. Fr. **3**: 92. 1887.

Apothecia gregarious or scattered, sessile, concave-hemispheric or nearly discoid, regular or irregularly contorted, the margin crenate or even, externally minutely warted, dark-purple or almost black, reaching a diameter of 5–8 mm.; hymenium concave or nearly plane, a little paler than the outside of the apothecium; substance bright-purple with transmitted light, the color being easily extracted from the dried plants with water; asci cylindric or subcylindric, tapering below into a slender stem, reaching a length of 200 μ and a diameter of 13–15 μ; spores 1-seriate, at first hyaline, becoming pale-purplish, smooth or very minutely sculptured at maturity, containing one to several oil-drops, reaching a diameter of 10–12 μ; spore-sculpturing consisting of very minute warts which barely roughen the surface of the spore; paraphyses clavate, reaching a diameter of 6 μ at their apices, filled with purple granules.

On sandy soil by roadsides, hillsides and on sand-dunes.

TYPE LOCALITY: France.

DISTRIBUTION: Very common in the Bermudas but not known from the mainland of North America; also in Europe.

ILLUSTRATIONS: Bull. Soc. Myc. Fr. **3**: *pl. 8;* Boud. Ic. Myc. *pl. 309.*

28. **Lamprospora lobata** (Berk. & Curt.) Seaver, Mycologia **6**: 22. 1914. (PLATE 3, FIG. 10.)

Peziza lobata Berk. & Curt.; Berk. Jour. Linn. Soc. **10**: 365. 1868.
Barlaea lobata Sacc. Syll. Fung. **8**: 117. 1889.

Apothecia scattered, at first subglobose, becoming shallow cup-shaped, finally almost scutellate, with the margin undulated or lobed, pale-yellow, reaching a diameter of 5–12 mm.; hymenium concave or nearly plane, a little darker than the outside of the apothecium; asci cylindric or subcylindric, reaching a length of 250 μ and a diameter of 15–18 μ; spores 1-seriate, at first smooth, becoming sculptured, reaching a diameter of 12–15 μ,

hyaline; spore-sculpturing consisting of four to six tubercles of variable size which extend beyond the periphery of the spore and several indistinct lines or bands extending across the spore and appearing like pieces of coarse twine wound about its surface, the inequality of the tubercles giving to the spore a rather irregular form; paraphyses very slender, reaching a diameter of 1–2 μ and scarcely enlarged at their apices.

On the ground.

TYPE LOCALITY: Cuba.

DISTRIBUTION: Known only from the type locality.

ILLUSTRATIONS: Cooke, Mycographia *pl. 69, f. 265;* Mycologia **6**: *pl. 114, f. 12.*

29. **Lamprospora exasperata** (Berk. & Curt.) Seaver, comb. nov.

Peziza exasperata Berk. & Curt. Grevillea **3**: 152. 1874.
Barlaea exasperata Sacc. Syll. Fung. **8**: 112. 1889.

Apothecia at first globose or subglobose, expanding and becoming shallow cup-shaped, with the margin inflexed, externally minutely verrucose, reaching a diameter of a little more than 1 cm.; hymenium concave, red; asci cylindric or subcylindric; spores 1-seriate, globose, becoming sculptured, reaching a diameter of 12–14 μ; spore-sculpturing consisting of minute warts; paraphyses rather stout.

On burnt ground.

TYPE LOCALITY: Alabama.

DISTRIBUTION: Known only from the type locality.

ILLUSTRATION: Cooke, Mycographia *pl. 6, f. 21.*

30. **Lamprospora polytrichina** (Rehm) Seaver, Mycologia **6**: 23. 1914.

?Peziza Polytrichi Schum. Enum. Pl. Saell. **2**: 423. 1803.
Peziza leucoloma Karst. Not. Fauna Fl. Fenn. **10**: 122. 1869. (excl. syn.)
Barlaea Polytrichi Sacc. Syll. Fung. **8**: 113 (in part). 1889.
Detonia polytrichina Rehm in Rab. Krypt.-Fl. 1^3: 1269. 1896.
Plicariella Polytrichi Lindau in E. & P. Nat. Pfl. 1^1: 180. 1897.

Apothecia gregarious, sessile, at first subglobose, becoming scutellate or shallow cup-shaped, margin often wavy, externally paler than the hymenium, whitish and more or less pruinose, reaching a diameter of 5–10 mm.; hymenium concave or nearly plane, bright-orange; asci cylindric or subcylindric, gradually tapering toward the base, reaching a length of 200–225 μ and a diameter of 17 μ; spores 1-seriate, smooth, reaching a diameter

of 10–13 μ, hyaline, often containing one or several oil-drops; paraphyses slender, slightly enlarged above, straight, slightly curved or more rarely hooked.

On bare soil among mosses, especially *Polytrichum*.

TYPE LOCALITY: Europe.

DISTRIBUTION: Minnesota and (Washington?); also in Europe.

ILLUSTRATION: Pat. Tab. Fung. *f. 277*.

DOUBTFUL AND EXCLUDED SPECIES

Peziza globifera Berk. & Curt.; Berk. Jour. Linn. Soc. **10**: 366. 1868. A Cuban species with pale-yellow apothecia and smooth spores, the spores reaching a diameter of 12–14 μ. A microscopic examination of the spores of this species has been made, but our knowledge is too meager to permit of a more complete description.

Barlaea fuliginea (Schum.) Sacc. Syll. Fung. **8**: 117. 1889. *Peziza fuliginea* Schum. Enum. Pl. Saell. **2**: 427. 1803. This species is reported from North America by Saccardo. No specimen of the species has been seen and the identity of the species is uncertain.

6. **PITHYA** Fuckel, Symb. Myc. 317. 1869.

Plants saprophytic or semiparasitic on the foliage and bark of coniferous trees; apothecia bright-colored, whitish externally, stipitate or sessile, the substance rather tough, not shrinking much in drying; hymenium bright-orange or yellow, even or lacunose, plane, concave or convex; asci 8-spored; spores perfectly globose, smooth, hyaline; paraphyses filiform or slightly enlarged above.

Type species, *Peziza pithya* Pers.

Plants occurring on the bark or foliage of *Abies*.	
Hymenium smooth.	1. *P. pithya*.
Hymenium lacunose.	2. *P. lacunosa*.
On foliage of *Juniperus*, *Cupressus*, etc.	3. *P. Cupressi*.

1. **Pithya pithya** (Pers.) Gill. Champ. Fr. Discom. 225 (as synonym). 1887. (PLATE 4, FIG. 3.)

Peziza pithya Pers. Ic. Descr. 43. 1800.
Pithya vulgaris Fuckel, Symb. Myc. 317. 1869.
Peziza Leineri Gonnerm. & Rab. Myc. Eu. **1**: 9. 1869.
Helotium pithyum Gill. Champ. Fr. Discom. 155. 1883.
Humaria pityina Quél. Ench. Fung. 289. 1886.
Peziza pithyna Pat. Tab. Fung. **7**: 73. 1889.

Apothecia gregarious, sessile or short-stipitate, at first subglobose, expanding and becoming subdiscoid, regular in form

when young, becoming irregular with age, externally pale, whitish or yellowish-white, surrounded at the base by a growth of white mycelium, margin of the apothecium free or adnate to the substratum, reaching a diameter of 12 mm.; hymenium plane or convex, bright-orange; asci cylindric or subcylindric, gradually attenuated below, reaching a length of 300–325 μ and a diameter of 15 μ; spores 1-seriate, at first granular within, later containing one large oil-drop, reaching a diameter of 12–14 μ; paraphyses slightly enlarged above, reaching a diameter of 3–4 μ at their apices.

On foliage and bark of species of *Abies*, *Sequoia*, and *Libocedrus*.

TYPE LOCALITY: Germany.

DISTRIBUTION: California; also in Europe.

ILLUSTRATIONS: Boud. Ic. Myc. *pl. 320;* Gonnerm. & Rab. Myc. Eu. 1: 5, *f. 4;* Pers. Ic. Descr. *pl. 11, f. 2;* Pat. Tab. Fung. *f. 698.*

While this species has been recorded from North America, most of the specimens examined belong with *P. Cupressi* rather than here. The present form is larger and occurs on bark more often than on foliage. The species is less frequent in America than in Europe, so far as we can judge from the specimens examined.

2. **Pithya lacunosa** (Ellis & Ev.) Seaver, comb. nov. (PLATE 4, FIG. 2.)

Barlaea lacunosa Ellis & Ev. Proc. Acad. Nat. Sci. Phila. **1894**: 347.

Apothecia sessile, spreading over the substratum and closely adhering to it, except the extreme margin which is free and slightly elevated, circular or more or less irregular in outline, reaching a diameter of 5 mm. to 1.5 cm., externally yellowish-white and attached to the substratum by a white mycelium; hymenium uneven, undulated or deeply lacunose, bright-orange; asci cylindric above, tapering below into a long stem-like base, reaching a length of 200–225 μ and a diameter of 12–17 μ; spores 1-seriate, smooth, reaching a diameter of 12–14 μ, granular within, hyaline; paraphyses gradually thickened above, reaching a diameter of 3–4 μ at their apices.

On the bark of some species of *Abies balsamea.*

TYPE LOCALITY: Notre Dame Bay, Newfoundland.

DISTRIBUTION: Known only from the type locality.

Rather closely related to the preceding species but differing in its lacunose hymenium.

3. **Pithya Cupressi** (Batsch) Rehm in Rab. Krypt.-Fl. 1^3: 926. 1896. (PLATE 4, FIG. 1.)

Peziza Cupressi Batsch, Elench. Fung. **1**: 119. 1783.
Peziza cupressina Fries, Syst. Myc. **2**: 135. 1822.
Pithya cupressina Fuckel, Symb. Myc. 317. 1869.
Humaria cupressina Quél. Ench. Fung. 289. 1886.
Helotium thujinum Peck, Ann. Rep. N. Y. State Mus. **26**: 82. 1874.
Phialea cupressina Gill. Champ. Fr. Discom. 107. 1882.
Lachnella Cupressi Phill. Brit. Discom. 240. 1887.
Pithya thujina Sacc. Syll. Fung. **8**: 210. 1889.

Apothecia gregarious, sessile or short-stipitate, at first subglobose, expanding, becoming circular in form or elongated and irregular; hymenium plane or slightly concave, deep-orange, darker than the exterior of the apothecium; the cells of the excipulum vertically elongated and subhyaline, giving rise below to white mycelial threads by means of which the apothecia are attached to the substratum; asci cylindric or subcylindric above, tapering below into a long stem-like base, reaching a length of 250 μ and a diameter of 15 μ; spores 1-seriate, granular within and usually containing one large oil-drop, reaching a diameter of 10–12 μ; paraphyses branched below, only slightly enlarged above, reaching a diameter of 2–3 μ at their apices.

On recently killed foliage of various species of *Juniperus*, *Cupressus*, *Thuja* and *Sequoia*.

TYPE LOCALITY: Europe.

DISTRIBUTION: Maine to Washington, New Jersey and California, and in the Bermuda Islands; also in Europe.

ILLUSTRATIONS: Grevillea **3**: *pl. 40, f. 171;* Gonnerm. & Rab. Myc. Eu. **1**: *pl. 5, f. 9;* E. & P. Nat. Pfl. 1^1: 205, *f. 160, C–D;* Phill. Brit. Discom. *pl. 8, f. 45.*

EXSICCATI: N. Am. Fungi *2322;* Fungi Columb. *741;* Ravenel, Fungi Am. *629* (as *Peziza pithya* Pers.); Thüm. Mycoth. Univ. *718* (as *Pithya cupressina* Fuckel).

Commonly collected in Bermuda on foliage of Bermuda cedar. Also fairly common in continental North America.

EXCLUDED SPECIES

Pithya suecia (DeBary) Fuckel, Symb. Myc. Nacht. **3**: 32. 1875. According to Rehm, this is a synonym of *Lachnellula chrysophthalma* (Pers.) Karst., a species which occurs in North America. The species belongs with the inoperculate discomycetes and is not a true *Pithya*.



1. PITHYA CUPRESSI
2. PITHYA LACUNOSA
3. PITHYA PITHYA

7. **ASCODESMIS** Van Tiegh. Bull. Soc. Bot. Fr. **23**: 271. 1876.

Ascolobus § *Sphaeridiobolus* Boud. Bull. Soc. Myc. Fr. **14**: 126. 1898.
Sphaeridiobolus Boud. Hist. Class. Discom. Eu. 73. 1907.

Apothecia minute, less than 1 mm. in diameter, very simple, consisting of a cluster of asci and paraphyses springing from a clump of basal mycelium; excipulum almost entirely undeveloped; asci comparatively few to each plant, very broad, 8-spored; spores globose or subglobose, hyaline, then pale-brown to blackish, becoming sculptured at maturity; spore-sculpturing consisting of warts or ridges; paraphyses sparse.

Type species, *Ascodesmis nigricans* Van Tiegh.

Spores globose, reticulated. 1. *A. microscopica.*
Spores subglobose, rough but not reticulated. 2. *A. porcina.*

1. **Ascodesmis microscopica** (Crouan) Seaver, Mycologia **8**: 3. 1916. (PLATE 5, FIG. 1–5.)

Ascobolus microscopicus Crouan, Ann. Sci. Nat. IV. **7**: 175. 1857.
?Ascobolus caninus Fuckel, Hedwigia **5**: 3. 1866.
Ascobolus hyperboreus Karst. Not. Fauna Fl. Fenn. **11**: 204. 1870.
Ascodesmis nigricans Van Tiegh. Bull. Soc. Bot. Fr. **23**: 275. 1876.
Boudiera microscopica Cooke, Grevillea **6**: 76. 1877.
Ascobolus niveus Quél. Fr. Assoc. Av. Sci. Comp. Rendu **9**: 674. 1880.
Sphaeridiobolus hyperboreus Heimerl, Niederosterr. Ascoboleen 12. 1889.
?Boudiera canina Schröt. Krypt.-Fl. Schles. **3²**: 55. 1893.
Boudiera Claussenii P. Henn. Hedwigia **42**: (182). 1903.
?Ascodesmis reticulata Bainier, Bull. Soc. Myc. Fr. **23**: 137. 1907.
Sphaeridiobolus hyperboreus niveus Boud. Hist. Class. Discom. Eu. 73. 1907.

Apothecia minute, less than 1 mm. in diameter, scattered or thickly gregarious, superficial, consisting of a tuft of asci and very stout paraphyses, at first entirely white, becoming dotted over with black specks, the ends of the asci filled with colored spores, finally becoming entirely black; excipulum almost wanting or consisting of loose mycelium similar to the paraphyses; asci broad-clavate to ovoid, reaching a length of 80 μ and a diameter of 30 μ, 8-spored; spores 2-seriate or irregularly crowded together, perfectly globose, becoming pale-brown and sculptured, reaching a diameter of 10–12 μ; spore-sculpturing consisting of net-like reticulations, the meshes reaching a diameter of 3 μ, the ridges of the reticulations thin and projecting about the periphery of the spore as minute spines; paraphyses hyaline, stout, scarcely enlarged above, reaching a diameter of 5–6 μ.

On excrement of dogs and tigers, less frequently on that of other animals.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York; also in Europe.

ILLUSTRATIONS: Ann. Sci. Nat. IV. **7**: *pl. 4, f. 20–23;* Bull. Soc. Myc. Fr. **23**: *pl. 19, f. 1–3;* Hedwigia **42**: (182), *f. 1–3;* Massee, Brit. Fungus-Fl. **4**: 12, *f. 46;* Mycologia **8**: *pl. 172, f. 1–5;* Fr. Assoc. Av. Sci. Comp. Rendu **9**: *pl. 9, f. 18;* Zukal, Mycol. Unters. *pl. 2, f. 5–10.*

2. **Ascodesmis porcina** Seaver, Mycologia **8**: 3. 1916. (PLATE 5, FIG. 6–9.)

Apothecia very small, scarcely visible, scattered or thickly gregarious, at first subglobose, expanding and becoming subdiscoid, externally whitish or pallid, not exceeding 0.5 mm. in diameter; hymenium strongly convex, becoming dark, finally almost black by reason of the dark-colored spores; excipulum almost wanting; asci broad-clavate to ovoid, abruptly narrowed into a stem-like base, reaching a length of 75–90 μ and a diameter of 25–35 μ; spores 2-seriate or very irregularly bunched near the end of the ascus, subglobose, at first hyaline and smooth, becoming pale-brown sculptured, reaching a diameter of 10–13 μ, or occasionally 10–11 $\times$ 12–13 μ; spore-sculpturing very variable, consisting of minute rounded wart-like projections, short interrupted ridges, or often with one conspicuous simple or branched ridge extending across the visible surface of the spore; paraphyses stout, gradually enlarged at their apices, reaching a diameter of about 5 μ.

On excrement of pigs.

TYPE LOCALITY: Grown in the laboratories of The New York Botanical Garden on pig dung sent from Porto Rico.

DISTRIBUTION: Cultivated on pig dung from Porto Rico and New Jersey.

ILLUSTRATION: Mycologia **8**: *pl. 172, f. 6–9.*

The occurrence of this species on material collected in two such widely separated localities would indicate that the species is more dependent upon the substratum conditions than on climate, assuming that the material collected in Porto Rico was infected before it was received.

8. CUBONIA Sacc. Syll. Fung. **8**: 527. 1889.

Apothecia sessile, convex-hemispheric, naked or clothed with hairs, white or whitish; hymenium plane or convex, roughened



1–5. ASCODESMIS MICROSCOPICA
6–9. ASCODESMIS PORCINA

by the protruding asci; asci clavate or subpyriform, 8-spored; spores perfectly globose, minutely sculptured, hyaline.

Type species, *Lasiobolus brachyascus* March.

One unnamed species of this genus has been reported for North America by Dr. B. O. Dodge. This genus contains but three species and in the absence of suitable material it is impossible to determine the identity of the American species. All of the species are coprophilous.

9. **ASCOBOLUS** Pers.; J. F. Gmel. Syst. Nat. 1461. 1791.

Apothecia sessile or substipitate, superficial or partially immersed in the substratum, externally smooth or pilose; hymenium concave, plane or convex; substance soft, fleshy or waxy, usually greenish; asci cylindric to clavate or subovoid, 4–8-spored; spores becoming blue or purple, fading to brown or blackish, ellipsoid to subglobose, smooth or becoming sculptured; spore-sculpturing very variable, consisting of warts, ridges or crevices; paraphyses slender and usually adhering together.

Type species, *Peziza stercoraria* Bull.

Plants growing on the dung of animals.
- Apothecia medium or small, not exceeding 5 mm. in diameter.
  - Apothecia medium large, reaching 5 mm. in diameter. — 1. *A. stercorarius.*
  - Apothecia not exceeding 1–2 mm. in diameter.
    - Spores very large, 20–35 × 50–70 μ. — 2. *A. immersus.*
    - Spores medium large, 25–40 μ long.
      - Spores densely marked with ridges or crevices. — 3. *A. glaber.*
      - Spores very minutely roughened.
        - Spores becoming delicately reticulated. — 4. *A. Winteri.*
        - Spores not reticulated.
          - Spores 15–17 × 30–40 μ. — 5. *A. americanus.*
          - Spores 13–15 × 24–28 μ. — 6. *A. Leveillei.*
    - Spores small, 7–8 × 12–14 μ.
      - Apothecia greenish. — 7. *A. viridulus.*
      - Apothecia white. — 8. *A. candidus.*
- Apothecia large, 5 mm. to 2.7 cm. in diameter. — 9. *A. magnificus.*

Plants growing on soil.
- Restricted to burnt places. — 10. *A. carbonarius.*
- On damp soil, not on burnt places.
  - Apothecia brownish-black.
    - Spores ellipsoid. — 11. *A. geophilus.*
    - Spores subglobose. — 12. *A. subglobosus.*
  - Apothecia white. — 13. *A. albinus.*

Plants growing on wood.	14. *A. xylophilus.*
Plants growing on dead leaves.	15. *A. striisporus.*
Plants growing on fungi or slime-moulds.	16. *A. epimyces.*

1. **Ascobolus stercorarius** (Bull.) Schröt. Krypt.-Fl. Schles. 3^2: 56. 1893. (PLATE 45, FIG. 21.)

Peziza stercoraria Bull. Herb. Fr. *pl. 376, f. 1.* 1787.
Ascobolus pezizoides Pers.; J. F. Gmel. Syst. Nat. 1461. 1791.
Ascobolus furfuraceus Pers. Obs. Myc. **1**: 33. 1796.
Ascobolus stercorarius retisporus Clements, Bot. Surv. Nebr. **5**: 9. 1901.

Apothecia scattered or thickly gregarious, sessile, often partially buried in the substratum, later becoming superficial or subsuperficial, at first globose or subglobose and closed, opening by a circular aperture and expanding, finally scutellate to discoid, reaching a diameter of 5 mm., externally pale-yellow or greenish (whitish in dried specimens), furfuraceous; hymenium concave to plane, at first yellowish or greenish, becoming dotted over with the protruding asci which appear black, finally entirely black; asci clavate, gradually tapering below into a stem-like base, reaching a length of 200–250 μ and a diameter of 30 μ, 8-spored; spores partially 2-seriate or irregularly disposed, ellipsoid, thick-walled, at first hyaline and more or less granular within, becoming violet, later brown, smooth, becoming sculptured, 11–14 $\times$ 20–30 μ; spore-sculpturing taking the form of ridges and crevices which have a tendency to be longitudinally disposed, occasionally anastomosing and then giving the spore a reticulate appearance; paraphyses slender, about 2 μ in diameter, scarcely enlarged above, embedded in golden-yellow mucilaginous substance.

On dung of various kinds.

TYPE LOCALITY: Europe.

DISTRIBUTION: Connecticut to California, Louisiana and Bermuda, probably throughout North America; also in Europe and doubtless widely distributed.

ILLUSTRATIONS: Pers. Obs. Myc. **1**: *pl. 4, f. 3–6;* Ann. Sci. Nat. V. **10**: *pl. 6, f. 6;* Bull. Herb. Fr. *pl. 376, f. 1;* Cooke, Handb. **2**: 728, *f. 338;* Jour. Bot. **2**: 150, *f. 1;* Trans. Linn. Soc. **24**: *pl. 25, f. 9, 10;* Gill. Champ. Fr. Discom. *pl. 84, f. 2;* E. & P. Nat. Pfl. 1^1: 192, *f. 154, J–L;* Massee, Brit. Fungus-Fl. **4**: 162, *f. 1–7;* Rab. Krypt.-Fl. 1^3: 1112, *f. 1, 2, 3, 6;* Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 29, f. 2.*

EXSICCATI: Ravenel, Fungi Am. *312, 632.*

Common and widely distributed everywhere.

2. **Ascobolus immersus** Pers. Obs. Myc. 1: 35. 1796. (PLATE 7, FIG. 5.)

Ascobolus macrosporus Crouan, Ann. Sci. Nat. IV. 7: 173. 1857.
Ascobolus gigasporus De-Not. Comm. Critt. Ital. 1: 360. 1863.

Apothecia minute, scattered or thickly gregarious, but not usually crowded, at first immersed or partially immersed in the substratum, expanding and becoming subturbinate, not exceeding 1–2 mm. in diameter, externally greenish-yellow, becoming brownish, pilose; hymenium plane or more often convex, similar in color to the outside of the apothecium; asci comparatively few to each plant, very large, clavate, at maturity projecting half their length above the surface of the hymenium, 3–10 appearing at one time and easily visible with the hand-lens as several black, spike-like protuberances, reaching a length of 500–600 μ and a diameter of 90–100 μ, 8-spored but often with only a part of the spores developed; spores very large, 2-seriate or irregularly disposed, broad-ellipsoid and each surrounded by a hyaline, mucilaginous envelope, at first hyaline, then violet, finally brown, becoming sculptured, 20–35 $\times$ 50–70 μ; spore-sculpturing taking the form of sparse reticulations, consisting of one to several anastomosing bands which extend from one end of the spore to the other or diagonally across its surface; paraphyses very slender, scarcely enlarged above and embedded in a greenish-yellow mucilaginous substance.

On dung of various kinds, especially after it has weathered for a long time.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to Colorado and Bermuda, probably throughout North America; also in Europe.

ILLUSTRATIONS: Ann. Sci. Nat. IV. **7**: *pl. 4, f. B, 5–8;* Ann. Sci. Nat. V. **10**: *pl. 8, f. 17;* Rab. Krypt.-Fl. 1^3: 1112, *f. 4–5;* Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 31, f. 1;* Bull. Torrey Club **39**: *pl. 11, f. 20–25;* Ann. Mag. Nat. Hist. III. **15**: *pl. 17, f. 33;* E. & P. Nat. Pfl. 1^1: 192, *f. 154, D–F.*

3. **Ascobolus glaber** Pers. in Römer's Neues Mag. Bot. 1: 115. 1794.

Apothecia scattered or gregarious, at first subglobose, becoming short-cylindric, finally expanded and more or less discoid, tapering below into a stem-like base, reaching a diameter of 0.5 to 1 mm., externally greenish-yellow, smooth; hymenium

plane, becoming convex, at first similar in color to the outside of the apothecium, becoming almost entirely black; the component cells of the excipulum roundish or irregularly angular, reaching a diameter of 25 μ, the walls hyaline to pale-yellowish; asci clavate, strongly swollen above, tapering rather abruptly below into a short stem-like base, reaching a length of 300–350 μ and a diameter of 40 μ, 8-spored; spores 2-seriate or irregularly crowded, ellipsoid, with the ends rather strongly narrowed, hyaline, becoming violet, finally pale-brown, becoming sculptured, 12–13 $\times$ 23–38 μ; spore-sculpturing taking the form of longitudinally or irregularly disposed, anastomosing ridges or crevices; paraphyses slender, about 2 μ in diameter, slightly enlarged above.

On dung of various kinds.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to Iowa; also in Europe.

ILLUSTRATIONS: Ann. Sci. Nat. IV. **10**: *pl. 13, f. A;* Ann. Sci. Nat. V. **10**: *pl. 7, f. 13, 15;* Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 31, f. 2;* E. & P. Nat. Pfl. 1^1: 192, *f. 154, G, H.*

4. **Ascobolus Winteri** Rehm in Rab. Krypt.-Fl. 1^3: 1124. 1896.

Apothecia gregarious or densely crowded, sessile, subglobose, externally yellowish, smooth, reaching a diameter of 0.2–0.4 mm.; hymenium convex, yellowish-brown, finally almost black; asci cylindric or subcylindric to clavate, tapering below rather gradually into a long stem-like base, reaching a length of 180–200 μ and a diameter of 30–33 μ, when mature often stretching to 300 or 400 μ long; spores 2-seriate or irregularly disposed, ellipsoid, at first hyaline, then brown, for a long time smooth, becoming sculptured, 12–16 $\times$ 24–33 μ; spore-sculpturing consisting of minute warts and reticulations, the reticulations consisting of light lines which give rise to irregular or hexagonal meshes, the meshes ranging from 2 to 5 μ in diameter; paraphyses filiform, septate, about 2.5 μ thick, a little enlarged above, embedded in a yellow mucilaginous substance.

On goose dung.

TYPE LOCALITY: Leipzig, Germany.

DISTRIBUTION: New York; also in Europe.

ILLUSTRATIONS: Bull. Torrey Club **39**: *pl. 10, f. 12–14;* also *pl. 15.*

5. **Ascobolus americanus** (Cooke & Ellis) Seaver, sp. nov.

Ascobolus Leveillei americanus Cooke & Ellis, Grevillea **5**: 52. 1876.

Apothecia thickly gregarious, crowded, or more rarely scattered, often nestling in little cavities in the substratum or entirely superficial, at first globose or pyriform with the small end directed downward, gradually expanding and becoming subdiscoid above, never exceeding 1 mm. in diameter, externally yellowish; hymenium plane or convex, at first greenish-yellow, becoming almost black, strongly roughened by the protruding asci; the component cells of the excipulum irregularly 6-sided, reaching a diameter of 12–18 μ; asci broad-clavate, gradually tapering below into a stem-like base, reaching a length of 200–250 μ and a diameter of 35–40 μ, when mature often stretching to 400 μ, 8-spored; spores 2-seriate or irregularly disposed, ellipsoid, thick-walled, at first hyaline, becoming violet, then brown, for long time smooth, finally becoming delicately sculptured, 15–17 $\times$ 30–40 μ; spore-sculpturing taking the form of minute granules which give to the surface of the spore a minutely warted or pitted appearance; paraphyses about 2 or 3 μ in diameter, scarcely thickened above, embedded in a greenish-yellow mucilaginous substance.

On dung of various kinds.

TYPE LOCALITY: Newfield, New Jersey.

DISTRIBUTION: New Jersey to New York and Kansas.

EXSICCATI: N. Am. Fungi *2333* (as *Ascobolus glaber* Pers.); N. Am. Fungi *2620* (as *Ascobolus vinosus* Berk.).

6. **Ascobolus Leveillei** Boud. Ann. Sci. Nat. V. **10**: 225. 1869.

Ascobolus Boudieri Quél. Ench. Fung. 293. 1886.

Apothecia thickly gregarious, minute, sessile, subglobose, with the base partially immersed in the substratum or entirely superficial, externally brownish, slightly roughened, reaching a diameter of 0.5 mm.; hymenium becoming convex, at first greenish, finally becoming darker, almost black; the component cells of the excipulum roundish or angular, reaching a diameter of 15–20 μ; asci clavate, tapering below into a stem-like base, reaching a length of 140–160 μ and a diameter of 28–30 μ, 8-spored; spores 2-seriate or irregularly disposed, at first hyaline, then purple, later brown, ellipsoid, for a long time smooth, becoming minutely sculptured, 13–15 $\times$ 24–28 μ; spore-sculpturing taking the form of minute granules which give to the

surface of the spore a delicately warted or pitted appearance; paraphyses slender, septate, about 3 μ in diameter, slightly thickened above, embedded in a mucilaginous substance.

On dung of different kinds.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to Iowa; also in Europe.

ILLUSTRATIONS: Ann. Sci. Nat. V. **10**: *pl. 7, f. 16;* Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 30, f. 2;* Bull. Torrey Club **39**: *pl. 11, f. 27.*

7. **Ascobolus viridulus** Phill. & Plow. Grevillea **8**: 103. 1879.

?Ascobolus aerugineus Fries, Obs. Myc. **2**: 310. 1818.
?Ascobolus crenulatus Karst. Fungi Fenn. Exsicc. *763.* 1868.

Apothecia scattered, sessile, at first nearly globose, expanding and becoming discoid, reaching a diameter of 0.5–1 mm., externally pale yellowish-green, coarsely furfuraceous; hymenium at first concave, becoming plane or slightly convex, yellowish-green, becoming darker as the plants mature on account of the dark-colored spores, roughened by the protruding asci; asci clavate, reaching a length of 75 μ and a diameter of 12–15 μ, 8-spored; spores 1-seriate, or finally becoming 2-seriate or irregularly crowded, ellipsoid with the ends quite strongly narrowed, hyaline, then violet, finally pale-brown, becoming sculptured, 7–8 $\times$ 12–14 μ; spore-sculpturing consisting of anastomosing ridges or crevices which are mostly longitudinal; paraphyses very slender, slightly enlarged above.

On dung of dogs and birds.

TYPE LOCALITY: Great Britain.

DISTRIBUTION: New York; also in Europe.

8. **Ascobolus candidus** Schröt. Krypt.-Fl. Schles. **3**^{2}: 55. 1893.

Apothecia scattered, roundish, becoming subdiscoid, entirely pallid-white; hymenium becoming slightly convex, finally dotted over with the protruding asci and becoming darker with age; the excipular cells irregular and reaching a diameter of 12–15 μ; asci clavate, gradually tapering below into a stem-like base, reaching a length of 100–150 μ and a diameter of 20 μ, 8-spored; spores 1-seriate, 2-seriate or irregularly disposed, ellipsoid, becoming pale-violet, later pale-brown, 8–10 $\times$ 13–16 μ, becoming sculptured; spore-sculpturing consisting of minute warts; paraphyses strongly enlarged above, reaching a diameter of about 4 μ at their apices, hyaline.



ASCOBOLUS MAGNIFICUS

On the dung of rabbits and goats.
TYPE LOCALITY: Europe.
DISTRIBUTION: New York; also in Europe.

9. **Ascobolus magnificus** Dodge, Mycologia **4**: 218. 1912. (PLATE 6.)

Apothecia scattered or closely crowded, sessile, at first globose and appearing closed or nearly so, white or whitish, gradually opening, becoming cup-shaped with the smooth margin inrolled, finally becoming subscutellate, externally pruinose from the projecting tips of thin-walled, hair-like hyphal branches which finally become discolored and brownish, reaching a diameter of 0.5–2.7 cm.; hymenium concave or nearly plane, at first greenish-yellow, roughened by the protruding asci and becoming nearly black at maturity on account of the dark-colored spores; asci cylindric to clavate, reaching a length of 200–300 μ and a diameter of 18–25 μ, 8-spored; spores becoming irregularly 2-seriate, ellipsoid, at first hyaline, then pale-lilac, finally rose-purple or violet, fading to brown in aged specimens, smooth, becoming sculptured, 12–14 $\times$ 20–25 μ, germinating by one germ-pore; spore-sculpturing consisting of one faint line extending from one end of the spore to the other or obliquely across its surface; paraphyses filiform, slightly enlarged above, septate, reaching a diameter of 5–7 μ at their apices, filled with a greenish granular content.

On horse dung in damp-chamber cultures.
TYPE LOCALITY: New York City.
DISTRIBUTION: New York and Porto Rico.
ILLUSTRATIONS: Mycologia **4**: *pl. 72, f. 1–8*, and *pl. 73*.

10. **Ascobolus carbonarius** Karst. (?Fungi Fennici *463*. 1866); Not. Fauna Fl. Fenn. **11**: 202. 1870. (PLATE 7, FIG. 4.)

Ascobolus atrofuscus Phill. & Plow. Grevillea **2**: 186. 1873.
Ascobolus carbonicola Boud. Bull. Bot. Soc. Fr. **24**: 310. 1877.
Phaeopezia Nuttallii Ellis & Ev. N. Am. Fungi *2908*. 1893.

Apothecia scattered or more often crowded into congested masses several cm. in diameter, at first globose or subglobose, expanding and becoming scutellate, reaching a diameter of 4–5 mm., externally yellowish-green, becoming dark-brown, coarsely furfuraceous; hymenium plane or nearly plane with the margin slightly elevated, at first greenish, becoming dotted over with

the ends of the protruding asci, gradually becoming darker, finally almost black; asci clavate, gradually tapering below into a long, slender, stem-like base, reaching a length of 200–225 μ and a diameter of 23–27 μ, 8-spored; spores 1-seriate or crowded together so as to become partially 2-seriate, ellipsoid with the ends rather strongly narrowed, thick-walled, the ends often capped with a thickening of the epispore, giving the spore a truncate appearance, hyaline, becoming violet, finally brown or brownish-black, becoming sculptured, 12–14 × 20–24 μ; spore-sculpturing taking the form of wart-like projections, the spaces between appearing lighter, giving rise to reticulate markings; paraphyses about 3 μ in diameter and scarcely thickened above, embedded in greenish-yellow mucilaginous substance.

On burnt places and on charcoal beds.

TYPE LOCALITY: Finland.

DISTRIBUTION: New York to Iowa and West Virginia; also in Europe.

ILLUSTRATIONS: Ann. Sci. Nat. V. **10**: *pl. 5, f. 4* (as *Ascobolus viridis* Curr.); Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 29, f. 1;* Bull. Torrey Club **39**: *pl. 10, f. 9, 10, 11;* Mycologia **8**: *pl. 184, f. 7–8.*

EXSICCATI: N. Am. Fungi *2908* (as *Phaeopezia Nuttallii* Ellis & Ev.).

11. **Ascobolus geophilus** Seaver, Mycologia **8**: 96. 1916. (PLATE 7, FIG. 2.)

Apothecia scattered, gregarious or more rarely crowded, at first subglobose, expanding, becoming scutellate to discoid, reaching a diameter of 5 mm., externally greenish-yellow, furfuraceous to nearly smooth; hymenium plane or slightly concave, similar in color to the outside of the apothecium, roughened by the protruding asci and becoming darker as the spores mature, finally almost black; asci clavate, gradually tapering below into a stem-like base, reaching a length of 200–250 μ and a diameter of 16–18 μ, 8-spored but often with only 4 spores developed; spores 1-seriate, becoming 2-seriate or irregularly crowded, ellipsoid, at first hyaline, becoming violet, then brown, 12–13 μ × 22–25 μ, becoming sculptured; spore-sculpturing consisting of verrucose markings and reticulations, the reticulations consisting of light lines which give rise to an irregular network; paraphyses very slender, sparingly branched, reaching a diameter of 2 μ, scarcely

enlarged above, embedded in golden-yellow mucilaginous substance.

On damp soil.

TYPE LOCALITY: The New York Botanical Garden.

DISTRIBUTION: New York to Iowa and Colorado.

ILLUSTRATIONS: Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 30, f. 1;* Mycologia **8**: *pl. 84, f. 1–2.*

This species has been mistaken for *Ascobolus viridis* of Europe which is very different in spore characters (see PLATE 7).

Since the above account was prepared, a fine specimen of this species has been received from Dr. G. R. Bisby, who collected it on damp soil at the Agricultural College in Winnipeg, Canada.

12. **Ascobolus subglobosus** Seaver, Mycologia **8**: 96. 1916. (PLATE 7, FIG. 3.)

Apothecia gregarious or thickly crowded, at first nearly globose, expanding and becoming almost discoid, reaching a diameter of 3–5 mm., externally smooth, greenish-yellow; hymenium plane, convex or irregularly convolute, at first similar in color to the outside of the apothecium, becoming dotted over with the ends of the protruding asci, finally entirely black; asci subcylindric to clavate, gradually tapering below into a stem-like base, 8-spored; spores at first 1-seriate, becoming crowded and partially 2-seriate as they mature, at first hyaline and containing one oil-drop which is often surrounded by numerous smaller ones, becoming violet, finally almost black and opaque, becoming sculptured, reaching a diameter of 18–20 μ or 18 $\times$ 20 μ; spore-sculpturing taking the form of numerous vein-like reticulations, which consist apparently of crevices, the spaces between the crevices giving rise to prominent projections visible about the periphery of the spore, the spaces between appearing as notches, giving the spore a very ragged appearance; paraphyses scarcely enlarged above, embedded in golden-yellow mucilaginous substance.

On damp soil in woods and on the margin of a small pond.

TYPE LOCALITY: Woods near Yonkers, New York.

DISTRIBUTION: Known only from the type locality.

ILLUSTRATION: Mycologia **8**: *pl. 84, f. 5–6.*

13. **Ascobolus albinus** Seaver, Mycologia **8**: 95. 1916.

Apothecia gregarious, at first subglobose, expanding and becoming scutellate, reaching a diameter of 4–5 mm., externally

pure white, minutely rough; hymenium plane or nearly plane, at first whitish, becoming darkened by the maturing spores; asci cylindric or subcylindric, 8-spored, reaching a length of about 200 μ and a diameter of about 25 μ; spores at first obliquely 1-seriate, becoming 2-seriate or irregularly crowded, ellipsoid, becoming purple, then brown, becoming sculptured, 12 $\times$ 20-26 μ; spore-sculpturing consisting of minute warts; paraphyses very slender, hyaline.

On damp soil in woods.

TYPE LOCALITY: Woods near Yonkers, New York.

DISTRIBUTION: Known only from the type locality.

Apparently differing from *A. carbonarius* in color and occurrence on damp soil where there was no trace of fire.

14. **Ascobolus xylophilus** Seaver, Mycologia **3**: 61. 1911.

Apothecia gregarious, sessile, at first subglobose, expanding and becoming scutellate to discoid, reaching a diameter of 1–2 mm., externally brownish; hymenium plane or slightly concave, greenish, becoming darker with age, finally almost black; asci cylindric to clavate, tapering below into a stem-like base, reaching a length of 175–235 μ and a diameter of 30–35 μ, 8-spored; spores 1-seriate or becoming partially 2-seriate above, large, ellipsoid, with the ends rather strongly narrowed, at first hyaline, becoming beautifully purple, then brown, smooth, becoming sculptured, 13–15 $\times$ 35–38 μ; spore-sculpturing taking the form of minute warts; paraphyses slender, scarcely enlarged above.

On the weathered surface of the wood of some coniferous tree.

TYPE LOCALITY: Geneva Creek Canyon, Colorado (elevation about 9,000 feet).

DISTRIBUTION: Known only from the type locality.

One of the few species of the genus known to occur on wood.

15. **Ascobolus striisporus** (Ellis & Dearn.) Seaver, comb. nov.

Chlorosplenium striisporum Ellis & Dearn. Proc. Acad. Sci. Phila. **1895**: 429. 1896.

Apothecia sessile, hemispheric, with the margin incurved and notched, externally greenish-yellow, verrucose-squamose, reaching a diameter of 1–1.5 mm.; hymenium concave or nearly plane, pale-yellowish, becoming darker in dried specimens; asci clavate,

SPECIES OF ASCOBOLUS

reaching a length of 150–175 μ and a diameter of 15–18 μ; spores at first 1-seriate, becoming 2-seriate or irregularly crowded in the upper part of the ascus, at first hyaline, becoming purple, then brown, ellipsoid, about 10–12 $\times$ 20 μ at maturity, becoming sculptured; spore-sculpturing consisting of longitudinal anastomosing ridges; paraphyses about 2 μ in diameter, slightly enlarged above.

On decaying leaves.

TYPE LOCALITY: London, Ontario, Canada.

DISTRIBUTION: Known only from the type locality.

Doubtless on account of its greenish color this was mistaken by Ellis for a *Chlorosplenium*. Our studies show it to be a true *Ascobolus*.

Since this work was submitted to the press, the writer has received and examined a specimen of this species collected on old wet leaves at Manitoba Agricultural College, Winnipeg, Canada, by Dr. G. R. Bisby, I. L. Conners, and W. L. Gordon, No. *3647*. So far as we are aware, this is only the second collection of this species on record. The species is rather obscure but is probably more common than would be indicated by the above records.

16. **Ascobolus epimyces** (Cooke) Seaver, comb. nov.

Chlorosplenium epimyces Cooke, Bull. Buffalo Acad. Sci. **2**: 299. 1875.

Apothecia gregarious, sessile, at first nearly globose, expanding and becoming discoid or subdiscoid, scarcely exceeding 1 mm. in diameter, greenish-yellow, then olivaceous, pulverulent, nearly black in dried specimens; hymenium becoming plane, a little lighter in color than the outside of the apothecium; asci clavate, reaching a length of 120 μ and a diameter of 12–15 μ, gradually tapering below into a slender stem-like base; spores 2-seriate above or irregularly crowded, 1-seriate below, fusoid, at first hyaline (then purple?), finally brown, 8–9 $\times$ 10–18 μ, becoming sculptured; spore-sculpturing consisting of delicate lines (apparently ridges) which freely anastomose and give the spore a decidedly striate appearance; paraphyses slender, slightly enlarged above, filled with yellowish-green coloring matter.

Originally reported on old *Corticium*, but apparently on the remains of some slime mould. Except for habitat and a slight difference in size and form of the spores, the species is similar to *Ascobolus viridulus* Phill. & Plow.

TYPE LOCALITY: Newfield, New Jersey.
DISTRIBUTION: Known only from the type locality.

DOUBTFUL AND EXCLUDED SPECIES

Ascobolus atrofuscus Phill. & Plow.; Harkness & Moore, Pacific Coast Fungi 38. 1880. This species was listed in the above work as a new species occurring on horse dung. The name is untenable since it was previously used by the same authors for a species occurring on burnt ground which has been made a synonym of *Ascobolus carbonarius* Karst. The application of the same name to a second species is doubtless due to an oversight. Nothing is known of the plant to which this name was applied in the Harkness and Moore report.

Ascobolus viridis Curr. Trans. Linn. Soc. **24**: 154. 1863. This species was reported by the writer from Iowa. Later studies showed that the plant to which this name was applied is quite different from the European species (see PLATE 7, FIG. 1) and the American species has therefore been redescribed under the name of *Ascobolus geophilus*. So far as known the European species has not been collected in America.

Ascobolus purpurascens Pers.; Calkins, Jour. Myc. **2**: 106. 1886. This species is listed by Calkins in his report of Florida fungi. We have no knowledge of the nature of the plants reported under this name.

Ascobolus brunneus Cooke, Fungi Brit. Exsicc. **3**: 87; Hedwigia **6**: 154. 1867. American specimens listed under this name have been referred to other species.

Ascobolus cubensis Berk. & Curt. Jour. Linn. Soc. **10**: 370. 1868. No specimen of this species has been seen and its identity is uncertain. From the description it appears to be close to *Ascobolus viridulus* Phill. & Plow.

Ascobolus vinosus Berk. in Hooker's Engl. Fl. 5^2: 209. 1836. American specimens referred to this species scarcely differ from *Ascobolus stercorarius* (Bull.) Schröt.

Ascobolus major Berk. & Curt. Grevillea **4**: 6. 1875. No specimen of this species has been seen and the description is so brief and incomplete as to give no clue to its identity.

Ascobolus conglomeratus Schw. Trans. Am. Phil. Soc. II. **4**: 178. 1832. This is one of the Hysteriaceae and has usually been treated as a synonym of *Angelina rufescens* (Schw.) Duby.

Ascobolus Trifolii Biv. Stirp. Rar. Sic. **4**: 27. 1816. This is not a true *Ascobolus*. It was made the type of the genus *Pseudopeziza* by Fuckel.

Ascobolus pusillus Boud. Bull. Soc. Bot. Fr. **24**: 310. 1877. The species has been reported from North America by Dodge.

10. **SACCOBOLUS** Boud. Ann. Sci. Nat. V. **10**: 228. 1869.

Apothecia minute, sessile, superficial or subsuperficial, externally smooth or pilose; hymenium concave, plane or convex, smooth or becoming roughened by the protruding asci; substance soft, fleshy or waxy, variable in color; asci cylindric to clavate, 4–8-spored; spores hyaline, then purple or bluish, later brown,

ellipsoid to subglobose, smooth or sculptured; spore-sculpturing consisting of minute warts or faint reticulations, united into a spore-mass; spore-mass elongated or subglobose; paraphyses filiform to clavate.

Type species, *Ascobolus Kerverni* Crouan.

Apothecia golden-yellow.	
Spores ellipsoid to subfusoid.	1. *S. Kerverni.*
Spores subglobose.	2. *S. portoricensis.*
Apothecia violaceous to blackish.	
Spores 7–8 × 15 μ.	3. *S. violascens.*
Spores 5 × 10–12 μ.	4. *S. depauperatus.*
Apothecia white.	5. *S. globuliferellus.*

1. Saccobolus Kerverni (Crouan) Boud. Ann. Sci. Nat. V. **10**: 229. 1869.

Ascobolus Kerverni Crouan, Ann. Sci. Nat. IV. **10**: 193. 1858.

Apothecia gregarious or thickly crowded, seated on a rather dense growth of yellowish or whitish mycelium, at first globose, expanding and becoming discoid or subdiscoid, externally golden-yellow, reaching a diameter of 1–2 mm.; hymenium at first concave or plane, becoming strongly convex, at first similar in color to the outside of the apothecium, becoming dotted over with the ends of the protruding asci which appear black on account of the dark-colored spores, entirely black when mature; asci very broad-clavate, gradually narrowed below into a stem-like base, reaching a length of 125 μ and a diameter of 30 μ, 8-spored; spore-mass elongated, reaching a length of 50–70 μ and a diameter of 20–27 μ; spores ellipsoid, with the ends narrowed, strongly swollen near the center, often subfusoid and slightly unequal-sided, hyaline, becoming violet, later brown, often delicately sculptured, 10 × 20–23 μ; spore-sculpturing taking the form of very delicate reticulations; paraphyses slender, branched rather freely, golden-yellow, thickened above.

On dung of various kinds.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to Colorado, Bermuda and Porto Rico, probably common throughout North America; also in Europe.

ILLUSTRATIONS: Ann. Sci. Nat. IV. **10**: *pl. 13, f. B;* Ann. Sci. Nat. V. **10**: *pl. 8, f. 18;* Ann. Mag. Nat. Hist. III. **15**: *pl. 17, f. 34;* Jour. Bot. **2**: 153, *f. 8;* Gill. Champ. Fr. Discom. *pl. 85, f. 1;* Rab. Krypt.-Fl. 1^3: 1111, *f. 6–7;* Massee, Brit. Fungus-Fl. **4**: 156, *f. 11;* Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 28, f. 2.*

2. **Saccobolus portoricensis** Seaver, sp. nov.

Apothecia gregarious or scattered, superficial, at first perfectly globose, becoming short-cylindric, finally discoid, smooth, pale-amber, reaching a diameter of 1 mm.; hymenium plane, becoming convex, at first similar in color to the outside of the apothecium, dotted over with the protruding asci, finally almost entirely black; hypothecium composed of rather loosely united roundish or angular cells; excipulum poorly developed, consisting of a palisade of mycelium; asci clavate, tapering below into a rather long slender base, reaching a length of 170–200 μ and a diameter of 35–40 μ, 8-spored; spore-mass at first elongated, finally subglobose, closely compact and opaque or nearly so, surrounded by a mass of purple granules, reaching a length of 40 μ and a diameter of 27–30 μ; spores at first subfusoid, finally subglobose, at first hyaline, then purple, finally pale-brown, becoming sculptured, about 17 $\times$ 20 μ; spore-sculpturing taking the form of very delicate reticulations, scarcely roughening the surface of the spore; paraphyses about 3 μ thick, scarcely enlarged above, filled with orange granules.

Cultivated on cow dung.

Type grown on dung sent from Porto Rico, autumn, 1915 (Herb. The New York Botanical Garden).

DISTRIBUTION: Known only from the type locality.

Distinguished by its subglobose spores.

3. **Saccobolus violascens** Boud. Ann. Sci. Nat. V. **10**: 230. 1869.

?Saccobolus neglectus Boud. Ann. Sci. Nat. V. **10**: 231. 1869.

Apothecia scattered, superficial, pale-violaceous, with a whitish mycelial growth about the base, not exceeding 1 mm. in diameter; hymenium becoming strongly convex, roughened by the protruding asci, at first similar in color to the outside of the apothecium, finally almost black; excipulum poorly developed, consisting of loosely interwoven hyphae which, with transmitted light, appear very dilutely violaceous; asci very broad-clavate, gradually tapering below into a stem-like base, reaching a length of 175 μ and a diameter of 30–35 μ, 8-spored; spore-mass densely compact, elongated, reaching a length of 40–60 μ and a diameter of 20 μ; spores elongated, slightly unequal-sided or almond-shaped, at first hyaline or very dilutely violaceous. 7-8 × 15 μ

On dung of various kinds, especially on goat dung.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to Iowa; also in Europe.

ILLUSTRATIONS: Ann. Sci. Nat. V. **10**: *pl. 8, f. 19*, and *pl. 9, f. 20;* Phill. Brit. Discom. *pl. 9, f. 55;* Rab. Krypt.-Fl. 1^3: 1111, *f. 1–5;* E. & P. Nat. Pfl. 1^1: 192, *f. 154, B, C.*

4. **Saccobolus depauperatus** (Berk. & Br.) Phill. Brit. Discom. 296. 1887.

Ascobolus depauperatus Berk. & Br. Ann. Mag. Nat. Hist. III. **15**: 448. 1865.

Apothecia scattered or gregarious, with a narrowed base which is immersed in the substratum, expanding and becoming discoid, scarcely exceeding 0.3 mm. in diameter; hymenium even with, or slightly elevated above the surface of the substratum, at first whitish with a tinge of violet, finally almost black; the excipular cells angular or rounded and reaching a diameter of 5–7 μ, the walls beautifully lilac, fading to brown, giving rise above to a palisade of mycelia which resemble paraphyses except that they are more deeply colored; asci very broad-clavate, the ends strongly truncate, tapering gradually below into a stem-like base, reaching a length of 65–70 μ and a diameter of 15–17 μ, 8-spored; spore-mass elongated, reaching a length of 30 μ and a diameter of 11–12 μ; spores ellipsoid, the ends strongly narrowed, smooth or minutely sculptured, at first hyaline, becoming deep-violet, finally blackish-brown, 5 $\times$ 10–12 μ; paraphyses slender, slightly enlarged above, reaching a diameter of 3–4 μ, hyaline, or very faintly colored.

On horse dung.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York; also in Europe.

ILLUSTRATION: Ann. Mag. Nat. Hist. III. **15**: *pl. 14, f. 6.*

5. **Saccobolus globuliferellus** Seaver, sp. nov.

Apothecia widely scattered, minute, scarcely visible, not exceeding 0.3 mm. in diameter, milk-white; hymenium becoming convex, at first white like the outside of the apothecium, finally dotted over with the ends of the asci which appear black on account of the dark-colored spores; asci broad-ovoid above, tapering below into a stem-like base, reaching a length of 60 μ and a diameter of 23–25 μ, 8-spored; spore-mass nearly rounded, about 23 $\times$ 20 μ; spores at first loosely disposed in the ascus, finally collecting into the loosely united spore-mass, ellipsoid, at first hyaline, later assuming a faded-blue color which finally

changes to smoky-blackish, becoming sculptured, 7 × 12–14 μ; spore-sculpturing consisting of minute scattered warts; paraphyses branched, rather strongly enlarged above where they reach a diameter of 7–8 μ, hyaline.

On horse dung.

TYPE LOCALITY: New York City.

DISTRIBUTION: Known only from the type locality.

The species, which was collected during the autumn of 1914, seems to differ in several particulars from any of the other species of the genus studied. The apothecia are white and very small. The spore masses are nearly rounded and the spores themselves more blue than purple and distinctly warted rather than reticulated. The species has been encountered only once.

DOUBTFUL SPECIES

Saccobolus obscurus (Cooke) Phill. Brit. Discom. 295. 1887; *Ascobolus obscurus* Cooke, Grevillea **4**: 112. 1876. Specimens were distributed under this name in Clements, Crypt. Form. Colo. *301.* No *Saccobolus* could be found in our copy of this number.

11. **ALEURIA** Fuckel, Symb. Myc. 325. 1869.

Apothecia gregarious, scattered or cespitose, sessile or stipitate, externally smooth or clothed with a delicate white tomentum; hymenium concave or plane, even or lacunose, usually bright-orange or red; asci cylindric or subcylindric, 8-spored; spores ellipsoid, at first smooth, becoming sculptured; spore-sculpturing assuming the form of reticulations, often also with apiculi, rings, etc., at the ends of the spores; paraphyses slender, usually strongly enlarged at their apices.

Type species, *Peziza aurantia* Pers.

Apothecia sessile or subsessile.
- Spores apiculate but not marked with a ring at the ends.
  - Apothecia reaching a diameter of several cm.; spores 18–20 μ long. — 1. *A. aurantia.*
  - Apothecia not exceeding a diameter of 2 cm.; spores 9–15 μ long.
    - Spores 7 × 14–15 μ. — 2. *A. wisconsinensis.*
    - Spores 5 × 9–10 μ. — 3. *A. cestrica.*
- Spores marked with a ring at either end. — 4. *A. bicucullata.*

Apothecia strongly stipitate.
- Occurring in coniferous woods; stems springing from a dense mass of mycelium. — 5. *A. rhenana.*
- Occurring usually singly among mosses in deciduous woods. — 6. *A. rutilans.*

1. **Aleuria aurantia** (Pers.) Fuckel, Symb. Myc. 325. 1869. (PLATE 8, FIG. 1.)

Elvela coccinea Schaeff. Fung. Bavar. **4**: 100. 1774. Not *Elvela coccinea* Scop. 1772.
Peziza coccinea Bull. Herb. Fr. *pl. 474.* 1789.
Peziza aurantia Pers. Obs. Myc. **2**: 76. 1797.
Scodellina aurantiaca S. F. Gray, Nat. Arrang. Pl. 668. 1821.
Otidea aurantia Massee, Brit. Fungus-Fl. **4**: 448. 1895.

Apothecia gregarious or cespitose, sessile, at first globose, gradually expanding, finally reaching a diameter of 5–6 cm. though often smaller, shallow, cup-shaped and usually regular in form when young, becoming irregular and often variously contorted with age or from mutual pressure, occasionally discoid and rarely one-sided and *Otidea*-like, externally whitish, pruinose; hymenium concave or almost plane, bright-orange, the color fading in dried specimens; asci cylindric or subcylindric, reaching a length of 175–250 μ and a diameter of 12–15 μ; spores 1-seriate, usually obliquely arranged in the ascus, the ends often overlapping, at first smooth and usually containing two large oil-drops, finally becoming sculptured, 9–10 $\times$ 18–22 μ; spore-sculpturing taking the form of reticulations which are regular in form and shallow, except at the ends of the spore where the ridges project, giving rise to apiculi; paraphyses strongly and rather abruptly enlarged at their apices, the ends often subglobose, reaching a diameter of 7–8 μ, filled with orange granules.

On damp soil in woods and open places, occasionally on lawns but more often on naked clayey soil.

TYPE LOCALITY: Europe.

DISTRIBUTION: Newfoundland to Washington, California and West Virginia; also in Europe.

ILLUSTRATIONS: Bolton, Fungi Halifax *pl. 100;* Bull. Herb. Fr. *pl. 474;* Cooke, Mycographia *pl. 52, f. 203;* Fl. Danici *pl. 157;* Schaeff. Fung. Bavar. *pl. 148;* Sow. Engl. Fungi *pl. 78;* Gill. Champ. Fr. Discom. *pl. 37, f. 2;* Pat. Tab. Fung. *f. 278;* E. & P. Nat. Pfl. 1^1: 186, *f. 150, J–L;* Phill. Brit. Discom. *pl. 3, f. 14;* Rab. Krypt.-Fl. 1^3: 920, *f. 1–3;* Batsch, Elench. Fung. **2**: *pl. 28, f. 158;* Boud. Ic. Myc. *pl. 313;* Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 17, f. 1.*

EXSICCATI: N. Am. Fungi *836;* Ellis & Ev. Fungi Columb. *15.*

One of the commonest and most widely distributed of the

larger cup-fungi, occurring from the Atlantic to the Pacific and probably throughout temperate North America as well as abroad. Also it is attractive and easily recognized by means of its bright-orange color. If there is any doubt of its identity, this can be easily dispelled by an examination of the beautifully reticulated spores.

2. **Aleuria wisconsinensis** Rehm, Ann. Myc. **2**: 34. 1904.

Apothecia gregarious, sessile, becoming discoid, contracted at the base into a slightly stem-like base, the margin thick and entire, at length becoming more or less convolute, externally light-colored, yellowish, tomentose, reaching a diameter of 0.5–2 cm.; hymenium slightly concave or plane, even or convolute, bright orange-red; the excipular cell walls subhyaline; asci cylindric or subcylindric, reaching a length of 200 μ and a diameter of 10 μ, 8-spored; spores 1-seriate, ellipsoid, hyaline, containing two large oil-drops, becoming sculptured, about 7 $\times$ 14–15 μ; spore-sculpturing taking the form of reticulations; paraphyses hyaline, filiform, septate, reaching a diameter of 5 μ at their apices and about 3 μ below.

On damp soil.

TYPE LOCALITY: Blue mounds, Wisconsin.

DISTRIBUTION: Wisconsin and Minnesota.

3. **Aleuria cestrica** (Ellis & Ev.) Seaver, comb. nov.

Peziza cestrica Ellis & Ev. Jour. Myc. **1**: 152. 1885.
Humaria cestrica Sacc. Syll. Fung. **8**: 133. 1889.
Aleurina Lloydiana Rehm, Ann. Myc. **2**: 35. 1904.
Peziza Commonsii Ellis & Ev. (in herb.).

Apothecia gregarious or densely crowded, sessile or subsessile, and circular or becoming irregular from mutual pressure, reaching a diameter of 5–6 mm. or rarely 1 cm., externally yellowish-orange; hymenium shallow-concave or becoming nearly plane, orange; asci cylindric or clavate, reaching a length of 150 μ and a diameter of 10 μ, 8-spored; spores 1-seriate, ellipsoid, at first smooth and containing two distinct oil-drops, 5–7 $\times$ 10–11 μ, becoming sculptured; spore-sculpturing taking the form of very fine reticulations, with one or two projections at either end and giving the spore an apiculate appearance (the mature spores resembling those of *Aleuria aurantia* except that they are much smaller), hyaline or subhyaline; paraphyses slender, rather



1. ALEURIA AURANTIA
2. ALEURINA ATROVINOSA

abruptly thickened above, where they reach a diameter of 3–4 μ.

On damp ground, usually among mosses.

TYPE LOCALITY: West Chester, Pennsylvania.

DISTRIBUTION: Known only from the type locality.

EXSICCATI: Fungi Columb. *1733.*

4. **Aleuria bicucullata** Boud. Bull. Soc. Bot. Fr. **28**: 93. 1881.

Peziza bicucullata Boud. Bull. Soc. Myc. Fr. **1**: 103. 1885.
Humaria bicucullata Quél. Ench. Fung. 288. 1886.

Apothecia gregarious or crowded, sessile, at first subglobose, expanding and becoming subdiscoid or irregular from mutual pressure, externally pale-orange or whitish, pruinose, reaching a diameter of 5 mm. to 1 cm.; hymenium concave or nearly plane, bright-orange or yellow; asci cylindric or subcylindric, reaching a length of 200 μ and a diameter of 12–15 μ; spores 1-seriate, obliquely disposed, at first smooth and containing one or two oil-drops, finally becoming sculptured, 10–12 × 20–23 μ, including sculpturing; spore-sculpturing consisting of a ring-like or hood-like process at one or both ends of the spore (especially where the spores come into contact with each other in the ascus) and irregular or often broken reticulations, the ridges of the reticulations conspicuous and giving rise to the rather sharp-pointed, spine-like projections, the terminal projections larger and giving the spore an apiculate appearance; paraphyses rather strongly enlarged above, where they reach a diameter of 5 μ.

On bare ground or among mosses.

TYPE LOCALITY: France.

DISTRIBUTION: Wisconsin and Ontario; also in Europe.

ILLUSTRATIONS: Bull. Soc. Bot. Fr. **28**: *pl. 3, f. 1;* Boud. Ic. Myc. *pl. 318;* Mycologia **6**: *pl. 144, f. 3;* Pat. Tab. Fung. *f. 371.*

5. **Aleuria rhenana** Fuckel, Symb. Myc. 325. 1869. (PLATE 9, FIG. 2.)

?Peziza radiculata Sow. Engl. Fungi *pl. 114* (with descr.). 1797.
Peziza splendens Quél. Champ. Jura 388. 1873.
?Peziza sordescens Berk. & Curt. Grevillea **3**: 150. 1875.
Sarcoscypha rhenana Sacc. Syll. Fung. **8**: 157. 1889.
?Sarcoscypha radiculata Sacc. Syll. Fung. **8**: 156. 1889.
?Geopyxis sordescens Sacc. Syll. Fung. **8**: 69. 1889.

Apothecia gregarious or cespitose, stipitate, reaching a diameter of 1–2 cm. and about half as deep, externally white or

whitish, tomentose, the tomentum consisting of poorly developed hairs; stems very variable in length, reaching 1–2 cm., irregular, often clinging together in clumps, attached at the base by a dense mass of white mycelium which penetrates into the substratum binding together the leaf-mould in which the plants grow, expanding abruptly above into the apothecium; hymenium concave, bright-orange, the color fading in dried specimens; asci cylindric above, tapering below into a stem-like base, often becoming spirally twisted, especially in dried specimens, reaching a length of 300–350 μ and a diameter of 15–17 μ; spores 1-seriate, obliquely arranged and with the ends overlapping, ellipsoid, at first smooth, becoming sculptured, 12 × 20–23 μ; spore-sculpturing assuming the form of reticulations, the meshes of the reticulations reaching a diameter of 3 μ or more, rarely 5 or 6 μ, the edges of the ridges extending 1–2 μ beyond the periphery of the spore; paraphyses enlarged above, where they reach a diameter of 6 μ, filled with orange granules.

On the ground in coniferous woods.

TYPE LOCALITY: Europe.

DISTRIBUTION: Pennsylvania to Alabama and west to Washington; also in Europe.

ILLUSTRATIONS: Boud. Ic. Myc. *pl. 314;* Cooke, Mycographia *pl. 112, f. 400;* Mycologia **6**: *pl. 142* (lower figure) and *pl. 144, f. 2.*

6. **Aleuria rutilans** (Fries) Gill. Champ. Fr. Discom. 53. 1879. (PLATE 9, FIG. 1; 45, FIG. 22.)

Peziza rutilans Fries, Syst. Myc. **2**: 68. 1822.
Leucoloma rutilans Fuckel, Symb. Myc. 318. 1869.
Humaria rutilans Sacc. Syll. Fung. **8**: 133. 1889.
Sarcoscypha albovillosa Rehm, Ann. Myc. **2**: 33. 1904.

Apothecia gregarious or scattered, stipitate, at first closed and about the same diameter as the stem, gradually expanding and becoming turbinate, the margin crenate and fringed with a few delicate hairs, externally whitish or pale-orange, reaching a diameter of 1 cm.; stem rather slender and short, reaching a maximum length of 5 mm., gradually expanding above into the apothecium; hairs delicate, hyaline, septate, reaching a length of 300 μ and a diameter of 10–12 μ; asci cylindric above, tapering below into a stem-like base, reaching a length of 300–350 μ and a diameter of 20 μ; spores 1-seriate, obliquely arranged and often with the ends overlapping, ellipsoid, containing one or two (usually one) large oil-drops, at first smooth, becoming

sculptured, 12–14 × 20–25 μ; spore-sculpturing taking the form of reticulations which are often interrupted or broken; paraphyses enlarged at their apices, where they reach a diameter of 6 μ, filled with orange granules.

On soil among mosses (especially *Polytrichum*), apparently growing on the dead leaves and often hidden by the living plants.

TYPE LOCALITY: Europe.

DISTRIBUTION: New Hampshire and New York to Iowa; also in Europe.

ILLUSTRATIONS: Boud. Ic. Myc. *pl. 315;* Grevillea **22**: 108, *f. 1–6;* Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 17, f. 2;* Mycologia **6**: *pl. 142* (upper figure) and *pl. 144, f. 4.*

DOUBTFUL AND EXCLUDED SPECIES

The genus *Aleuria* is here restricted to those species in which the spores become reticulate as used by Fuckel. The genus has been used in a much broader sense by both Gillet and Boudier, so that many of the species treated as *Aleuria* by those authors will be included in the present work with the genus *Peziza* and other genera of the tribe Pezizeae.

12. **ALEURINA** Sacc. & Sydow in Sacc. Syll. Fung. **16**: 738. 1902.

Phaeopezia § *Aleurina* Sacc. Syll. Fung. **8**: 472. 1889.

Apothecia medium-sized, often reaching a diameter of several cm., cup-shaped to scutellate, fleshy or subfleshy, dark-colored, brownish-black or entirely black, fleshy or subfleshy; asci cylindric to clavate, 8-spored; spores ellipsoid, at first hyaline, becoming smoky-brown, sculptured; spore-sculpturing taking the form of regular or indistinct reticulations; paraphyses rather stout, usually filled with dark granules.

Type species, *Peziza retiderma* Cooke.

Spores broad-ellipsoid, 10 × 15–17 μ. 1. *A. atrovinosa.*
Spores ellipsoid to subfusoid, 10–12 × 22–25 μ. 2. *A. aquehongensis.*

1. **Aleurina atrovinosa** (Cooke) Seaver, comb. nov. (PLATE 8, FIG. 2.)

Peziza atrovinosa Cooke, Bull. Buffalo Soc. Nat. Sci. **2**: 288. 1875.
Peziza retiderma Cooke, Mycographia 176. 1877.
Peziza chlamydospora Ellis & Ev. Bull. Torrey Club **10**: 98. 1883.
Phaeopezia retiderma Sacc. Syll. Fung. **8**: 472. 1889.
Aleurina retiderma Sacc. & Sydow in Sacc. Syll. Fung. **16**: 739. 1902.

Apothecia gregarious or occasionally cespitose, rather deep cup-shaped, regular in form or becoming irregularly contorted

with age or from mutual pressure, at first pale-brown or smoky, becoming darker with age, reaching a diameter of 2–5 cm.; hymenium concave, darker than the outside of the apothecium, becoming almost black with age or with an olive tint from the spores which have dusted out over its surface; asci cylindric or subcylindric, tapering rather abruptly below into a stem-like base, reaching a length of 275 μ and a diameter of 12–14 μ; spores 1-seriate, ellipsoid, at first hyaline and smooth, containing one or two oil-drops, finally becoming sculptured and smoky-brown, 10 × 15–17 μ; spore-sculpturing usually consisting of one large protuberance at either end of the spore and irregular reticulate markings over the surface of the spore, with irregular lumps at various points on its surface, the ridges of the reticulations so arranged as to give the spore a somewhat striate appearance; paraphyses strongly enlarged above, where they reach a diameter of 8 μ, minutely granular within and dilutely colored.

On the ground in woods, often among mosses.

TYPE LOCALITY: New Jersey.

DISTRIBUTION: New York, New Jersey, Maine and Montana.

ILLUSTRATIONS: Cooke, Mycographia *pl. 79, f. 306;* Mycologia 6: *pl. 143* (lower figure) and *pl. 144, f. 6.*

2. **Aleurina aquehongensis** Seaver, Mycologia 6: 278. 1914.

Apothecia gregarious or scattered, sessile, discoid to shallow cup-shaped, externally slightly roughened, pale brownish-black, reaching a diameter of 1 cm.; hymenium concave, dark-brown or becoming almost black or with a slight greenish tinge which seems to be due to the spores which have been dusted out over the hymenium; asci cylindric above, tapering below into an irregular stem-like base, reaching a length of 300–350 μ and a diameter of 15–17 μ; spores 1-seriate or occasionally slightly crowded, ellipsoid to subfusoid, containing one or two oil-drops, at first smooth, becoming sculptured, pale smoky-brown, 10–12 × 22–25 μ; spore-sculpturing taking the form of irregular reticulations, the ridges of the reticulations strongly projecting, giving the spore a decidedly roughened surface; paraphyses rather strongly enlarged above, reaching a diameter of 6 μ at their apices, pale-brown.

On the ground in damp places.

TYPE LOCALITY: Staten Island, New York.

DISTRIBUTION: Known only from the type locality.

1. ALEURIA RUTILANS
2. ALEURIA RHENANA

13. **MELASTIZA** Boud. Bull. Soc. Myc. Fr. **1**: 106. 1885.

Apothecia cup-shaped or scutellate, with the hymenium bright-colored, externally clothed with flexuous or bristle-like hairs; asci 8-spored; spores hyaline, at first smooth, at maturity sculptured; spore-sculpturing assuming the form of reticulations; paraphyses filiform to clavate.

Type species, *Peziza miniata* Fuckel.

Hairs inconspicuous, closely adpressed to the sides of the apothecium. 1. *M. Charteri.*

Hairs conspicuous, extending far beyond the margin of the apothecium.

Plants tropical; spores 30–35 μ long. 2. *M. asperrima.*

Plants temperate; spores 25–30 μ long. 3. *M. pennsylvanica.*

1. **Melastiza Charteri** (W. G. Smith) Boud. Hist. Class. Discom. Eu. 64. 1907. (PLATE 45, FIG. 17.)

Peziza Charteri W. G. Smith, Gard. Chron. **1872**: 9. 1872.
Humaria miniata Fuckel, Symb. Myc. Nacht. **3**: 32. 1875.
Peziza miniata Cooke, Mycographia 71. 1876. Not *Peziza miniata* Batsch. 1786.
Humaria Charteri Rehm, Ascom. *455*. 1878.
Lachnea miniata Gill. Champ. Fr. Discom. 210. 1886.
Scutellinia miniata Kuntze, Rev. Gen. Pl. **2**: 869. 1891.
Melastiza miniata Boud. Ic. Myc. exp. pl. **1**: 16. 1905.

Apothecia scattered, gregarious, or crowded, expanding and becoming nearly plane or more rarely cup-shaped, regular in outline when young, the margin often becoming wavy or irregularly convolute with age or from mutual pressure, reaching a diameter of 1.5 cm., though often smaller, externally clothed with clumps of imperfectly developed hairs; hymenium concave or nearly plane, bright-red or almost scarlet; the excipular cells roundish or angular, the cell walls pale-brown; hairs usually clinging together in fascicles, flexuous, thin-walled, with blunt ends, 3–4-septate, pale-brown, reaching a length of 150–175 μ and a diameter of 10–12 μ; asci cylindric or subcylindric, tapering gradually below, reaching a length of 300 μ and a diameter of 12–14 μ; spores 1-seriate, parallel with the ascus or more rarely diagonally disposed, ellipsoid, hyaline, usually containing two oil-drops, finally becoming sculptured, 10–13 × 17–20 μ; spore-sculpturing assuming the form of reticulations, the reticulations often indistinct and broken in young specimens, and often giving rise to an apiculus at either end; paraphyses enlarged above, where they reach a diameter of 6–8 μ.

On bare soil or among mosses.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to Colorado; also in Europe.

ILLUSTRATIONS: Boud. Ic. Myc. *pl. 386;* Cooke, Mycographia *pl. 33, f. 127;* Gardner's Chron. **1872**: 9, *f. 1;* Grevillea **1**: *pl. 8, f. 1, 2;* Gill. Champ. Fr. Discom. (suites); Pat. Tab. Fung. *f. 276.*

This species has been frequently collected about New York on bare soil in open woods. As this goes to press, specimens of the species have been received from Winnipeg, Canada, with the statement that it is collected there every autumn in abundance on sawdust in an old ice-house after the ice has been removed. The material was collected by A. H. R. Buller and G. R. Bisby.

2. **Melastiza asperrima** (Ellis & Ev.) Seaver, comb. nov.

Lachnea barbata Massee, Jour. Bot. **30**: 163. 1892. Not *L. barbata* (Kunze) Gill. 1879.

Lachnea asperrima Ellis & Ev. (in herb.).

Apothecia scattered or gregarious, scutellate, the margins slightly elevated, regular in form or with the margins wavy, externally sparingly clothed with hairs, reaching a diameter of 5–8 mm.; hymenium slightly concave, bright-red; hairs very long and projecting far out about the margin of the apothecium, dark reddish-brown, septate, thick-walled, gradually tapering above into a bristle-like apex, reaching a length of 2 mm. and a diameter of 35 μ near the base; asci cylindric or subcylindric, reaching a length of 250 μ and a diameter of 17–20 μ; spores 1-seriate, ellipsoid, containing one large oil-drop, 15–17 $\times$ 30–35 μ, becoming sculptured; spore-sculpturing assuming the form of very regular, usually hexagonal reticulations, the reticulations consisting of ridges which extend about 2 μ beyond the periphery of the spore, strongly roughening its surface, the meshes of the reticulations about 3 or 4 μ in diameter; paraphyses slender below, rather strongly thickened above.

On rotten wood.

TYPE LOCALITY: Nicaragua.

DISTRIBUTION: Nicaragua to Mexico and the West Indies.

ILLUSTRATION: Jour. Bot. **30**: *pl. 321, f. 5–8.*

3. **Melastiza pennsylvanica** Seaver, sp. nov.

Apothecia scattered or sessile, scutellate, the margin slightly elevated and sparingly clothed with long hairs, reaching a

diameter of 5–8 mm.; hymenium slightly concave, bright-red; hairs dark-brown, thick-walled, septate, reaching a length of more than 1 mm. and a diameter of 35–40 μ near the base, gradually contracted below and often forked at the base, tapering above into a bristle-like apex; asci cylindric or subcylindric above, reaching a length of 250 μ and a diameter of 20 μ; spores 1-seriate, parallel with the ascus or diagonally disposed, at first smooth, becoming sculptured at an early stage, 14–17 × 25–30 μ; spore-sculpturing taking the form of distinct and regular reticulations, the ridges of the reticulations 1–2 μ deep and giving to the spore surface a very rough appearance; paraphyses about 2 μ in diameter below, rather strongly enlarged above.

On old wood.

Type collected in deciduous woods along the Youghiogheny River, Pennsylvania, by W. A. Murrill (Herb. The New York Botanical Garden).

DISTRIBUTION: Pennsylvania.

14. **PSILOPEZIA** Berk. London Jour. Bot. II. 6: 325. 1847.

Fleischhakia Rab. Hedwigia **17**: 114. 1878.
Peltidium Kalchbr. Hedwigia **2**: 58. 1862. Not *Peltidium* Zollik. 1820.

Apothecia medium-sized, usually not exceeding 1 cm. in diameter, becoming flattened or subdiscoid, closely adhering to the substratum and usually surrounded by a sterile border, occasionally with the margin free and slightly elevated; hymenium slightly concave, plane or more often convex, usually dark-colored, brown or brownish, more rarely bright-colored, yellow or orange; asci cylindric or subcylindric, 8-spored, usually spirally marked; spores ellipsoid, smooth or more rarely sculptured, hyaline or subhyaline; paraphyses stout.

Type species, *Psilopezia nummularia* Berk.

Apothecia dark-brown or reddish-brown.
  Spores smooth.
    Spores 12 × 20–25 μ. — 1. *P. nummularia.*
    Spores 15–17 × 22–25 μ. — 2. *P. hydrophila.*
    Spores 15–17 × 30–36 μ. — 3. *P. deligata.*
  Spores becoming rough. — 4. *P. trachyspora.*
Apothecia yellow or orange.
  Spores 7 × 12–14 μ. — 5. *P. flavida.*
  Spores 12–13 × 20–24 μ. — 6. *P. aquatica.*

1. **Psilopezia nummularia** Berk. London Jour. Bot. **6**: 325. 1847.

Peziza Babingtonii Berk. & Br. Ann. Mag. Nat. Hist. II. **7**: 179. 1851.
Psilopezia myrothecioides Berk. & Br. Ann. Mag. Nat. Hist. IV. **15**: 39. 1875.
Fleischhakia rhizinoides Rab. Hedwigia **17**: 114. 1878.
Peziza depressa Phill.; Cooke, Mycographia 233. 1879. Not *P. depressa* Pers. 1796.
Humaria depressa Sacc. Syll. Fung. **8**: 145. 1889.
Rhizina myrothecioides Massee, Brit. Fungus-Fl. **4**: 455. 1895.
Rhizina Babingtonii Massee, Brit. Fungus-Fl. **4**: 455. 1895.
Peziza nummularia Morgan, Jour. Myc. **8**: 190. 1902.

Apothecia gregarious, becoming discoid or subdiscoid and closely adhering to the substratum to the extreme margin and surrounded by a whitish mycelial growth, reaching a diameter of 1–5 mm. or rarely 1 cm.; hymenium plane or slightly convex, reddish-brown to brown, almost black in dried specimens, more or less velvety from the protruding paraphyses; asci cylindric above, tapering below and marked with a distinct spiral which appears to be due to thickening or possibly to a twisting of the ascus, reaching a length of 200–250 μ and a diameter of 15–18 μ; spores 1-seriate at maturity, ellipsoid, containing one or two oil-drops, hyaline or very faintly yellowish, about 12 $\times$ 20–25 μ; paraphyses stout, much thickened above and containing coarse brown granules, pale below, reaching a diameter of 10 μ at their apices.

On rotten wood.

TYPE LOCALITY: Great Britain.

DISTRIBUTION: Ontario to New Hampshire and Greenland?; also in Europe.

ILLUSTRATIONS: Ann. Mag. Nat. Hist. IV. **15**: *pl. 2, f. 5;* Cooke, Mycographia *pl. 110, f. 392;* Phill. Brit. Discom. *pl. 5, f. 23;* Rab. Krypt.-Fl. 1^3: 1135, *f. 1–3;* Jour. Linn. Soc. **31**: *pl. 16, f. 13.*

2. **Psilopezia hydrophila** (Peck) Seaver, comb. nov.

Peziza hydrophila Peck, Ann. Rep. N. Y. State Mus. **34**: 51. 1881.
Humaria hydrophila Sacc. Syll. Fung. **8**: 140. 1889.
?Peziza rivularis Clements, Bot. Surv. Nebr. **3**: 8. 1894.

Apothecia scattered, sessile, becoming expanded, at maturity plane or even convex; hymenium reddish-brown when moist, black when dry, reaching a diameter of 2–5 mm.; asci cylindric above, tapering below; spores 1-seriate, ellipsoid, generally con-

taining two oil-drops, reaching a diameter of 15–17 μ and a length of 22–25 μ; paraphyses numerous, thickened above, compact and adhering together at their apices.

On decaying wood, lying in the water.

TYPE LOCALITY: Adirondack Mountains, New York.

DISTRIBUTION: New York and (? Nebraska).

3. **Psilopezia deligata** (Peck) Seaver, comb. nov.

Bulgaria deligata Peck, Ann. Rep. N. Y. State Mus. **32**: 49. 1879. Not *Peziza deligata* Peck.

Peziza leucobasis Peck, Bull. N. Y. State Mus. **2**: 20. 1887.

Pyronema leucobasis Sacc. Syll. Fung. **8**: 110. 1889.

Apothecia sessile, closely adhering to the substratum and surrounded by a whitish or yellowish border of superficial mycelium, becoming almost convex-hemispheric, reaching a diameter of 3 mm.; hymenium covering all of the exposed surface of the apothecium, convex, brownish-black with a tinge of olive; asci cylindric to subclavate, rather broad; spores 1-seriate, broad-ellipsoid, thick-walled, smooth, slightly yellowish, usually containing two large oil-drops, reaching a diameter of 15–17 μ and a length of 30–36 μ; paraphyses enlarged above, reaching a diameter of 10 μ at their apices, yellowish or brownish but not usually containing the conspicuous granules characteristic of the other species.

On wet hemlock wood and the bark of an unknown tree.

TYPE LOCALITY: Catskill Mountains, New York.

DISTRIBUTION: New York and New Hampshire.

ILLUSTRATION: Bull. N. Y. State Mus. **2**: *pl. 2, f. 1–3.*

4. **Psilopezia trachyspora** Ellis & Ev. Erythea **1**: 200. 1893.

Apothecia gregarious, becoming subdiscoid, adhering closely to the substratum except the extreme margin which is free and slightly elevated, externally pale-brown and becoming slightly wrinkled at least in dried specimens, reaching a diameter of 1–2 cm.; hymenium dark-brown; asci cylindric above, gradually tapering below, reaching a length of 300–325 μ and a diameter of 15–18 μ, becoming spirally marked, the spiral markings extending the whole length of young asci but in older asci restricted to the lower extremity only; spores 1-seriate or slightly crowded, ellipsoid, at first smooth and containing two oil-drops, becoming sculptured, 13–15 $\times$ 25–28 μ; spore-sculpturing taking the form

of prominent wart-like protuberances; paraphyses stout, much enlarged above, filled with dark-brown coloring matter.

On decaying cedar wood.

TYPE LOCALITY: North Bend, King County, Washington.

DISTRIBUTION: Known only from the type locality.

5. **Psilopezia flavida** Berk. & Curt. Grevillea **4**: 1. 1875.

Apothecia crowded and occasionally confluent, orbicular in form when solitary, reaching a diameter of 3–6 mm., the margin free and often raised when mature; hymenium concave, plane or convex, yellow; asci narrowly cylindric, reaching a length of 120 μ and a diameter of 7–8 μ; spores 1-seriate, hyaline, smooth, ellipsoid, 7 $\mu \times$ 12–14 μ, containing two oil-drops; paraphyses stout, scarcely thickened above, not longer than the asci.

On rotten wood of *Quercus alba*.

TYPE LOCALITY: Alabama.

DISTRIBUTION: Alabama.

ILLUSTRATION: Jour. Linn. Soc. **31**: *pl. 16, f. 10.*

6. **Psilopezia aquatica** (DC.) Rehm; Sacc. & D. Sacc. in Sacc. Syll. Fung. **18**: 12. 1906.

Peziza aquatica DC. in Lam. Encyc. **5**: 216. 1804.
Peltidium Oocardii Kalch. Hedwigia **2**: 58. 1862.
Peziza Oocardii lignaria P. Karst. Not. Fauna Fl. Fenn. **10**: 128. 1869.
Peziza Oocardii Karst. Myc. Fenn. **1**: 84. 1871.
Peziza lechithina Cooke, Grevillea **4**: 110. 1876.
Peltidium lignarium Karst.; Hazsl. in Oesterr. Bot. Zeits. **32**: 7. 1882.
Peltidium Cookei Hazsl. in Oesterr. Bot. Zeits. **32**: 7. 1882.
Humaria lechithina Sacc. Syll. Fung. **8**: 126. 1889.
Humaria Oocardii Sacc. Syll. Fung. **8**: 144. 1889.
Humaria aquatica Rehm in Rab. Krypt.-Fl. 1^3: 954. 1896.
Psilopezia Oocardii Sacc. & D. Sacc. in Sacc. Syll. Fung. **18**: 11. 1906.

Apothecia gregarious, sessile, becoming discoid and closely adhering to the substratum, surrounded about the margin by a white radiating mycelium, reaching a diameter of 3–5 mm.; hymenium plane or slightly convex, bright-orange or becoming dingy with age, blackish when dry; asci cylindric above, gradually tapering below, reaching a length of 275 μ and a diameter of 16–20 μ, becoming strongly spirally twisted; spores ellipsoid, 1-seriate, usually containing two oil-drops or occasionally with numerous smaller ones, hyaline or faintly yellowish, 12–13 $\times$ 20–24 μ, smooth or becoming minutely sculptured; spore-sculpturing, when present, consisting of very minute granules;

paraphyses rather strongly enlarged above, reaching a diameter of 7–8 μ.

On stones and water-soaked logs.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York and New Jersey to Minnesota and Utah; also in Europe.

ILLUSTRATIONS: Hedwigia **2**: *pl. 10, f. 3;* Cooke, Mycographia *pl. 12, f. 47* and *pl. 23, f. 89;* Massee, Brit. Fungus-Fl. **4**: 290, *f. 10–13;* Fest. Asch. 483, *f. 1–6.*

DOUBTFUL AND EXCLUDED SPECIES

Psilopezia mirabilis Berk. & Curt.; Berk. Jour. Linn. Soc. **10**: 364. 1869. The species, which is very imperfectly described, is said to occur on twigs and to look like a *Corticium.*

15. **PYRONEMA** Carus, Nov. Act. Acad. Nat. Cur. **17**: 370. 1835.

Vegetative mycelium superficial, hyaline (white in mass), thin-walled, septate, branched, the branches proceeding almost at right angles to the main hypha, frequently anastomosing, filled with conspicuous vacuoles, giving rise to paired sex organs; sex organs occurring in clusters of several pairs each, each pair consisting of a clavate antheridium and an inflated oögonium surmounted by a slender trichogyne, which at a later stage usually fuses with the oögonium, each cluster of sex organs finally giving rise to a compound apothecium; apothecia open from the first, never enclosed by the excipulum; asci cylindric, 8-spored; spores ellipsoid, hyaline, smooth; paraphyses filiform.

Type species, *Pyronema Marianum* Carus?

1. **Pyronema omphalodes** (Bull.) Fuckel, Symb. Myc. 319. 1869. (PLATE 10.)

Peziza omphalodes Bull. Herb. Fr. *pl. 485, f. 1.* 1790. Hist. Champ. 264. 1791.
Peziza confluens Pers. Obs. Myc. **2**: 81. 1799.
Pyronema Marianum Carus, Nov. Act. Acad. Nat. Cur. **17**: 370. 1835.
Pyronema confluens Tul. Fung. Carp. **3**: 197. 1865.
Aleuria omphalodes Gill. Champ. Fr. Discom. 48. 1879.
?Pyronema aurantiorubrum Sacc. Syll. Fung. **8**: 108. 1889.
Tapesia omphalodes Quél. Ench. Fung. 291. 1886.
Humaria omphalodes Massee, Brit. Fungus-Fl. **4**: 410. 1895.

Apothecia small, not usually exceeding 1–2 mm. in diameter, soon becoming confluent and forming congested masses several

cm. in diameter; apothecial masses circular or irregular, often interrupted, pale-orange, surrounded by a dense superficial white mycelial growth; hymenium usually convex, the color varying greatly with conditions, sometimes only slightly yellowish, to bright-orange, occasionally with a purplish tinge; asci cylindric or subcylindric, 150 μ long and 10–14 μ in diameter; spores 1-seriate, ellipsoid, smooth, hyaline, 5–8 $\times$ 10–13 μ; paraphyses rather stout, very slightly enlarged above, reaching a diameter of 6–7 μ at their apices, filled with orange granules.

Occurring in nature on burned places; in greenhouses on steamed soil or soil which has been sterilized by dry heat.

TYPE LOCALITY: Europe.

DISTRIBUTION: Throughout North America, Europe and Australia; probably world-wide.

ILLUSTRATIONS: Bull. Herb. Fr. *pl. 485, f. 1;* Boud. Ic. Myc. *pl. 419;* Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 7, f. 1;* Cooke, Mycographia *pl. 17, f. 65;* Gill. Champ. Fr. Discom. *pl. 51;* E. & P. Nat. Pfl. 1^1: 177, *f. 145;* Mycologia **1**: *pl. 9–12;* Mycologia **2**: *pl. 26;* Tul. Fung. Carp. **3**: *pl. 17, f. 11, 12;* Pat. Tab. Fung. *f. 287;* Cooke, Austr. Fung. *pl. 19, f. 149;* Rab. Krypt.-Fl. 1^3: 919, *f. 1–5;* Phill. Brit. Discom. *pl. 5, f. 22;* Ann. Sci. Nat. V. **6**: *pl. 11, f. 10–13* and *pl. 12;* Heidler, Pfl. Marienbad, *pl. 2.*

EXSICCATI: Fungi Columb. *1308.*

Pyronema confluens inigneum W. H. Brown, Am. Jour. Bot. **2**: 297. 1915.

This variety according to Brown seems to differ from the species described above through the fact that the plants will develop on soil or pots which have not been sterilized or heated, a condition which has been shown to be absolutely essential to the development of the species. The variety is also said to differ from the species in the entire absence of fusion between the antheridium and trichogyne.

EXCLUDED SPECIES

Pyronema leucobasis (Peck) Sacc. Syll. Fung. **8**: 110. 1889. The only character about this plant which suggests *Pyronema* is the superficial mycelium which surrounds the apothecia, and it was doubtless due to this resemblance that Saccardo placed the species in that genus. The species is a true *Psilopezia*. The superficial resemblance of the apothecia in *Psilopezia* to those of *Pyronema* was mentioned by Berkeley when he proposed the genus *Psilopezia*.



PYRONEMA OMPHALODES

16. **ASCOPHANUS** Boud. Ann. Sci. Nat. V. **10**: 241. 1869.

Coprobia Boud. Bull. Soc. Myc. Fr. **1**: 107. 1885.

Plants coprophilous (occurring on the dung of animals or stercoriated mud); apothecia small, rarely exceeding 1–2 mm. in diameter and often less than 1 mm., sessile or short-stipitate, scutellate, externally naked (not clothed with hairs) and smooth or minutely rough, usually bright-colored or more rarely brownish or blackish; hymenium concave, plane or more often strongly convex, leaving the margin indistinct, usually similar in color to the outside of the apothecium; asci cylindric, clavate or ovoid, 4–8-spored; spores 1–2-seriate, hyaline or subhyaline, ellipsoid, smooth or minutely rough, in one species apiculate; paraphyses slender below, their ends clavate to subglobose, hyaline or colored.

Type species, *Peziza subfusca* Crouan.

Spores large, 25 μ or more in length.
- Apothecia white or whitish.
  - Spores remaining permanently smooth.
    - Spores apiculate. — 1. *A. Holmskjoldii.*
    - Spores not apiculate. — 2. *A. isabellinus.*
  - Spores becoming rough.
    - Spores 18–20 × 25–30 μ; spore-sculpturing wart-like. — 3. *A. sarcobius.*
    - Spores 23–25 × 33–38 μ; spore-sculpturing like tack-heads. — 4. *A. bermudensis.*
- Apothecia ochraceous-carneous. — 5. *A. vicinus.*

Spores medium-sized, 10–25 μ in length.
- Apothecia white.
  - Asci cylindric; spores 1-seriate. — 6. *A. lacteus.*
  - Asci clavate to subovoid; spores 2-seriate. — 7. *A. argenteus.*
- Apothecia colored, reddish, brown, or lilac.
  - Apothecia reddish or orange.
    - Spores 6–8 × 10–12 μ. — 8. *A. Aurora.*
    - Spores 10–16 × 15–26 μ.
      - Asci clavate; spores 2-seriate. — 9. *A. carneus.*
      - Asci cylindric or subcylindric; spores 1-seriate. — 10. *A. granulatus.*
  - Apothecia yellowish to ochraceous.
    - Spores 7–8 × 10–13 μ. — 11. *A. granuliformis.*
    - Spores 8–10 × 15–20 μ. — 12. *A. ochraceus.*
    - Spores 10–11 × 5–6 μ. — 13. *A. gallinaceus.*
  - Apothecia cinereous or blackish.
    - Spores 10–12 × 20–24 μ. — 14. *A. cinereus.*
    - Spores 7–8 × 15–17 μ. — 15. *A. tetraonalis.*
  - Apothecia lilac. — 16. *A. lilacinus.*

Spores small, less than 10 μ long.
  Apothecia cinereous. 17. *A. cinerellus.*
  Apothecia whitish to yellowish to brown.
    Paraphyses globose at their apices. 18. *A. microsporus.*
    Paraphyses only slightly enlarged above.
      Apothecia externally brown; hymenium pallid. 19. *A. Cesatii.*
      Apothecia whitish or yellowish. 20. *A. glaucellus.*

1. **Ascophanus Holmskjoldii** E. C. Hansen, Vidensk. Medd. 1876: 290. Jan. 1877. (PLATE 11, FIG. 8.)

Ascobolus incanus Phill. Grevillea **5**: 117. March, 1877.
Ascophanus incanus Sacc. Syll. Fung. **8**: 529. 1889.

Apothecia scattered or gregarious, at first turbinate, becoming expanded and reaching a diameter of about 1 mm., grayish; hymenium plane or nearly so, of the same color as the outside of the apothecium, roughened by the large protruding asci; asci broad-clavate, attenuated below into a stem-like base, reaching a length of 250 μ and a diameter of 40 μ; spores 2-seriate or irregularly disposed, ellipsoid, often with a small knob or apiculus at one or both ends, hyaline, granular within and often surrounded by an accumulation of granules which give to the outside of the spore a granular appearance but apparently with no true sculpturing, 14–16 × 30–36 μ; paraphyses slender, scarcely enlarged above, hyaline or subhyaline.

On cow dung.

TYPE LOCALITY: Denmark.

DISTRIBUTION: New York, California; also in Europe.

ILLUSTRATIONS: Grevillea **5**: *pl. 88, f. 10;* Vidensk. Medd. 1876: *pl. 6, f. 1–8;* Rab. Krypt.-Fl. 1^3: 1079, *f. 1–2;* Bull. Soc. Myc. Fr. **34**: *pl. 3, f. 1.*

2. **Ascophanus isabellinus** Clements, Bot. Surv. Nebr. **5**: 9. 1901. (PLATE 11, FIG. 13.)

Apothecia at first nearly doliiform, becoming expanded, white, later becoming dirty-pallid, reaching a diameter of 1–3 mm.; hymenium becoming plane or slightly convex; asci broad-clavate, attenuated below into a stem-like base, reaching a length of 250 μ and a diameter of 30–40 μ, 8-spored; spores 2-seriate, ellipsoid, hyaline, smooth, with a wall 2–3 μ thick, reaching a size of 12–13 × 27–30 μ; paraphyses linear and clavate, the latter reaching a diameter of 3 μ at their apices.

On stercoriated mud.

TYPE LOCALITY: Otawanie woods, Nebraska.

DISTRIBUTION: New York to Nebraska.

3. **Ascophanus sarcobius** Boud. Bull. Soc. Myc. Fr. **18**: 143. 1902. (PLATE 11, FIG. 4.)

Apothecia gregarious, at first subglobose, smooth, white, gradually expanding, at maturity with the hymenium convex and margin indistinct, reaching a diameter of 1–2 mm.; asci clavate, about 250–300 × 35–38 μ, 8-spored; spores 2-seriate or irregularly crowded near the end of the ascus, ellipsoid, at first smooth, becoming sculptured hyaline, about 18–20 × 25–30 μ; spore-sculpturing taking the form of minute warts; paraphyses strongly enlarged above, septate, hyaline, reaching a diameter of 10–12 μ at their apices.

On cow dung.

TYPE LOCALITY: France.

DISTRIBUTION: Bermuda; also in Europe.

ILLUSTRATIONS: Bull. Soc. Myc. Fr. **18**: *pl. 8, f. 2;* Boud. Ic. Myc. *pl. 413.*

4. **Ascophanus bermudensis** Seaver, Mem. N. Y. Bot. Gard. 6: 505. 1916. (PLATE 11, FIG. 2.)

Apothecia gregarious or scattered, at first subglobose, expanding and becoming subdiscoid, reaching a diameter of 1–2 mm., white or more often with a delicate pinkish tint; hymenium at first slightly concave, becoming plane, finally convex, roughened by the protruding asci, similar in color to the outside of the apothecium; asci clavate, reaching a length of 325 μ and a diameter of 35–40 μ, 8-spored; spores 1-seriate or partially 2-seriate, or occasionally irregularly disposed, at first smooth, becoming rough, 23–25 × 33–38 μ; spore-roughenings assuming the form of scattered protuberances which are enlarged outwards, giving them the appearance of protruding tack- or nail-heads, 2–3 μ long and 2 μ broad; paraphyses strongly enlarged above, hyaline.

On cow dung.

Type collected on cow dung near Harrington Sound, Bermuda, December, 1912 (Herb. The New York Botanical Garden).

DISTRIBUTION: Known only from the type locality.

5. **Ascophanus vicinus** Boud. Ann. Sci. Nat. V. **10**: 246. 1869.

Apothecia scattered or more rarely aggregated, becoming discoid, reaching a diameter of 0.5–1 mm., ochraceous-carneous; hymenium plane or convex, similar in color to the outside of the apothecium, roughened by the protruding asci; asci broad-clavate, attenuated at the base, 8-spored; spores 2-seriate or irregularly disposed, hyaline, 20–22 × 30–32 μ; paraphyses clavate at their apices, septate, hyaline.

On cow dung.

TYPE LOCALITY: France.

DISTRIBUTION: New Jersey; also in Europe.

ILLUSTRATION: Ann. Sci. Nat. V. **10**: *pl. 11, f. 33.*

6. **Ascophanus lacteus** (Cooke & Phill.) Sacc. Syll. Fung. **8**: 528. 1889. (PLATE 11, FIG. 1.)

Ascobolus lacteus Cooke & Phill. Grevillea **5**: 119. 1876.

Apothecia small, not exceeding 0.3 mm. in diameter, discoid, tapering below, white or with a very delicate pinkish tinge; hymenium concave, minutely roughened by the protruding asci; the excipular cells irregularly polygonal, hyaline or subhyaline, reaching a diameter of 12–15 μ; asci cylindric to clavate, gradually tapering below into a stem-like base, reaching a length of 60–75 μ and a diameter of 12–15 μ, 8-spored; spores usually 1-seriate or more rarely crowded, ellipsoid, smooth, about 7 × 10 μ; paraphyses about 1 μ thick below, enlarged above, branched and curved at their apices.

On dung of cows and horses.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York; also in Europe.

ILLUSTRATIONS: Heimerl, Niederöst. Ascoboleen *f. 9, 10.*

7. **Ascophanus argenteus** (Curr.) Boud. Ann. Sci. Nat. V. **10**: 245. 1869. (PLATE 11, FIG. 5.)

Ascobolus argenteus Currey, Trans. Linn. Soc. **24**: 496. 1864.

Apothecia gregarious, minute, not exceeding 0.2–0.3 mm. in diameter, sessile, discoid, translucent and silvery-white; hymenium nearly plane or convex, minutely roughened by the protruding asci; excipular cells hyaline, irregularly polygonal and 7–10 μ in diameter; asci broadly clavate to subovoid and abruptly tapering below into a stem-like base, reaching a length of 50–60 μ and a diameter of 20 μ, 8-spored, tapering into a short

stem-like base; spores irregularly 2-seriate, ellipsoid, hyaline, smooth, 6–7 × 10–12 μ; paraphyses slender, septate, hyaline, clavate at their apices.

On cow and horse dung.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to Colorado and Porto Rico; also in Europe.

ILLUSTRATIONS: Trans. Linn. Soc. **24**: *pl. 51, f. 21;* Ann. Mag. Nat. Hist. III. **15**: *pl. 17, f. 32;* Ann. Sci. Nat. V. **10**: *pl. 11, f. 32;* Jour. Bot. **2**: 152, *f. 6.*

8. **Ascophanus Aurora** (Crouan) Boud. Ann. Sci. Nat. V. **10**: 248. 1869. (PLATE 11, FIG. 7.)

Peziza Aurora Crouan, Fl. Finist. 53. 1867.
Aleuria Aurora Gill. Champ. Fr. Discom. 54. 1879.

Apothecia scattered, sessile, at first subglobose, becoming subdiscoid, reaching a diameter of 0.5–1.5 mm., externally smooth, bright-orange; hymenium at first concave, becoming plane, similar in color to the outside of the apothecium but a little brighter, roughened by the protruding asci; asci cylindric to clavate, 8-spored; spores 1-seriate or becoming irregularly 2-seriate, ellipsoid, hyaline, 6–7 × 10–12 μ; paraphyses slender, septate, simple or branched, usually curved at their apices, scarcely enlarged above, filled with orange granules.

On dung of various kinds.

TYPE LOCALITY: Europe.

DISTRIBUTION: Delaware and Nebraska; also in Europe.

ILLUSTRATION: Ann. Sci. Nat. V. **10**: *pl. 11, f. 36.*

9. **Ascophanus carneus** (Pers.) Boud. Ann. Sci. Nat. V. **10**: 250. 1869.

Ascobolus carneus Pers. Syn. Fung. 676. 1801.
Ascobolus saccharinus Berk. & Curt.; Berk. Outl. Brit. Fungol. 374. 1860.
Ascophanus saccharinus Boud. Ann. Sci. Nat. V. **10**: 251. 1869.
Peziza humosoides Peck, Ann. Rep. N. Y. State Mus. **32**: 46. 1879.
Ascophanus humosoides Peck, Bull. N. Y. State Mus. **2**: 22. 1887.
Ascophanus carneus saccharinus Phill. Brit. Discom. 309. 1887.
Pyronema carneum Schröt. Krypt.-Fl. Schles. **3**^{2}: 34. 1893.
Ascophanus carneus difformis Clements, Bot. Surv. Nebr. **5**: 22. 1901. Not *Peziza difformis* Karst.

Apothecia gregarious or occasionally crowded in clusters of variable size, minute, usually not exceeding 1 mm. in diameter

or rarely as large as 2 mm., flesh-colored, the color becoming brighter in dried specimens; hymenium at first concave, becoming convex, similar in color to the outside of the apothecium but a little brighter, roughened by the protruding asci; excipulum giving rise above to a palisade of mycelium the loose ends of which form a very delicate fringe-like border, the ends of the peridial mycelium reaching a diameter of 12–15 μ; asci rather broad-clavate, reaching a length of 200–300 μ and a diameter of 25–40 μ, tapering below into a stem-like base, 8-spored; spores 2-seriate above or becoming irregularly crowded, ellipsoid, hyaline, at first smooth, becoming sculptured, 11–12 $\times$ 17–20 μ, or rarely 14–16 $\times$ 24–26 μ; spore-sculpturing consisting of very minute warts barely roughening the surface of the spore; paraphyses septate, stout, enlarged at their apices, where they reach a diameter of 7–8 μ, densely filled with orange granules.

On dung of various kinds.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to North Dakota, Florida, Colorado, Bermuda and Porto Rico; also in Europe.

ILLUSTRATIONS: Ann. Mag. Nat. Hist. III. **15**: *pl. 14, f. 5;* Ann. Sci. Nat. V. **10**: *pl. 12, f. 38–40;* Rab. Krypt.-Fl. 1^3: 1080, *f. 1–5;* Pat. Tab. Fung. *f. 379;* Trans. Linn. Soc. **24**: *pl. 51, f. 20.*

A specimen grown on dung from Porto Rico shows asci 40 $\times$ 300 μ and spores 14 $\times$ 26 μ; otherwise it appears to be typical of *A. carneus.*

10. **Ascophanus granulatus** (Bull.) Speg. Michelia 1: 235. 1879. (PLATE 11, FIG. 12.)

?Peziza scabra Müll. Fl. Dan. **11**: 7. 1775.
Peziza granulata Bull. Herb. Fr. *pl. 438, f. 3.* 1789.
Peziza granulosa Pers. Syn. Fung. 667. 1801.
Ascobolus granulatus Fuckel, Symb. Myc. 287. 1869.
Aleuria granulata Gill. Champ. Fr. Discom. 56. 1879.
Humaria granulata Quél. Ench. Fung. 290. 1886.
Coprobia granulata Boud. Hist. Class. Discom. Eu. 69. 1907.

Apothecia gregarious or crowded, at first closed and subglobose, expanding and becoming scutellate to subdiscoid, externally pale-orange, coarsely granulose, reaching a diameter of 1–3 mm., color fading in dried specimens to nearly white; hymenium slightly concave to plane, orange, brighter than the outside of the apothecium; asci cylindric or subcylindric, gradu-

ally tapering below, reaching a length of 175–200 μ and a diameter of 15–17 μ, 8-spored; spores 1-seriate, diagonally arranged in the ascus, narrow-ellipsoid, smooth, 7–9 $\times$ 14–18 μ; paraphyses stout, strongly thickened above, reaching a diameter almost equal to that of the ascus.

On cow dung.

TYPE LOCALITY: Europe.

DISTRIBUTION: Connecticut to Iowa, South Carolina and Mexico; also in Europe.

ILLUSTRATIONS: Bull. Herb. Fr. *pl. 438, f. 3;* Gill. Champ. Fr. Discom. *pl. 47, f. 1;* Cooke, Mycographia *pl. 15, f. 59;* ?Fl. Dan. *pl. 655, f. 2.*

EXSICCATI: Rav. Fungi Am. Exsicc. *630;* N. Am. Fungi *2037.*

11. **Ascophanus granuliformis** (Crouan) Boud. Ann. Sci. Nat. V. **10**: 245. 1869. (PLATE 11, FIG. 10.)

Ascobolus granuliformis Crouan, Ann. Sci. Nat. IV. **10**: 195. 1858.

Apothecia minute, usually less than 1 mm. in diameter, sessile, at first globose, becoming discoid; hymenium pale yellow-ochraceous; asci broad-clavate to ovoid, tapering below, 8-spored; spores hyaline, smooth, ellipsoid, 7–8 $\times$ 10–13 μ; paraphyses either simple or branched, hyaline, septate, pear-shaped at their apices.

On cow dung.

TYPE LOCALITY: France.

DISTRIBUTION: Iowa and Colorado; also in Europe.

ILLUSTRATIONS: Ann. Sci. Nat. IV. **10**: *pl. 13, f. F;* Ann. Sci. Nat. V. **10**: *pl. 10, f. 31;* Ann. Mag. Nat. Hist. III. **15**: *pl. 17, f. 31;* Jour. Bot. **2**: 152, *f. 5;* Pat. Tab. Fung. *f. 173.*

EXSICCATI: Clements, Crypt. Form. Colo. *300.*

12. **Ascophanus ochraceus** (Crouan) Boud. Ann. Sci. Nat. V. **10**: 247. 1869.

Ascobolus ochraceus Crouan, Fl. Finist. 57. 1867.
?Peziza cremoricolor Berk. Grevillea **3**: 151. 1875.
Ascophanus subgranuliformis Rehm; Voss, Verh. Zool.-Bot. Ges. Wien **37**: 224. 1887.
Ascophanus flavus Karst. Medd. Soc. Fauna Fl. Fenn. **16**: 105. 1889.
?Humaria cremoricolor Sacc. Syll. Fung. **8**: 136. 1889.

Apothecia thickly gregarious, sessile, at first globose or subglobose, expanding and becoming subdiscoid, pale-yellow, ex-

ternally smooth, reaching a diameter of 0.3–1 mm.; hymenium plane or a little concave, finally convex, the color a little brighter than that of the outside of the apothecium, roughened by the protruding asci; asci cylindric above, slightly tapering below, reaching a diameter of 12–14 μ, 8-spored; spores 1-seriate, parallel with the ascus or diagonally disposed, more rarely crowded, ellipsoid, smooth, subhyaline or faintly yellowish, 8–10 $\times$ 15–20 μ; paraphyses slender, enlarged above, where they reach a diameter of 5–6 μ.

On dung of various kinds.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to West Virginia and Bermuda; also in Europe.

ILLUSTRATIONS: Verh. Zool.-Bot. Ges. Wien **37**: *pl. 5, f. 7;* Ann. Sci. Nat. V. **10**: *pl. 11, f. 34;* E. & P. Nat. Pfl. 1^1: 189, *f. 152, D.*

13. **Ascophanus gallinaceus** (Peck) Seaver, comb. nov.

Peziza gallinacea Peck, Ann. Rep. N. Y. State Mus. **31**: 46. 1879.
Humaria gallinacea Sacc. Syll. Fung. **8**: 135. 1889.

Apothecia gregarious or scattered, yellowish to ochraceous, sessile or with the base slightly projected below, externally slightly furfuraceous, the margin often wavy or irregular, reaching a diameter of 2–3 mm.; hymenium smooth, similar in color to the outside of the apothecium; asci cylindric, reaching a length of 150 μ and a diameter of 10 μ, 8-spored; spores ellipsoid, smooth, 1-seriate, occupying the upper part of the ascus only, about 5–6 $\times$ 10–11 μ; paraphyses slender, enlarged above at their apices.

On partridge dung.

TYPE LOCALITY: Oneida, New York.

DISTRIBUTION: Known only from the type locality.

14. **Ascophanus cinereus** (Crouan) Boud. Ann. Sci. Nat. V. **10**: 249. 1869. (PLATE 11, FIG. 11.)

Ascobolus cinereus Crouan, Ann. Sci. Nat. IV. **10**: 194. 1858.
Peziza cinerea Karst. Myc. Fenn. **1**: 59. 1871.
Ascophanus crustaceus Starb. Bot. Not. **1898**: 216. 1898.

Apothecia gregarious or closely crowded, at first subglobose or pyriform, becoming short-cylindric, finally expanding to discoid or subdiscoid and narrowed below into a stem-like base,

at first grayish or cinereous, but at maturity darker and appearing almost black and so closely resembling the substratum in color that they are detected with difficulty, reaching an extreme diameter of 2 mm.; hymenium becoming plane, finally strongly convex, roughened by the protruding asci, often surrounded by a slightly elevated margin; excipular cells rounded or angular, the walls smoky-brown, variable in size but reaching a diameter of 12–15 μ, giving rise above to a palisade of mycelium, the ends of which are enlarged, reaching a diameter of 15–17 μ, the palisade mycelium often appearing as rudimentary hairs; asci clavate, gradually tapering below into a stem-like base, with an abrupt swelling at the extreme base, reaching a length of 250–300 μ partly due to stretching at maturity and a diameter of 15–17 μ; spores at first 1-seriate, finally becoming partially to entirely 2-seriate, each spore surrounded by a granular protoplasm which sometimes collects about the outer surface, so as to give them a roughened appearance, ellipsoid, with the ends strongly narrowed, perfectly hyaline, 10–12 × 20–24 μ; paraphyses very slender, about 1.3 μ thick, rather strongly enlarged above, where they reach a diameter of 4 μ, smoky-brown above, hyaline below.

On excrement of horses.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to Colorado; also in Europe.

ILLUSTRATIONS: Ann. Sci. Nat. IV. **10**: *pl. 13, f. D;* Ann. Sci. Nat. V. **10**: *pl. 11, f. 37;* Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 28, f. 1;* Ann. Mag. Nat. Hist. III. **15**: *pl. 17, f. 30.*

15. **Ascophanus tetraonalis** Peck, Bull. N. Y. State Mus. **2**: 22. 1887.

Peziza tetraonalis Peck, Ann. Rep. N. Y. State Mus. **32**: 46. 1879.

Apothecia sessile, externally cinereous, the margin often wavy or flexuous, reaching a diameter of 1–3 mm.; hymenium blackish or blackish-brown; asci cylindric, truncate at the apex, 8-spored; spores 1-seriate, ellipsoid, smooth, colorless, reaching a length of 7–8 × 15–17 μ; paraphyses filiform, slightly enlarged above.

On the excrement of partridge or grouse.

TYPE LOCALITY: Catskill Mountains.

DISTRIBUTION: Known only from the type locality.

16. **Ascophanus lilacinus** Cooke, Grevillea **21**: 74. 1893.

Apothecia minute, gregarious, smooth, reaching a diameter of 0.3–0.8 mm.; hymenium slightly concave, pale-lilac, roughened by the protruding asci; asci rather fusoid-clavate, 8-spored; spores ellipsoid, hyaline, smooth, 10 × 20 μ; paraphyses filiform, scarcely thickened above, longer than the asci.

On horse dung.

TYPE LOCALITY: New York.

DISTRIBUTION: Known only from the type locality.

17. **Ascophanus cinerellus** (Karst.) Speg. Anal. Soc. Ci. Argent. **10**: 29. 1880.

Peziza cinerella Karst. Myc. Fenn. **1**: 51. 1871.

Apothecia scattered or subgregarious, sessile, cinereous-pallid, reaching a diameter of 0.3–0.4 mm.; hymenium convex, similar in color to the outside of the apothecium; asci cylindric-clavate, reaching a length of 38–42 μ and a diameter of 6–7 μ, 8-spored; spores 1-seriate or 2-seriate, ellipsoid or globose-ellipsoid, not containing oil-drops, 3–4 × 5–7 μ; paraphyses not visible.

On cow dung.

TYPE LOCALITY: Finland.

DISTRIBUTION: Greenland; also in Europe.

18. **Ascophanus microsporus** (Crouan) Phill. Brit. Discom. 307. 1887. (PLATE 11, FIG. 9.)

Ascobolus microsporus Berk. & Br. Ann. Mag. Nat. Hist. III. **15**: 449. 1865.
?Ascophanus minutissimus Boud. Ann. Sci. Nat. V. **10**: 243. 1869.
?Ascophanus Coemansii Ann. Sci. Nat. V. **10**: 244. 1869.

Apothecia gregarious but not usually crowded, at first globose, expanding and becoming discoid, with the hymenium plane or a little concave and the margin slightly elevated, externally dark brownish-black, when dry entirely black and appearing as minute black specks on the substratum, entire plant not exceeding 0.2–0.5 mm. in diameter; hymenium paler, reddish-brown; excipulum pseudoparenchymatous, the component cells 2–5 μ in diameter; asci clavate, 7–10 × 50–70 μ, tapering gradually below; spores minute, usually 2-seriate or partially 2-seriate, ellipsoid, smooth, perfectly hyaline, 4 × 6–7 μ; paraphyses very slender below, about 1 μ thick, abruptly expanding above into a subglobose head which is filled with golden-yellow coloring matter, reaching a diameter of 7–8 μ.



SPECIES OF ASCOPHANUS

On cow dung.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to Colorado; also in Europe.

ILLUSTRATIONS: Ann. Mag. Nat. Hist. III. **15**: *pl. 16, f. 28;* Ann. Sci. Nat. V. **10**: *pl. 27, f. 29–30;* Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 27, f. 2.*

19. **Ascophanus Cesatii** (Carest) Sacc. Syll. Fung. **8**: 533. 1889.

Ascobolus Cesatii Carest in Rab. Fungi Eu. *976.* 1866.
Pezizula Cesatii Karst. Myc. Fenn. **1**: 83. 1871.

Apothecia becoming discoid, externally brownish; hymenium pallid, nearly plane; asci clavate, 8-spored, reaching a length of 32 μ and a diameter of 8 μ; spores hyaline, ovoid, reaching a length of 8 μ; paraphyses simple, straight.

On bird dung.

TYPE LOCALITY: Europe.

DISTRIBUTION: Greenland; also in Europe.

20. **Ascophanus glaucellus** Rehm in Rab. Krypt.-Fl. 1^3: 1086. 1896.

Apothecia gregarious or scattered, sessile, becoming discoid, white or whitish, reaching a diameter of 0.2–0.5 mm.; asci clavate or subovoid, reaching a length of 40–50 μ and a diameter of 8–12 μ; spores 2-seriate, ellipsoid, smooth, hyaline, 3.5–4.5 $\times$ 6–9 μ; paraphyses slender, reaching a diameter of 1.5 μ below and 2.5 μ above, hyaline or yellowish, strongly curved at their apices.

On dung.

TYPE LOCALITY: Germany.

DISTRIBUTION: New York; also in Europe.

ILLUSTRATION: Rab. Krypt.-Fl. 1^3: 1080, *f. 5.*

17. **HUMARINA** Seaver, Mycologia **19**: 87. 1927.

Peziza § *Humaria* Fries, Syst. Myc. **2**: 42. 1822.
Humaria Sacc. Syll. Fung. **8**: 118. 1889. Not *Humaria* Fuckel. 1869.
Leucoloma Fuckel, Symb. Myc. 317. 1869. Not *Leucoloma* Brid. 1827.

Apothecia minute to medium-sized, usually less than 1 cm. in diameter and often less than 1 mm., usually light-colored, white, yellow, orange, or purple, more rarely dark-colored, brownish or blackish, usually discoid or more rarely cup-shaped,

occurring on humus often among mosses, or more rarely on the stems and leaves of the higher plants; asci clavate, 4–8-spored; spores 1–2-seriate, ellipsoid to fusoid or fusiform, smooth or becoming sculptured, hyaline; paraphyses slender or stout, usually containing a granular coloring matter.

Type species, *Octospora leucoloma* Hedw.

It is difficult to separate this from the preceding genus on morphological characters, while it is doubtful if they should be separated on any other. However, as the coprophilous forms have usually been segregated, in this case, we follow precedent and leave the reader to satisfy himself in the matter.

Apothecia light-colored, white, yellow, red, lilac, or violaceous.
Spores ellipsoid, usually not more than twice as long as broad.
Growing on the living stems of mosses and liverworts.
Occurring in the axils of the leaves of mosses. 1. *H. axillaris.*
Occurring on the thallus of *Marchantia.* 2. *H. ithacaensis.*
Growing on soil and on decaying plant remains.
Apothecia white or dirty-whitish.
Apothecia rounded or discoid. 3. *H. pallens.*
Apothecia hysteriform, partially closed. 4. *H. clausa.*
Apothecia orange, yellow, or lilac.
Occurring on old paper, cloth, twine, etc. 5. *H. testacea.*
Occurring on soil, usually among mosses.
Apothecia minute, rarely exceeding 1 mm. in diameter.
Spores becoming rough. 6. *H. orthotricha.*
Spores remaining permanently smooth.
Apothecia orange.
Spores 6–7 × 12 μ. 7. *H. phycophila.*
Spores 10–12 × 16–18 μ. 8. *H. rubens.*
Spores 12–14 × 20–25 μ. 9. *H. convexula.*
Apothecia with a lilac tinge. 10. *H. lilacina.*
Apothecia medium-sized, 2–10 mm. in diameter.
Spores broad-ellipsoid, usually containing one large oil-drop or none.
Apothecia bright-orange or red.
Apothecia entirely sessile. 11. *H. leucoloma.*

Apothecia short-stipitate. 12. *H. araneosa.*
Apothecia yellow to pale-orange.
Apothecia sessile, partially immersed. 13. *H. semiimmersa.*
Apothecia short-stipitate, superficial. 14. *H. ochroleuca.*
Spores narrow-ellipsoid, containing two oil-drops. 15. *H. rufa.*
Occurring on decaying plant remains.
Spores minute, 3 × 6–7 μ, on *Carex*. 16. *H. wisconsinensis.*
Spores medium-sized, 12–20 μ long.
Apothecia 1 mm. or less in diameter.
On brassicaceous plants. 17. *H. nectrioides.*
On unknown herbaceous plants. 18. *H. deligata.*
Apothecia 1 to several mm. in diameter.
Spores 10–11 × 20 μ; on chaff. 19. *H. glumarum.*
Spores 8 × 12 μ; on damp hay. 20. *H. Peckii.*
Spores 12 × 18–20 μ; on leaves. 21. *H. phyllogena.*
Spores fusoid, approximately three times as long as broad.
Asci 4-spored. 22. *H. tetraspora.*
Asci 8-spored.
Apothecia with a rose-colored hymenium. 23. *H. permuda.*
Apothecia with an orange hymenium.
Apothecia reaching a diameter of 5–8 mm. 24. *H. ollaris.*
Apothecia not exceeding 2–3 mm. in diameter.
Spores 22–25 μ long; on burned areas. 25. *H. aggregata.*
Spores 25–35 μ long.
Occurring on soil, but not on burned areas.
Spores almost fusiform. 26. *H. fusispora.*
Spores fusoid, but with ends blunt. 27. *H. coccinea.*
Occurring on dead sticks. 28. *H. Cookeina.*
Apothecia violaceous. 29. *H. Gerardi.*
Apothecia dark-colored, brown, brownish-black or more rarely purplish.
Spores becoming rough; apothecia purplish. 30. *H. purpurea.*
Spores remaining permanently smooth.
Apothecia liver-colored. 31. *H. hepatica.*
Apothecia dark but not liver-colored. 32. *H. trachyderma.*

1. **Humarina axillaris** (Nees) Seaver, comb. nov.

Peziza axillaris Nees, Syst. Pilze 258. 1817.
Leucoloma axillaris Fuckel, Symb. Myc. 318. 1869.
Humaria axillaris Sacc. Syll. Fung. **8**: 130. 1889.
Helotium axillaris Boud. Hist. Class. Discom. Eu. 114. 1907.

Apothecia sessile, subglobose, becoming slightly depressed, reaching a diameter of 1 mm., externally pale-orange, surrounded at the base by coarse mycelium which attaches the apothecia to the moss plants on which they grow; hymenium plane or nearly so, a little darker than the outside of the apothecium; asci cylindric above, narrowed below into a stem-like base, 8-spored; spores obliquely 1-seriate, hyaline, smooth, ellipsoid, containing one or two large oil-drops, 7–10 × 17–24 μ; paraphyses numerous, slender, slightly enlarged above.

Growing from the axils of leaves of mosses.

TYPE LOCALITY: Europe.

DISTRIBUTION: Colorado; also in Europe.

ILLUSTRATIONS: Nees, Syst. Pilze *pl. 37, f. 267;* Cooke, Mycographia *pl. 23, f. 91.*

EXSICCATI: Clements, Crypt. Form. Colo. *112.*

2. **Humarina ithacaensis** (Rehm) Seaver, comb. nov.

Humaria ithacaensis Rehm, Ann. Myc. **2**: 35. 1904.

Apothecia scattered, sessile, at first globose and closed, then cup-shaped, finally expanded, flesh-red, reaching a diameter of 0.5–1 mm., smooth, attached to the substratum by a coarse white mycelium; hymenium becoming plane or slightly convex; asci cylindric, reaching a length of 150–180 μ and a diameter of 15 μ, 8-spored; spores 1-seriate, ellipsoid, containing two large oil-drops, hyaline, 8–9 × 15–17 μ, becoming minutely sculptured; spore-sculpturing consisting of minute warts; paraphyses filiform, septate, hyaline, 3 μ in diameter below, enlarged above, reaching a diameter of 4 μ at their apices.

On the living thallus of *Marchantia polymorpha.*

TYPE LOCALITY: Ithaca, New York.

DISTRIBUTION: Known only from the type locality.

3. **Humarina pallens** (Boud.) Seaver, comb. nov.

Ascophanus pallens Boud. Bull. Soc. Myc. Fr. **4**: XLVIII. 1888.

Apothecia gregarious but not crowded, at first subglobose, expanding and becoming discoid or subdiscoid, entirely white or

dirty-white, becoming darker when dry, reaching a diameter of 2–3 mm.; hymenium plane or strongly convex, roughened by the large protruding asci; asci clavate, reaching a length of 350–400 μ and a diameter of 35–50 μ at their apices, tapering below into a slender stem-like base, 8-spored; spores 1–3-seriate or more often irregularly crowded together, granular within and usually surrounded by a hyaline membrane, large, narrow-ellipsoid or with the ends rather abruptly narrowed, about 15–20 × 35–45 μ, hyaline; paraphyses slender, strongly enlarged above.

On sandy soil which is sparingly overrun with algae.

TYPE LOCALITY: France.

DISTRIBUTION: New York; also in Europe.

ILLUSTRATIONS: Bull. Soc. Myc. Fr. **4**: *pl. 2, f. 2;* Boud. Ic. Myc. *pl. 414.*

4. **Humarina clausa** (Clem.) Seaver, comb. nov.

Humaria clausa Clements, Bot. Surv. Nebr. **4**: 10. 1896.

Apothecia cespitose, fleshy, subglobose or nearly hemispheric, pruinose, reaching a diameter of 2 mm.; hymenium somewhat hysteriform, gaping slightly, drab; asci clavate, reaching a length of 250 μ and a diameter of 35 μ, 8-spored; spores large, ellipsoid, smooth, devoid of oil-drops, 12.5–16 × 32–37 μ; paraphyses abundant, filiform, scarcely enlarged at their apices.

On moist ground.

TYPE LOCALITY: Otawaine woods, Lancaster County, Nebraska.

DISTRIBUTION: Known only from the type locality.

5. **Humarina testacea** (Moug.) Seaver, comb. nov.

Peziza testacea Moug.; Fries, Elench. Fung. **2**: 11. 1828.
Helotium testaceum Berk. Outl. Brit. Fungol. 372. 1860.
Ascobolus testaceus Berk. Ann. Mag. Nat. Hist. III. **15**: 447. 1865.
Ascophanus testaceus Phill. Brit. Discom. 310. 1887.
Humaria testacea Schröt. Krypt.-Fl. Schles. **3**²: 36. 1893.
Ascophanus carneus testaceus Massee, Brit. Fungus-Fl. **4**: 178. 1895.

Apothecia gregarious or densely crowded, at first subglobose, becoming discoid or subdiscoid, flesh-red, not usually exceeding 1–2 mm. in diameter; hymenium at first slightly concave, becoming plane or more often convex, roughened by the protruding asci, similar in color to the outside of the apothecium or a little brighter; the excipular cells irregular and reaching a

diameter of 12–15 μ, giving rise above to palisade-like fascicles of club-shaped mycelial threads; asci rather broad-clavate, reaching a length of 200 μ and a diameter of 20–25 μ, 8-spored; spores at first quite regularly 2-seriate, becoming irregularly disposed, ellipsoid, hyaline, smooth or very minutely sculptured, 10–12 × 17–20 μ; spore-sculpturing consisting of minute granules; paraphyses slightly enlarged above, reaching a diameter of 7–8 μ at their apices, filled with orange granules.

On old building paper, sacking, cloth, twine, leather, and on weathered fibers of *Yucca* leaves.

TYPE LOCALITY: Europe.

DISTRIBUTION: Massachusetts to New Jersey, Louisiana and North Dakota; also in Europe.

ILLUSTRATIONS: Phill. Brit. Discom. *pl. 9, f. 58;* Ann. Mag. Nat. Hist. III. **15**: *pl. 14, f. 5;* Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 27, f. 1.*

On morphological characters it is difficult, if not impossible, to separate this species from *Ascophanus carneus*, its nearest coprophilous relative. In fact, Massee treats this species as a variety of the latter. While some workers seem to see a difference in spore measurements, such a difference is scarcely apparent so far as our observations go.

The habitat of this species is unique. The persistence with which it occurs on cloth, string, and building paper is interesting. It was first encountered by the writer early in his studies on the Discomycetes at Lafayette, Indiana, where it was found in abundance on old burlap sacking, the search for it having been prompted by Phillips' illustration of the species in his British Discomycetes. About the same time it was found on old rag carpet at Pocahontas, Iowa. In 1905 it was found in abundance around Mt. Pleasant, Iowa, on old paper and sacking, and in 1907 on a similar habitat in Fargo, North Dakota. In 1913 it was collected at Portland, Connecticut, on old paper, twine and sacking.

In 1916, a fine collection was made at Yonkers, New York, on decaying paper and the surrounding soil in a dumping ground. This collection showed unusually large spores, measuring 16–18 × 28–30 μ, and was listed as a new variety under the name *Humarina testacea macrospora*. The spores in this form were decidedly rough.

Further study is necessary in order to determine whether

this species and *Ascophanus carneus* are really distinct but for the present they are allowed to stand.

6. **Humarina orthotricha** (Cooke & Ellis) Seaver, comb. nov.

Peziza orthotricha Cooke & Ellis; Cooke, Grevillea **6**: 7. 1877.
Humaria orthotricha Sacc. Syll. Fung. **8**: 119. 1889.

Apothecia widely scattered, at first globose and closed, opening and at maturity becoming scutellate or subdiscoid, reaching a diameter of 1 mm., externally pale-orange; hymenium plane or a little concave, orange-vermillion; asci cylindric or subcylindric to clavate, 8-spored, reaching a length of 200–250 μ and a diameter of 18–20 μ; spores 1-seriate or becoming irregularly crowded, ellipsoid, becoming sculptured, 10–12 $\times$ 18–22 μ; spore-sculpturing taking the form of minute warts not exceeding 1 μ in diameter; paraphyses gradually thickened above, reaching a diameter of 5 μ at their apices, often strongly curved.

On soil and bark among mosses, especially *Orthotrichum*.

TYPE LOCALITY: New Jersey.

DISTRIBUTION: New York to New Jersey and Colorado.

ILLUSTRATIONS: Grevillea **6**: *pl. 96, f. 33;* Cooke, Mycographia *pl. 98, f. 355.*

7. **Humarina Clementsii** Seaver, nom. nov.

Humaria phycophila Clements, Bot. Surv. Nebr. **4**: 10. 1896. Not *Humaria phycophila* Oudem. Beih. Bot. Centr. **11**: 526. 1902.

Apothecia very minute, gregarious, sessile, scutellate, finally becoming subdiscoid, reaching a diameter of 0.3–0.5 mm.; hymenium at first concave, finally becoming convex, orange to reddish, darker than the outside of the apothecium; asci clavate-cylindric, obliquely operculate, 8-spored, reaching a length of 60–67 μ and a diameter of 9–15 μ; spores ellipsoid, obliquely 1-seriate or often becoming 2-seriate, usually containing one oil-drop, smooth, hyaline, 6–7.5 $\times$ 10–12.5 μ; paraphyses filiform, filled with orange granules, strongly curved at their apices, reaching a diameter of 1–2 μ.

Among filaments of *Lyngbya* on moist ground.

TYPE LOCALITY: Beatrice, Nebraska.

DISTRIBUTION: Known only from the type locality.

8. **Humarina rubens** (Boud.) Seaver, comb. nov.

Humaria rubens Boud. Bull. Soc. Myc. Fr. **12**: 13. 1896.

Apothecia gregarious but not usually crowded, at first subglobose, expanding and becoming discoid or subdiscoid, reaching

a diameter of 1–3 mm., externally pale-orange; hymenium becoming plane or occasionally a little convex, orange to bright-red; asci cylindric above, gradually tapering below into a rather short stem-like base, reaching a length of 200 μ and a diameter of 16 μ, 8-spored; spores 1-seriate, ellipsoid or ovoid, each containing one large oil-drop, hyaline, smooth, 10–12 × 16–18 μ; paraphyses rather stout, simple or branched, often swollen near the joints, reaching a diameter of 8–9 μ at their apices.

On damp soil among mosses.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York and Colorado; also in Europe.

ILLUSTRATION: Bull. Soc. Myc. Fr. **12**: *pl. 3, f. 3.*

9. **Humarina convexula** (Pers.) Seaver, comb. nov.

Peziza convexula Pers. Obs. Myc. **2**: 85. 1799.
Leucoloma convexula Fuckel, Symb. Myc. 318. 1869.
Peziza chrisophthalma Ger. Bull. Torrey Club **4**: 48. 1873. Not *Peziza chrysophthalma* Pers. 1822.
Aleuria convexula Gill. Champ. Fr. Discom. 55. 1879.
Humaria convexula Sacc. Syll. Fung. **8**: 139. 1889.

Apothecia gregarious but not usually crowded, at first globose or short-cylindric, soon opening, pale- to bright-orange, rarely exceeding 1–2 mm. in diameter and often less; hymenium becoming convex, similar in color to the outside of the apothecium; asci cylindric above, tapering rather abruptly below, reaching a length of 200 μ and a diameter of 18–20 μ, 8-spored; spores obliquely or irregularly disposed, broad-ellipsoid, hyaline, smooth, containing either one or two large oil-drops which reach a diameter of about 10 μ, entire spore 12–14 × 22–25 μ; paraphyses straight or very slightly curved, rather stout, about 2 or 3 μ thick below, strongly enlarged above, where they reach a diameter of 7–8 μ, filled with rather coarse orange granules.

On bare soil or among mosses.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York; also in Europe.

ILLUSTRATIONS: Cooke, Mycographia *pl. 9, f. 36;* Grevillea **3**: *pl. 31, f. 81;* Gill. Champ. Fr. Discom. *pl. 56, f. 1.*

10. **Humarina lilacina** Seaver, sp. nov.

Apothecia rather thickly gregarious or rarely with several in close contact but not usually congested, at first globose or subglobose, expanding and becoming subdiscoid, reaching a diameter

of 0.5 mm. or rarely as large as 1 mm., reddish, with a decided lilac tinge; hymenium at first concave, finally plane or convex, the margin becoming indistinct, roughened by the protruding asci, the color similar to that of the outside of the apothecium; asci clavate or subclavate, reaching a length of 200 μ and a diameter of 15–17 μ, 8-spored; spores 1-seriate or becoming irregularly crowded, ellipsoid, usually containing one large often eccentric oil-drop half the diameter of the spore, 10 $\times$ 15–17 μ; paraphyses strongly enlarged above, septate, filled with coarse granules which are subhyaline or with a pale-lilac tinge.

On the ground which has been overrun with mosses and algae.

Type collected in the New York Botanical Garden, September 5, 1912 (Herb. The New York Botanical Garden).

DISTRIBUTION: New York and New Jersey.

11. **Humarina leucoloma** (Hedw.) Seaver, comb. nov.

Octospora leucoloma Hedw. Descr. **2**: 13. 1788.
Peziza leucoloma Rebent. Fl. Neom. 386. 1804.
Peziza humosa Fries, Obs. Myc. **2**: 308. 1818.
Leucoloma Hedwigii Fuckel, Symb. Myc. 317. 1869.
Aleuria humosa Gill. Champ. Fr. Discom. 56. 1879.
Aleuria leucoloma Gill. Champ. Fr. Discom. 56. 1879.
Humaria humosa Quél. Ench. Fung. 289. 1886.
Humaria leucoloma Quél. Ench. Fung. 289. 1886.
Neottiella leucoloma Massee, Brit. Fungus-Fl. **4**: 374. 1895.
Leucoscypha excipulata Clements, Crypt. Form. Colo. *117*. 1906.

Apothecia gregarious or scattered, sessile and often with the base nestling in the substratum, at first globose and closed, expanding and becoming scutellate to subdiscoid, reaching a diameter of 2–5 mm., externally pale-orange; hymenium concave or often becoming plane or a little convex, bright-orange; asci cylindric or subclavate, gradually tapering below into a long stem-like base, reaching a length of 200–250 μ and a diameter of 20 μ, 8-spored; spores 1-seriate or occasionally crowded, usually diagonally disposed, ellipsoid or with one or both ends slightly narrowed, 12–15 $\times$ 18–24 μ, a few often larger, smooth, usually containing one large oil-drop surrounded by several smaller ones; paraphyses 3–4 μ in diameter below, enlarged above, reaching a diameter of 8–9 μ at their apices, branched, often curved and filled with numerous granules.

On the ground among mosses.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to Delaware and Colorado; also in Europe.

ILLUSTRATIONS: Hedw. Descr. **2**: *pl. 4, f. 1–7;* Cooke, Mycographia *pl. 7, f. 28;* Boud. Ic. Myc. *pl. 391* and *395;* Gill. Champ. Fr. Discom. *pl. 56, f. 2;* Nees, Syst. Pilze *pl. 37, f. 268;* Ann. Mag. Nat. Hist. III. **18**: *pl. 3, f. 16.*

EXSICCATI: Fungi Columb. *3815;* Sydow, Fungi Exotici Exsicc. *139;* Clements, Crypt. Form. Colo. *117* (as *Leucoscypha excipulata*).

12. **Humarina araneosa** (Peck) Seaver, comb. nov.

Peziza araneosa Bull. Herb. Fr. *pl. 280.* 1785.
Leucoloma araneosa Fuckel, Symb. Myc. Nacht. **1**: 50. 1871.
Peziza rubra Peck, Ann. Rep. N. Y. State Mus. **24**: 95. 1872.
Aleuria araneosa Gill. Champ. Fr. Discom. 55. 1872.
Humaria araneosa Sacc. Syll. Fung. **8**: 122. 1889.

Apothecia subglobose, finally becoming hemispheric, the mouth constricted and often slightly irregular, short-stipitate, externally pale-whitish or reddish, reaching a diameter of 4–5 mm.; stem short, stout; hymenium bright-red, concave; asci cylindric or subcylindric, 8-spored; spores 1-seriate, ellipsoid and containing no oil-drops, hyaline, $7 \times 14\ \mu$; paraphyses filiform, clavate.

On damp soil and on burnt ground.

TYPE LOCALITY: Highlands, New York.

DISTRIBUTION: New York; also in Europe.

ILLUSTRATIONS: Ann. Rep. N. Y. State Mus. **24**: *pl. 2, f. 20, 21;* Sow. Engl. Fungi *pl. 369, f. 5;* Bull. Herb. Fr. *pl. 280.*

13. **Humarina semiimmersa** (Karst.) Seaver, comb. nov.

Peziza semiimmersa Karst. Not. Fauna Fl. Fenn. **10**: 117. 1869.
Humaria semiimmersa Sacc. Syll. Fung. **8**: 143. 1889.
Sepultaria semiimmersa Massee, Brit. Fungus-Fl. **4**: 391. 1895.

Apothecia gregarious, sessile and partially immersed in the ground, at first subglobose and closed, expanding and becoming shallow cup-shaped to scutellate, reaching a diameter of 1 cm., externally pale-yellow, more or less floccose or tomentose, the margin irregular and often torn; hymenium concave, bright-yellow to pale-ochraceous; asci cylindric or subcylindric above, reaching a diameter of 17–20 μ, 8-spored; spores 1-seriate, obliquely arranged in the ascus or occasionally parallel with

the ascus, ellipsoid, hyaline or very faintly yellowish, smooth, 10–11 × 18–20 μ; paraphyses slender, slightly enlarged above, where they reach a diameter of 2 μ, yellowish.

On bare ground.

TYPE LOCALITY: Europe.

DISTRIBUTION: Delaware and Kentucky; also in Europe.

ILLUSTRATIONS: Cooke, Mycographia *pl. 12, f. 46;* Grevillea **3**: *pl. 31, f. 75.*

14. **Humarina ochroleuca** (Clements) Seaver, comb. nov.

Humaria ochroleuca Clements, Bull. Torrey Club **30**: 91. 1903.

Apothecia solitary, sessile or contracted below into a stem-like base, cupulate, finally expanded, reaching a height of 2–6 mm. and a diameter of 3–6 mm., yellowish-white, margin crenulate; hymenium at first concave, becoming plane or occasionally convex, the color similar to that of the outside of the apothecium; asci cylindric above, long-stipitate below, reaching a length of 275 μ and a diameter of 11–12 μ, 8-spored; spores irregularly 1-seriate, ellipsoid, smooth, hyaline, 11 × 19 μ; paraphyses slender, about 3 μ in diameter, slightly enlarged above.

Among mosses.

TYPE LOCALITY: Engelmann Canyon, Colorado.

DISTRIBUTION: Colorado.

EXSICCATI: Clements, Crypt. Form. Colo. *113.*

15. **Humarina rufa** (Pers.) Seaver, comb. nov.

Peziza rufa Pers. Obs. Myc. **2**: 79. 1799.
Peziza rubricosa Fries, Syst. Myc. **2**: 72. 1822.
Leucoloma rubricosa Fuckel, Symb. Myc. 318. 1869.
Humaria rubricosa Sacc. Syll. Fung. **8**: 138. 1889.

Apothecia scattered, sessile or tapering below into a stem-like base, externally rough, orange, reaching a diameter of 1–4 mm.; hymenium a little concave to nearly plane, orange, a little darker than the exterior of the apothecium, excipulum giving rise to a palisade of mycelium above, the loose ends of which form a fringe-like border, are clavate and reach a diameter of 10 μ, hyaline; asci rather broad-cylindric or clavate, reaching a length of 75 μ and a diameter of 20 μ at their apices, 8-spored; spores 2-seriate or irregularly crowded above, partially 1-seriate below, narrow-ellipsoid, often curved or unequal-sided, containing two oil-drops, hyaline, 9–10 × 20–23 μ; paraphyses enlarged above, reaching a diameter of 7–8 μ.

Among mosses.

TYPE LOCALITY: Europe.

DISTRIBUTION: Delaware; also in Europe.

ILLUSTRATIONS: Cooke, Mycographia *pl. 18, f. 71;* Boud. Ic. Myc. *pl. 397;* Rab. Krypt.-Fl. 1^3: 916, *f. 5.*

16. **Humarina wisconsinensis** (Rehm) Seaver, comb. nov.

Humaria wisconsinensis Rehm, Ann. Myc. **3**: 517. 1905.

Apothecia gregarious, sessile, at first globose and closed, expanding and becoming discoid, externally pale-yellow, smooth, attached to the substratum by conspicuous white mycelium, reaching a diameter of 5 mm.; hymenium plane or nearly plane, orange-yellow; asci clavate, the apex rounded, reaching a length of 40–45 μ and a diameter of 5–6 μ, 8-spored; spores 2-seriate, ellipsoid, hyaline, 3 $\times$ 6–7 μ; paraphyses filiform, septate, reaching a diameter of 2–2.5 μ, scarcely enlarged above, yellow.

On decaying *Carex.*

TYPE LOCALITY: Blue Mountains, Wisconsin.

DISTRIBUTION: Known only from the type locality.

17. **Humarina nectrioides** Seaver, sp. nov.

Apothecia scattered or gregarious, sessile, almost globose, becoming somewhat discoid, bright-orange, not exceeding 1 mm. in diameter and closely resembling a *Nectria;* hymenium plane or slightly concave; asci clavate, 8-spored, reaching a length of 150 μ and a diameter of 20 μ; spores 2-seriate or irregularly disposed, ellipsoid, hyaline, smooth, about 12 $\times$ 20 μ; paraphyses stout, septate, thickened above, reaching a diameter of 10 μ at their apices, filled with minute orange granules.

Type collected on old cabbage stems associated with *Pseudombrophila deerata* (Karst.) Seaver, Kulm, North Dakota, July 4, 1911 (Herb. The New York Botanical Garden).

DISTRIBUTION: Known only from the type locality.

EXSICCATI: Brenckle, Fungi Dakotenses *140* (in part).

18. **Humarina deligata** (Peck) Seaver, comb. nov.

Peziza deligata Peck, Ann. Rep. N. Y. State Mus. **30**: 61. 1879.
Humaria deligata Sacc. Syll. Fung. **8**: 123. 1889.

Apothecia gregarious, minute, sessile, with radiating, hyaline mycelial threads at the base, at first subglobose, then opening and becoming scutellate to subdiscoid, externally pale-reddish;

hymenium slightly concave or plane, reddish, a little brighter than the outside of the apothecium; asci subcylindric to clavate, reaching a length of 150 μ and a diameter of 20 μ, 8-spored; spores 2-seriate above or irregularly disposed, smooth, hyaline, 10–12 $\times$ 18–20 μ; paraphyses very slender, reaching a diameter of about 3 μ below, slightly enlarged at their apices.

On dead herbs lying on the ground.

TYPE LOCALITY: Bethlehem, New York.

DISTRIBUTION: Known only from the type locality.

19. **Humarina glumarum** (Desm.) Seaver, comb. nov.

Peziza glumarum Desm. Ann. Sci. Nat. II. **15**: 129. 1841.
Aleuria glumarum Gill. Champ. Fr. Discom. 55. 1879.
Humaria glumarum Sacc. Syll. Fung. **8**: 130. 1889.

Apothecia gregarious, at first closed, gradually expanding and becoming scutellate, reaching a diameter of 4–8 mm., orange; hymenium concave or nearly plane, a little darker than the outside of the apothecium; asci cylindric above, tapering below, reaching a length of 235 μ and a diameter of 15–16 μ, 8-spored; spores ellipsoid, 1-seriate or partially 2-seriate, smooth, hyaline or very faintly yellowish, 10–11 $\times$ 20 μ; paraphyses slender, enlarged above, reaching a diameter of 7–8 μ.

On chaff and straw.

TYPE LOCALITY: Europe.

DISTRIBUTION: Iowa; also in Europe.

ILLUSTRATION: Cooke, Mycographia *pl. 10, f. 39*.

20. **Humarina Peckii** (House) Seaver, comb. nov.

Humaria Peckii House, Bull. N. Y. State Mus. **205–206**: 38. 1919.

Apothecia at first closed, finally opening and becoming shallow cup-shaped to scutellate, externally pallid, subpruinose, surrounded at the base with a mass of white mycelium, reaching a diameter of 4 mm.; hymenium slightly concave, pale-yellow or subochraceous; asci cylindric or subcylindric, reaching a length of 160 μ and a diameter of 12 μ, attenuated below into a rather long stem-like base; spores ellipsoid or ovoid, 1-seriate, each containing one oil-drop, about 8 $\times$ 12 μ; paraphyses not exceeding 2 μ in diameter, scarcely thickened above.

On damp decaying hay accompanied by a *Sclerotium*.

TYPE LOCALITY: Menandos, Albany County, New York, April, 1890 (C. H. Peck).

DISTRIBUTION: Known only from the type locality.

21. **Humarina phyllogena** Seaver, comb. nov.

Humaria phyllogena Seaver, Mycologia **17**: 46. 1925.

Apothecia sessile or subsessile, discoid or with the margin slightly elevated, reaching a diameter of 3–5 mm., flesh-red, brighter when dry; asci clavate, operculate, reaching a length of 200 μ and a diameter of 12 μ; spores broad-ellipsoid, 12 $\times$ 18–20 μ, usually containing two oil-drops, smooth.

On dead leaves.

TYPE LOCALITY: El Yunque, Porto Rico.

DISTRIBUTION: Known only from the type locality.

22. **Humarina tetraspora** (Fuckel) Seaver, comb. nov.

Ascobolus tetrasporus Fuckel, Hedwigia **5**: 4. 1866.
Leucoloma tetraspora Fuckel, Symb. Myc. 317. 1869.
Peziza tetraspora Cooke, Grevillea **3**: 73. 1874.
Aleuria tetraspora Gill. Champ. Fr. Discom. 207. 1886.
Humaria tetraspora Sacc. Syll. Fung. **8**: 121. 1889.
Humaria silvosa Clements, Bot. Surv. Nebr. **5**: 7. 1901.

Apothecia gregarious or scattered, sessile, at first closed and subglobose, expanding and becoming subdiscoid, reaching a diameter of 2–3 mm.; hymenium at first concave, becoming nearly plane or slightly convex, surrounded by a delicate fringe-like border, bright-orange; asci clavate, reaching a length of 150–200 μ and a diameter of 15–22 μ, 4-spored; spores 1-seriate, with the ends strongly overlapping, narrow, ellipsoid to subfusoid, containing two or three oil-drops, 10–12 $\times$ 22–26 μ, or rarely reaching a length of 30 μ, hyaline; paraphyses strongly enlarged at their apices, where they reach a diameter of 7–8 μ, filled with orange granules.

On the ground usually among mosses.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to Iowa and Minnesota; also in Europe.

ILLUSTRATIONS: Hedwigia **5**: *pl. 1, f. 5;* Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 13, f. 1;* Boud. Ic. Myc. *pl. 393;* Cooke, Mycographia *pl. 12, f. 45;* Gill. Champ. Fr. Discom. *pl. 52, f. 1;* Pat. Tab. Fung. *f. 482.*

23. **Humarina permuda** (Cooke) Seaver, comb. nov.

Peziza fusispora var. *permuda* Cooke, Mycographia 209. 1878.
Peziza permuda Cooke, Mycographia 209. 1878 (as synonym).
Humaria permuda Sacc. Syll. Fung. **8**: 134. 1889.

Apothecia gregarious or scattered, at first cup-shaped, expanding and becoming discoid, whitish with a tinge of pink, reaching a diameter of 1–2 mm.; hymenium slightly concave or nearly plane; asci cylindric or subcylindric, 8-spored; spores obliquely 1-seriate, fusoid, each containing one or two oil-drops, reaching a size of 10 × 30 μ; paraphyses slender, only slightly enlarged above.

On the ground.

TYPE LOCALITY: South Carolina.

DISTRIBUTION: Known only from the type locality.

ILLUSTRATION: Cooke, Mycographia *pl. 98, f. 353.*

24. **Humarina ollaris** (Fries) Seaver, comb. nov.

Peziza ollaris Fries, Syst. Myc. **2**: 68. 1822.
Aleuria ollaris Gill. Champ. Fr. Discom. 51. 1879.
Humaria ollaris Sacc. Syll. Fung. **8**: 131. 1889.
Neottiella ollaris Clements, Bot. Surv. Nebr. **5**: 22. 1901.

Apothecia gregarious or more commonly crowded, sessile, at first closed and hemispherical, finally expanding and becoming scutellate, reaching a diameter of 5–8 mm.; excipulum giving rise above to a palisade of parallel mycelial threads, the ends of which give to the border of the apothecium a minutely downy appearance, the ends of the mycelium; hymenium orange, a little darker than the exterior; asci cylindric above, tapering below into a stem-like base, 8-spored; spores obliquely 1-seriate, hyaline, smooth, containing one or two oil-drops, fusoid, 8–11 × 26–30 μ; paraphyses septate, strongly enlarged above, reaching a diameter of 8–10 μ at their apices.

On the ground among mosses and in plant-pots.

TYPE LOCALITY: Europe.

DISTRIBUTION: Nebraska and New York; also in Europe.

ILLUSTRATIONS: Cooke, Mycographia *pl. 14, f. 56;* Gonnerm. & Rab. Myc. Eu. **2**: *pl. 2, f. 2.*

The species described under this name has been reported by Clements from Nebraska. A fine collection obtained March 19, 1928, on well-manured soil in pots in greenhouse of The New York Botanical Garden has been referred here. Our plants reach a diameter of 5 mm. and as they expand they split so that the margin appears crenulate with a whitish border but scarcely pubescent. The spores are a little broader than usually indicated for this species. The color of the hymenium is a dirty-

orange. The paraphyses are strongly enlarged and several times branched.

There seems to be some confusion regarding this species in Europe and of course the question arises as to whether we have the same thing here. Perhaps future study will add to our knowledge of this form. Our specimens are preserved in the herbarium of The New York Botanical Garden.

25. **Humarina aggregata** (Berk. & Br.) Seaver, comb. nov.

Peziza aggregata Berk. & Br. Ann. Mag. Nat. Hist. III. **18**: 123. 1866.
Humaria aggregata Sacc. Syll. Fung. **8**: 134. 1889.

Apothecia gregarious or more often crowded together into confluent masses, subcylindric or turbinate, externally whitish or pale-orange, reaching a diameter of 2–3 mm., seated on a conspicuous mycelial growth; hymenium slightly concave or nearly plane, surrounded by a delicate fringe-like border, bright-orange; asci cylindric to clavate, tapering rather abruptly below, reaching a length of 200–225 μ and a diameter of 10–12 μ, 8-spored; spores 1-seriate, often with the ends strongly overlapping or irregularly crowded and partially 2-seriate, fusoid, containing two oil-drops, 8–9 $\times$ 22–25 μ, hyaline; paraphyses rather strongly enlarged above.

On the ground, on burned places and on humus in coniferous woods.

TYPE LOCALITY: Great Britain.

DISTRIBUTION: New Hampshire and Indiana; also in Europe.

ILLUSTRATIONS: Grevillea **3**: *pl. 50, f. 236;* Cooke, Mycographia *pl. 8, f. 31;* E. & P. Nat. Pfl. 1^1: 184, *f. 149, F–H.*

26. **Humarina fusispora** (Berk.) Seaver, comb. nov.

Peziza fusispora Berk. London Jour. Bot. **5**: 5. 1846.
Humaria fusispora Sacc. Syll. Fung. **8**: 133. 1889.
Leucoloma fusispora Rehm, Ascom. *1053;* Hedwigia **31**: 301. 1892.

Apothecia gregarious or crowded, at first subglobose, becoming scutellate to discoid, reaching a diameter of 2–3 mm., externally pale-orange, surrounded at the base by a rather conspicuous mycelial growth; hymenium concave or nearly plane, bright-orange; asci subcylindric to clavate, 8-spored, reaching a length of about 250 μ and a diameter of 12 μ; spores 1-seriate, with the ends overlapping or becoming partially 2-seriate, fusoid or almost fusiform, containing one or more commonly two oil-drops, 8–10 $\times$ 30–33 μ, hyaline; paraphyses filiform, slightly enlarged above.

On the ground.

TYPE LOCALITY: Tasmania.

DISTRIBUTION: New Jersey to South Carolina; also in Europe and Tasmania.

ILLUSTRATION: Cooke, Mycographia *pl. 8, f. 32.*

EXSICCATI: Ravenel, Fungi Am. Exsicc. *174.*

27. **Humarina coccinea** (Crouan) Seaver, comb. nov.

Ascobolus coccineus Crouan, Ann. Sci. Nat. IV. **7**: 174. 1857.
Leucoloma coccinea Fuckel, Symb. Myc. 318. 1869.
Peziza corallina Cooke, Mycographia 19. 1875.
Aleuria corallina Gill. Champ. Fr. Discom 51. 1879.
Peziza muralis Quél. Grevillea **8**: 116. 1880.
Humaria coccinea Sacc. Syll. Fung. **8**: 119. 1889.
Humaria muralis Sacc. Syll. Fung. **8**: 127. 1889.
Neottiella corallina Massee, Brit. Fungus-Fl. **4**: 372. 1895.
Plicaria muralis Rehm in Rab. Krypt.-Fl. 1^3: 1006. 1896.

Apothecia gregarious or scattered, at first closed and subglobose, later expanding and becoming nearly discoid, reaching a diameter of 2–3 mm.; hymenium slightly concave or nearly plane, surrounded by a delicate ragged fringe-like border, bright-orange, brighter than the outside of the apothecium; asci clavate, 8-spored; spores 2-seriate above or irregularly crowded, fusoid, with the ends strongly narrowed but not acute, containing two or three oil-drops, rarely one, 7–10 × 25–35 μ; paraphyses strongly enlarged above, reaching a diameter of 6–10 μ at their apices.

On sandy soil among mosses.

TYPE LOCALITY: France.

DISTRIBUTION: Iowa; also in Europe.

ILLUSTRATIONS: Ann. Sci. Nat. **7**: *pl. 4, f. 15–19;* Boud. Ic. Myc. *pl. 392;* Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 13, f. 2;* Cooke, Mycographia *pl. 8, f. 30;* Proc. Assoc. Fr. Av. Sci. **12**: *pl. 7, f. 7;* Gill. Champ. Fr. Discom. (suites); Massee, Brit. Fungus-Fl. **4**: 290, *f. 33–35;* E. & P. Nat. Pfl. 1^1: 184, *f. 149, E;* Rab. Krypt.-Fl. 1^3: 917, *f. 1–4;* Phill. Brit. Discom. *pl. 4, f. 21;* Grevillea **3**: *pl. 31, f. 83.*

28. **Humarina Cookeina** Seaver, comb. nov.

Humaria Cookeina Seaver, Mycologia **17**: 46. 1925.

Apothecia short-stipitate or subsessile, with the hymenium slightly concave, reaching a diameter of 2–3 mm., bright-orange, a little paler outside, the stem slightly grooved or furrowed;

asci clavate, 8-spored, 14–15 × 160 mm.; spores partially 2-seriate, fusoid, 6 × 30 μ.

On decaying sticks.

TYPE LOCALITY: El Yunque, Porto Rico.

DISTRIBUTION: Known only from the type locality.

29. **Humarina Gerardi** (Cooke) Seaver, comb. nov.

Peziza Gerardi Cooke, Hedwigia **14**: 81. 1875.
Peziza violacea Ger.; Cooke, Hedwigia **14**: 81 (as synonym). 1875. Not *Peziza violacea* Pers. 1797.
Peziza ionella Quél. Bull. Soc. Bot. Fr. **24**: 328. 1877.
Aleuria ionella Gill. Champ. Fr. Discom. 51. 1879.
Humaria ionella Sacc. Syll. Fung. **8**: 149. 1889.
Humaria Gerardi Sacc. Syll. Fung. **8**: 150. 1889.
Galactinia ionella Boud. Hist. Class. Discom. Eu. 47. 1907.

Apothecia gregarious, at first closed and subglobose, expanding and becoming scutellate or nearly discoid, sessile or substipitate, reaching a diameter of 4–5 mm., externally pale-violaceous and slightly roughened; hymenium slightly concave or nearly plane, violaceous; asci clavate, 8-spored; spores 1-seriate or partially 2-seriate above, occasionally becoming irregularly crowded, fusoid, smooth, containing one to several oil-drops, 9–10 × 30–35 μ, hyaline; paraphyses rather stout, enlarged above.

On the ground.

TYPE LOCALITY: New York.

DISTRIBUTION: New York and Minnesota; also in Europe.

ILLUSTRATIONS: Cooke, Mycographia *pl. 11, f. 44;* (?) Bull. Soc. Bot. Fr. **24**: *pl. 6, f. 4;* Grevillea **3**: *pl. 43, f. 206;* Boud. Ic. Myc. *pl. 301.*

30. **Humarina purpurea** Seaver, sp. nov.

Apothecia gregarious or closely crowded and often irregular from mutual pressure, sessile, becoming scutellate, externally rough and dark reddish-purple to blackish-brown, the margin crenate, reaching a diameter of 2–3 mm.; hymenium concave, reddish-purple, lighter than the outside of the apothecium; excipular cells elongated, rectangular or polyhedral, reaching a diameter of 25–30 μ, with brown walls, giving rise to fascicles of dark-brown hair-like structures which project and give rise to the crenate margin; asci cylindric or subcylindric, reaching a length of 150–175 μ and a diameter of 11–12 μ, 8-spored; spores 1-seriate, parallel with the ascus or slightly oblique, fusoid, at

first smooth and containing one oil-drop about half the diameter of the spore or rarely with two, 8–10 × 13–20 μ, becoming delicately sculptured at maturity; spore-sculpturing consisting of minute warts or very short irregular ridges, often with a minute apiculus at each end; paraphyses slender and often adhering together at their apices, purplish.

On damp soil.

Type collected in Colorado by E. Bethel (1914?) (Herb. The New York Botanical Garden).

DISTRIBUTION: Known only from the type locality.

31. **Humarina hepatica** (Batsch) Seaver, comb. nov.

Peziza hepatica Batsch, Elench. Fung. Contin. **1**: 199. 1786.
Aleuria hepatica Gill. Champ. Fr. Discom. 208. 1886.
Humaria hepatica Sacc. Syll. Fung. **8**: 140. 1889.
Ascophanus hepaticus Boud. Hist. Class. Discom. Eu. 76. 1907.

Apothecia gregarious or scattered, at first closed and subglobose, opening and becoming scutellate or subdiscoid, reaching a diameter of 2–4 mm., externally reddish-brown or liver-colored and minutely granular, the margin becoming irregularly crenate; hymenium concave or nearly plane, similar in color to the outside of the apothecium; excipular cells rounded or polygonal and reaching a diameter of 30–40 μ, the walls pale-brown or with a purplish tinge; asci broad-cylindric, reaching a length of 200 μ and a diameter of 20 μ, 8-spored; spores 1-seriate or irregularly crowded, ellipsoid, smooth, hyaline, rather thick-walled, 10–12 × 22–26 μ; paraphyses septate, enlarged at the septa, giving them a knotted appearance, enlarged at their apices, where they reach a diameter of 6 μ, pale-brown or purplish.

On soil and fire clay.

TYPE LOCALITY: Europe.

DISTRIBUTION: Colorado; also in Europe.

ILLUSTRATIONS: Batsch, Elench. Fung. Contin. **1**: *pl. 26, f. 138;* Ann. Mag. Nat. Hist. III. **15**: *pl. 15, f. 19;* Cooke, Mycographia *pl. 22, f. 85;* Pat. Tab. Anal. Fung. *f. 485.*

32. **Humarina trachyderma** (Ellis & Ev.) Seaver, comb. nov.

?Peziza tapeina Cooke, Mycographia 157. 1877.
?Peziza pertenuis Cooke, Mycographia 157. (as synonym). 1877.
Peziza trachyderma Ellis & Ev. Am. Nat. **31**: 426. 1897.
Humaria trachyderma Sacc. & Syd. in Sacc. Syll. Fung. **14**: 752. 1899.

Apothecia sessile, becoming shallow cup-shaped or nearly plane, externally furfuraceous-verrucose, when fresh nearly the

color of the substratum, with the hymenium darker when dry, margin incurved, reaching a diameter of 2–4 mm.; asci cylindric or subcylindric, about 12–14 × 200 μ, 8-spored; spores 1-seriate, ellipsoid, hyaline, smooth, 10–12 × 18–20 μ; paraphyses thickened at their apices.

On decaying and partially buried wood.

TYPE LOCALITY: Valentine, Nebraska.

DISTRIBUTION: Nebraska and ?South Carolina.

ILLUSTRATION: Cooke, Mycographia *pl. 70, f. 268.*

DOUBTFUL AND EXCLUDED SPECIES

Humaria bella (Berk. & Curt.) Sacc. Syll. Fung. **8**: 135. 1889; *Peziza bella* Berk. & Curt.; Berk. Jour. Linn. Soc. **10**: 366. 1868. This species is reported from Cuba by Berkeley and Curtis. No specimens have been seen, but the general description of the species suggests *Humarina leucoloma* (Hedw.) Seaver.

Humaria scatigena (Berk. & Curt.) Sacc. Syll. Fung. **8**: 147. 1889; *Peziza scatigena* Berk. & Curt.; Berk. Jour. Linn. Soc. **10**: 366. 1868. This species is also reported from Cuba by Berkeley and Curtis. No specimens have been seen, but both the description and habitat suggest an *Ascobolus.*

Humaria salmonicolor (Berk. & Br.) Sacc. Syll. Fung. **8**: 123. 1889; *Peziza salmonicolor* Berk. & Br. Ann. Mag. Nat. Hist. III. **18**: 124. 1866. This species is reported from Nebraska by Clements. No specimens have been found available for study.

Humaria olivatra (Ellis & Holw.) Sacc. Syll. Fung. **8**: 148. 1889; *Peziza olivatra* Ellis & Holw.; Arth. Bull. Geol. Nat. Hist. Surv. Minn. **3**: 36. 1887. This is an inoperculate species and therefore not a member of the present genus, which belongs to the operculate section of discomycetes.

Humaria fuscocarpa (Ellis & Holw.) Morgan, Jour. Myc. **8**: 189. 1902; *Peziza fuscocarpa* Ellis & Holw. Jour. Myc. **1**: 5. 1885. This also belongs to the inoperculate series of discomycetes. It is apparently a synonym of *Peziza olivacea* Batsch and belongs to the monotypic genus *Catinella* established by Boudier in 1907.

Humaria vitigena Massee & Morgan; Morgan, Jour. Myc. **8**: 189. 1902. This is not a member of this genus and probably also belongs with the inoperculate section of discomycetes.

Humaria subcrenulata Clements, Bot. Surv. Nebr. **4**: 11. 1896. The type of this species could not be located, so that the identity of the species is uncertain.

Humaria secreta (Phill.) Sacc. Syll. Fung. **8**: 123. 1889; *Peziza secreta* Phill. Grevillea **7**: 21. 1878. This species was described from material collected in California by Harkness. We have no knowledge of the species except that gained from the original description.

Humaria groenlandica Rostr. Medd. Groenland **18**: 55. 1896. This small blackish species is reported from Jamesons Land, Greenland. No specimen has been seen, but the description suggests a close relationship with *Peziza spissa* Berk.

Humaria microspora (Berk. & Curt.) Sacc. Syll. Fung. **8**: 131. 1889; *Peziza microspora* Berk. & Curt.; Berk. Grevillea **3**: 150. 1875. This species was first reported from South Carolina. Specimens collected in Pennsylvania which appear to agree with the original description are more closely related to the genus *Plectania* than to *Humarina*.

Humaria xanthomela americana (Rehm) Sacc. & Trott. in Sacc. Syll. Fung. **22**: 626. 1913; *Melachroia xanthomela americana* Rehm, Ann. Myc. **3**: 230. 1905. The American form of this species was reported by Rehm from Tacoma Park, Washington, D. C. No specimen has been seen.

Peziza elachroa Berk. & Curt.; Cooke, Mycographia 160. 1877. The species which is described from Cuban material is about one-fourth of an inch broad and of an olivaceous color. No specimen has been seen.

Humaria subgranulata (Berk. & Curt.) Sacc. Syll. Fung. **8**: 129. 1889; *Peziza subgranulata* Berk. & Curt.; Berk. Jour. Linn. Soc. **10**: 366. 1868. A Cuban species said to be closely related to *Ascophanus granulatus*.

18. **PSEUDOMBROPHILA** Boud. Hist. Class. Discom. Eu. 65. 1907.

Apothecia subsessile or short-stipitate, less than 1 cm. in diameter; substance rather tough and resembling an *Ombrophila;* asci cylindric or subcylindric, usually 8-spored; spores ellipsoid, hyaline; paraphyses delicate.

Type species, *Helotium Pedrottii* Bres.

1. **Pseudombrophila deerrata** (Karst.) Seaver, comb. nov.

Peziza deerrata Karst. Not. Fauna Fl. Fenn. **10**: 119. 1869.
Peziza rufescens Schröt.; Rab. Fungi Eu. *2311.* 1878.
Peziza Schröteri Cooke, Grevillea **6**: 110. 1878.
Helotium Pedrottii Bres. Fungi Trid. **1**: 14. 1881.
Humaria deerrata Sacc. Syll. Fung. **8**: 140. 1889.
Geoscypha Schröteri Rehm, Ascom. *901;* Hedwigia **27**: 163. 1888.
Humaria Schröteri Sacc. Syll. Fung. **8**: 141. 1889.
Pseudombrophila Pedrottii Boud. Hist. Class. Discom. Eu. 65. 1907.

Apothecia gregarious or crowded, sessile or short-stipitate, at first closed and subglobose, opening and becoming discoid or shallow cup-shaped, reaching a diameter of 3–4 mm., externally dark-brownish and subtomentose or squamulose; hymenium plane or slightly concave, pale-reddish or reddish-purple; asci cylindric or subcylindric, 8-spored; spores ellipsoid, 1-seriate, parallel with the ascus and apparently adhering together but easily detached, occasionally containing one central oil-drop, hyaline or slightly yellowish, 8–9 × 12–14 μ; paraphyses slender, rather strongly enlarged above, where they reach a diameter of 4 μ.

On dead herbaceous stems, especially on Brassicaceae.

TYPE LOCALITY: Finland.

DISTRIBUTION: North Dakota; also in Europe.

ILLUSTRATIONS: Boud. Ic. Myc. *pl. 390;* Bres. Fungi Trid. **1**: *pl. 15;* Cooke, Mycographia *pl. 110, f. 393.*

EXSICCATI: Brenckle, Fungi Dakotenses *140.*

19. STREPTOTHECA Vuill. Jour. de Bot. **1**: 33. 1887.

Ascobolus § *Ascozonus* Renny, Jour. Bot. **12**: 355. 1874.
Ascozonus Boud. Hist. Class. Discom. Eu. 79. 1907.

Apothecia minute, scarcely visible to the unaided eye, externally naked or clothed with poorly developed hyaline hairs, white or whitish; hymenium plane or convex, roughened by the large protruding asci; asci broad-clavate to ovoid, with a distinct collar near their apices, more or less attenuated at the apex, opening by a transverse slit which gives the open ascus a bilabiate appearance, many-spored; spores hyaline or subhyaline, small, ellipsoid, often enveloped in a mucilaginous material; paraphyses slender, scant.

Type species, *Streptotheca Boudieri* Vuill.

Asci 32-spored.	1. *S. Crouani.*
Asci 64-spored.	
Spores 6–7 × 12–14 μ.	2. *S. woolhopensis.*
Spores 4 × 8 μ.	3. *S. obscura.*

1. Streptotheca Crouani (Renny) Seaver, comb. nov.

Ascobolus Crouani Renny, Jour. Bot. **12**: 356. 1874.
Streptotheca Boudieri Vuill. Jour. de Bot. **1**: 34. 1887.
Ryparobius Crouani Sacc. Syll. Fung. **8**: 543. 1889.
Ascozonus Boudieri Boud. Hist. Class. Discom. Eu. 79. 1907.
Ascozonus Crouani Boud. Hist. Class. Discom. Eu. 79. 1907.

Apothecia minute, scarcely visible to the unaided eye, gregarious or widely scattered, at first subcylindric, becoming expanded and more or less discoid, not exceeding 0.5 mm. in diameter, white or whitish, externally clothed with a few hair-like hyphae; hymenium plane or convex, of about the same color as the outside of the apothecium; asci clavate, marked near the apex with a distinct thickened ring which projects inward, 32-spored, reaching a length of 120 μ and a diameter of 22 μ, opening by a transverse slit; spores ellipsoid, smooth, hyaline, 3 × 12 μ; paraphyses very slender.

On dung of different kinds.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York; also in Europe.

ILLUSTRATIONS: Jour. de Bot. 1: *34, f. A–G;* E. & P. Nat. Pfl. 1[1]: 191, *f. 153, A–B;* Jour. Bot. 12: *pl. 154, f. 6–10.*

2. **Streptotheca woolhopensis** (Renny) Seaver, comb. nov. (PLATE 12, FIG. 2–3.)

Ryparobius woolhopensis Renny; Berk. & Br. Ann. Mag. Nat. Hist. IV. **11**: 348. 1873.

Ascobolus woolhopensis Renny, Jour. Bot. **12**: 356. 1874.

Apothecia minute, less than 1 mm. in diameter, white, sessile or subsessile, externally clothed with minute whitish hairs; hymenium plane or nearly so, roughened by the large protruding asci; hairs blunt, thin-walled, septate, hyaline, reaching a length of 40–60 μ and a diameter of 8–20 μ; asci clavate, attenuated above into a rather acute apex, short-stipitate, reaching a length of 100–200 μ and a diameter of 20–22 μ (broader when under pressure), 64-spored; spores at first collected into a rather compact mass, becoming more or less separated in the older asci, ellipsoid to fusoid, hyaline, 6–7 $\times$ 12–14 μ; paraphyses rather stout, hyaline.

On the dung of birds and mice.

TYPE LOCALITY: Great Britain.

DISTRIBUTION: New York; also in Europe.

ILLUSTRATIONS: Ann. Mag. Nat. Hist. IV. **11**: *pl. 9, f. 12;* Jour. Bot. **12**: *pl. 153.*

3. **Streptotheca obscura** Seaver, sp. nov. (PLATE 12, FIG. 4.)

Apothecia minute, scarcely visible to the naked eye, cinereous under the lens and scarcely differing in color from the substratum; hymenium convex, dull; excipular cells irregular and the walls pale yellowish-brown; asci broad, clavate, stipitate below, slightly constricted above near the apex, the constriction marking the position of the ring or collar, reaching a length of 80 μ and a diameter of 20–22 μ, 64-spored; spores ellipsoid, hyaline, 4 $\times$ 8 μ; paraphyses slender, septate.

Type collected on rabbit dung at Tarrytown, New York, May, 1916 (Herb. The New York Botanical Garden).

DISTRIBUTION: Known only from the type locality.

20. **RYPAROBIUS** Boud. Ann. Sci. Nat. V. **10**: 237. 1869.

?Thelebolus Tode, Fungi Meckl. **1**: 41. 1790.
Pezizula Karst. Myc. Fenn. **1**: 81. 1871.

Apothecia minute, often scarcely visible, sessile, externally smooth or clothed with poorly developed hairs, white to whitish or yellowish; excipulum often remaining closed until ruptured by the maturing asci; hymenium concave, plane or more often convex, usually roughened by the protruding asci; asci large, broad-clavate to ovoid, opening by a circular lid or in a few cases the method of dehiscence uncertain, many-spored; spores hyaline or subhyaline, small, ellipsoid, often enveloped in a mucilaginous material with which they are discharged from the ascus; paraphyses usually very delicate.

Type species, *Ryparobius brunneus* Boud.

Asci 16-spored.	1. *R. sexdecimsporus.*
Asci 64-spored.	2. *R. crustaceus.*
Asci 72-spored.	3. *R. hyalinellus.*
Asci more than 72-spored.	
Apothecia always having more than one ascus.	4. *R. polysporus.*
Apothecia usually having a single ascus.	5. *R. monascus.*

1. **Ryparobius sexdecimsporus** (Crouan) Sacc. Syll. Fung. **8**: 541. 1889. (PLATE 12, FIG. 1.)

Ascobolus sexdecimsporus Crouan, Ann. Sci. Nat. IV. **10**: 195. 1858.
Ascophanus sexdecimsporus Boud. Ann. Sci. Nat. V. **10**: 247. 1869.

Apothecia scattered or gregarious, sessile, at first subglobose, expanding and becoming subdiscoid, smooth, at first white or whitish, finally dull, pallid or yellowish, reaching a diameter of 1–2 mm., or occasionally less than 1 mm.; hymenium at maturity plane or convex, roughened by the protruding asci, similar in color to the outside of the apothecium; excipular cells irregularly polygonal, reaching a diameter of 12–15 μ, giving rise above to a palisade of mycelium with loose, club-shaped ends; asci broadly clavate, reaching a length of 75–125 μ and a diameter of 20–25 μ, tapering below into a short stem-like base, 16-spored; spores 2-seriate or becoming irregularly crowded, ellipsoid, containing one central oil-drop, 7–10 $\times$ 11–14 μ, hyaline; paraphyses very slender, not usually exceeding 2 μ in diameter and scarcely thickened above.

On dung of various kinds.

TYPE LOCALITY: France.

RYPAROBIUS, STREPTOTHECA, AND THECOTHEUS

DISTRIBUTION: New York to North Dakota; also in Europe.

ILLUSTRATIONS: Ann. Sci. Nat. IV. **10**: *pl. 13, f. 21–26;* Ann. Sci. Nat. V. **10**: *pl. 11, f. 35;* Ann. Mag. Nat. Hist. III. **15**: *pl. 17, f. 35;* Rab. Krypt.-Fl. 1^3: 1082, *f. 5;* E. & P. Nat. Pfl. 1^1: 189, *f. 152, E–F;* Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 34, f. 1.*

2. **Ryparobius crustaceus** (Fuckel) Rehm, Ber. Naturh. Ver. Augsburg **26**: 17. 1881.

Ascobolus crustaceus Fuckel, Hedwigia **5**: 4. 1866.
Ascobolus Cookei Crouan, Fl. Finist. 56. 1867.
Ascobolus polysporus Auersw. Hedwigia **7**: 51. 1868.
Ryparobius Cookei Boud. Ann. Sci. Nat. V. **10**: 238. 1869.
Pezizula crustacea Karst. Myc. Fenn. **1**: 81. 1871.

Apothecia gregarious or scattered, very minute and scarcely visible to the unaided eye, at first subglobose, then becoming expanded and more or less discoid, at first white or whitish, becoming dull-pallid or dirty-white, not usually exceeding 0.5 mm. in diameter; hymenium convex and roughened by the strongly protruding asci; asci comparatively few to each plant and very broad-clavate to subovoid, many-spored, the exact number of the spores difficult to make out with certainty, but usually 64, entire ascus reaching a length of 90–100 μ and a diameter of about 25 μ, rather abruptly tapering below into a short stem-like base; spores ellipsoid, crowded together in the ascus with no apparent order, hyaline, often with one small oil-drop or with none apparent, above 4–6 $\times$ 7–10 μ; paraphyses sparse, slender and very slightly thickened above.

On dung of different kinds.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to Colorado; also in Europe.

ILLUSTRATIONS: Ann. Sci. Nat. V. **10**: *pl. 9, f. 24;* Hedwigia **5**: *pl. 1, f. 6;* Crouan Fl. Finist. pl. (Suppl.), *f. 3;* Massee, Brit. Fungus-Fl. **4**: 156, *f. 34–35;* Phill. Brit. Discom. *pl. 9, f. 57;* Rab. Krypt.-Fl. 1^3: 1083, *f. 1–4;* Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 33, f. 2.*

3. **Ryparobius hyalinellus** (Karst.) Sacc. Syll. Fung. **8**: 542. 1889.

Peziza hyalinella Karst. Myc. Fenn. **1**: 83. 1871.

Apothecia gregarious, subglobose, smooth, whitish, reaching a diameter of 100 μ; hymenium strongly convex, similar in color

to the outside of the apothecium; asci clavate, reaching a length of 55–100 μ and a diameter 18–28 μ, 72-spored; spores ellipsoid, hyaline, reaching a length of 5 μ and a diameter of 3 μ; paraphyses slender, enlarged above, reaching a diameter of 6 μ at their apices.

On bird dung.

TYPE LOCALITY: Finland.

DISTRIBUTION: Greenland; also in Europe.

4. **Ryparobius polysporus** (Karst.) Sacc. Syll. Fung. **8**: 539. 1889. (PLATE 12, FIG. 5.)

Ascobolus polysporus Karst. Fungi Fenn. *655*. 1867; Not. Fauna Fl. Fenn. **11**: 208. 1870.
Peziza polyspora Karst. Myc. Fenn. **1**: 82. 1871.
Ryparobius pachyascus Zukal; Rehm, Ascom. *914;* Hedwigia **27**: 167. 1888.
Ascozonus oligoascos Heimerl, Niederöst. Ascoboleen 27. 1889.

Apothecia gregarious or scattered, subglobose, becoming more or less expanded at the top, very minute, scarcely visible to the unaided eye and difficult to find even with a lens, often partially immersed in the substratum, yellowish to yellowish-brown, not exceeding 100–150 μ in diameter; excipulum for a long time entirely enclosing the apothecium, finally rupturing irregularly and exposing the asci and paraphyses, which constitute a poorly developed hymenium; asci comparatively few to each plant, not usually exceeding 3–5, broad-clavate to subovoid, the small end directed downward, reaching a length of 70–150 μ (often stretching when mature) and a diameter of 35–50 μ, with a faint suggestion of a ring near the apex, many-spored; spores irregularly disposed, small, ellipsoid, the exact number in each ascus difficult to determine with certainty, 3 × 5–7 μ; paraphyses minute and indistinct.

On dung of different kinds, especially abundant on goat dung.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to Colorado; also in Europe.

ILLUSTRATIONS: Rab. Krypt.-Fl. 1^3: 1083, *f. 1–3;* Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 34, f. 2;* Heimerl, Niederöst. Ascoboleen *pl. 1, f. 1.*

5. **Ryparobius monascus** Mouton, Bull. Soc. Royale Bot. Belg. **25**: 141. 1886. (PLATE 12, FIG. 6.)

?Thelebolus stercoreus Tode, Fungi Meckl. **1**: 41. 1790.
Thelebolus nanus Heimerl, Niederöst. Ascoboleen 30. 1889.
Thelebolus monascus Boud. Hist. Class. Discom. Eu. 79. 1907.
?Thelebolus lignicola Lloyd, Myc. Notes **52**: 738. 1917.

Apothecia gregarious or scattered, at first globose, then broad-ovoid, externally smooth, yellowish to brown, entirely closed, finally bursting open by a rupture of the excipulum through which the ascus protrudes, reaching a diameter of 200–300 μ; asci usually one to each apothecium, but occasionally two or three, ellipsoid to ovoid, often slightly constricted near the upper end and with a faint suggestion of a ring, reaching a length of 200–260 μ and a diameter of 150–175 μ, each containing a large number of spores, the exact number difficult to determine but exceeding one hundred, method of dehiscence uncertain since they do not dehisce readily but usually burst open irregularly; spores ellipsoid, hyaline, 3–3.5 $\times$ 5–7 μ; paraphyses slender, septate.

On dung of various kinds, especially abundant on goat dung; more rarely on wood.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York; also in Europe.

ILLUSTRATIONS: ?Tode, Fungi Meckl. **1**: *pl. 7, f. 56;* Rab. Krypt.-Fl. **1**^{3}: 1080, *f. 1–4;* Ann. Bot. **15**: *pl. 18, f. 41–44, 66;* Heimerl, Niederöst. Ascoboleen *pl. 1, f. 2.*

This and the preceding species differ from the other members of the genus examined in that they are for a long time enclosed in a surrounding sheath here referred to as the excipulum. Perhaps these forms should have been referred to the genus *Thelebolus* in which there is no true apothecium, although they are referred to as such by systematists.

DOUBTFUL AND EXCLUDED SPECIES

Ryparobius niveus (Fuckel) Rehm in Rab. Krypt.-Fl. **1**^{3}: 1102. 1896; *Ascobolus niveus* Fuckel, Hedwigia **5**: 4. 1886. This species has been reported for North America by Dodge. No specimens have been seen. The species is closely related to *R. crustaceus* (Fuckel) Rehm.

21. **THECOTHEUS** Boud. Ann. Sci. Nat. V. **10**: 235. 1869.

Apothecia sessile, short-cylindric, at least when young, white or whitish, minute, substance soft and waxy; asci very large,

broad-clavate, usually truncate at the apex, many-spored; spores ellipsoid or subellipsoid, smooth, each usually surrounded by a mucilaginous envelope; paraphyses present, slender.

Type species, *Ascobolus Pelletieri* Crouan.

1. **Thecotheus Pelletieri** (Crouan) Boud. Ann. Sci. Nat. V. **10**: 236. 1869. (PLATE 12, FIG. 7.)

Ascobolus Pelletieri Crouan, Ann. Sci. Nat. IV. **7**: 173. 1857.
Ryparobius Pelletieri Sacc. Syll. Fung. **8**: 542. 1889.

Apothecia gregarious or scattered, sessile, at first subconical, the top gradually expanding, finally short-cylindric or rarely approaching discoid, reaching a diameter of 1–2 mm. and of about the same height, white or whitish, externally more or less pruinose; hymenium plane or slightly convex, similar in color to the outside of the apothecium, roughened by the protruding asci; the excipular cells roundish, loosely united, reaching a diameter of 20 μ, the cell walls hyaline; asci comparatively few to each apothecium, very large, strongly exserted and easily visible with the hand lens, broad-cylindric to clavate, tapering below into a stem-like base, reaching a length of 300–350 μ and a diameter of 50–60 μ, 32-spored; spores 3–4-seriate or irregularly disposed, ellipsoid, the ends strongly narrowed, containing one central nucleus-like body or a number of small oil-drops, 23–24 $\times$ 35–38 μ, each surrounded by a mucilaginous envelope which is rather thick and of irregular outline; paraphyses very slender.

On dung of various kinds.

TYPE LOCALITY: France.

DISTRIBUTION: Massachusetts to North Dakota and Bermuda; also in Europe.

ILLUSTRATIONS: Ann. Sci. Nat. IV. **7**: *pl. 4, f. A;* Ann. Sci. Nat. V. **10**: *pl. 9, f. 22;* E. & P. Nat. Pfl. 1^1: 189, *f. 152, G–J;* Massee, Brit. Fungus-Fl. **4**: 156, *f. 38;* Rab. Krypt.-Fl. 1^3: 1082, *f. 1–4;* Phill. Brit. Discom. *pl. 9, f. 56;* Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 33, f. 1;* Pat. Tab. Fung. *f. 172.*

22. **SEPULTARIA** Massee, Brit. Fungus-Fl. **4**: 389. 1895.

Peziza § *Sepultaria* Cooke, Mycographia 259. 1879.

Apothecia medium-sized, reaching a diameter of 2–3 cm., cup-shaped or rarely subscutellate with age, clothed with flexuous or bristly hairs, partially to entirely buried in the ground, often becoming subsuperficial with age, entire or splitting *Geaster-*

like; hymenium concave, whitish or orange to red; asci 8-spored; spores ellipsoid; paraphyses present and variable.

Type species, *Peziza sepulta* Fries.

Apothecia large, 1–4 cm. in diameter, entirely buried and appearing as a hole in the ground.
- Spores ellipsoid, 15–17 × 25–30 μ. — 1. *S. arenicola.*
- Spores subglobose, 18–20 × 20–22 μ. — 2. *S. Longii.*

Apothecia medium-sized, seldom exceeding 1 cm., partially buried.
- Hymenium white or whitish. — 3. *S. arenosa.*
- Hymenium orange or reddish.
  - Spores 8–10 × 18–25 μ; hairs rigid. — 4. *S. aurantiaca.*
  - Spores 12–14 × 25–33 μ; hairs entirely flexuous. — 5. *S. pellita.*

1. Sepultaria arenicola (Lév.) Massee, Brit. Fungus-Fl. **4**: 390. 1895.

?Peziza lanuginosa Bull. Herb. Fr. *pl. 396, f. 2.* 1788.
Peziza arenicola Lév. Ann. Sci. Nat. III. **9**: 140. 1848.
Peziza sepulta Fries; Berk. & Br. Ann. Mag. Nat. Hist. II. **13**: 463. 1854.
Peziza Geaster Berk. & Br. Ann. Mag. Nat. Hist. III. **18**: 125. 1866.
Peziza lanuginosa Sumneri Berk. Trans. Linn. Soc. **25**: 432. 1866.
Lachnea arenicola Gill. Champ. Fr. Discom. 77. 1879.
Lachnea Sumneriana Phill. Brit. Discom. 213. 1887.
Lachnea lanuginosa Sacc. Syll. Fung. **8**: 167. 1889.
Lachnea Geaster Sacc. Syll. Fung. **8**: 172. 1889.
Lachnea sepulta Sacc. Syll. Fung. **8**: 170. 1889.
Scutellinia Geaster Kuntze, Rev. Gen. Pl. **2**: 869. 1891.
Scutellinia sepulta Kuntze, Rev. Gen. Pl. **2**: 869. 1891.
Scutellinia Sumneriana Kuntze, Rev. Gen. Pl. **2**: 869. 1891.
Scutellinia arenicola Kuntze, Rev. Gen. Pl. **2**: 869. 1891.
?Scutellinia lanuginosa Kuntze, Rev. Gen. Pl. **2**: 869. 1891.
Sarcosphaera arenicola Lindau in E. & P. Nat. Pfl. **1**[1]: 182. 1897.
Sarcosphaera Sumneriana Lindau in E. & P. Nat. Pfl. **1**[1]: 182. 1897.
Sarcosphaera Geaster Lindau in E. & P. Nat. Pfl. **1**[1]: 182. 1897.
Sepultaria heterothrix Clements, Bull. Torrey Club **30**: 91. 1903.
Lachnea heterothrix Sacc. & D. Sacc. in Sacc. Syll. Fung. **18**: 33. 1906.
Sepultaria Sumneri Boud. Hist. Class. Discom. Eu. 59. 1907.
Sepultaria sepulta Boud. Hist. Class. Discom. Eu. 59. 1907.
Sepultaria Geaster Boud. Hist. Class. Discom. Eu. 59. 1907.

Apothecia gregarious, at first closed and entirely immersed in the ground, at maturity opening and appearing as a hole in the ground, sometimes becoming semisuperficial, margin incurved and entire or splitting irregularly, reaching a diameter of 1–4 cm., externally brown and densely clothed with brown flexuous hairs which bind the substratum together so that the whole exterior of the apothecium is encrusted with sand, the exterior of the cup

woolly when the soil has been washed away; hymenium pallid, becoming yellowish or brownish when old or when dried; hairs thin-walled, sparingly septate, often branched, reaching a diameter of 10–15 μ, externally minutely roughened; asci cylindric or subcylindric, reaching a diameter of 20 μ; spores ellipsoid, often narrowed at the ends, smooth, usually containing one large oil-drop, or rarely two, 12–17 $\times$ 25–30 μ, usually about 12 $\times$ 25 μ; paraphyses slender, slightly enlarged at their apices.

On rather sandy soil.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to Colorado, and California; also in Europe and Africa.

ILLUSTRATIONS: Cooke, Mycographia *pl. 29, f. 111, 112, 114;* Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 9, f. 2;* Trans. Linn. Soc. **25**: *pl. 55, f. 1;* Boud. Ic. Myc. *p*. *358;* Phill. Brit. Discom. *pl. 6, f. 39;* Rab. Krypt.-Fl. 1^3: 1036, *f. 1–5;* E. & P. Nat. Pfl. 1^1: 181, *f. 147, H;* Ann. Mag. Nat. Hist. III. **18**: *pl. 4, f. 25–27.*

EXSICCATI: Fungi Columb. *2377.*

Since this manuscript was submitted to the printers, the writer has received from Dr. John Axel Nannfeldt of Sweden a collection, under the name of *Sepultaria sepulta* (Fries) Rehm, which he reports growing abundantly on (or rather in or under) the sidewalks in a suburb of Upsala just outside the Botanical Garden and only a few hundred meters from the type locality. He also states that it looks very insignificant like holes from earthworms.

Nannfeldt also states that after examination of the authentic specimens of *Peziza arenosa* Fuckel, he is convinced that this is identical with the above. *Sepultaria arenosa* has been treated as distinct in this work, the American specimens referred to this species having smaller apothecia which are only partially submerged. It is possible that this is only a form of *Sepultaria arenicola.*

2. **Sepultaria Longii** Seaver, Mycologia **7**: 199. 1915.

Apothecia densely gregarious or cespitose, at first closed and entirely buried, finally opening to the surface by an elongated or compressed aperture, or when the substratum is not too compact spreading so as to expose the hymenium, never protruding above the surface of the substratum but causing the soil to become



SEPULTARIA LONGII

slightly elevated as they mature, reaching a diameter of 4 cm., regular in form or becoming very much contorted from mutual pressure, clothed with hairs which extend into the substratum, binding the surrounding soil to the outside of the apothecium; hymenium deeply concave, nearly smooth, almost white; hairs flexuous, septate, brown, and of nearly uniform thickness throughout their entire length; asci subcylindric above, tapering gradually below into a stem-like base, reaching a length of 250–300 μ and a diameter of 20–22 μ; spores 1-seriate, short-ellipsoid or subglobose, at first containing one small oil-drop which gradually enlarges until it nearly fills the spore, about 18–20 × 20–22 μ; paraphyses stout, gradually enlarged above, where they reach a diameter of 4–6 μ, filled with numerous vacuoles or oil-drops, hyaline.

On bare ground.

TYPE LOCALITY: Albuquerque, New Mexico.

DISTRIBUTION: Known only from the type locality.

ILLUSTRATION: Mycologia **7**: *pl. 161.*

3. **Sepultaria arenosa** (Fuckel) Boud. Hist. Class. Discom. Eu. 59. 1907.

Peziza arenosa Fuckel, Fung. Rhen. *1212.* 1865; Hedwigia **5**: 15. 1866.
Humaria arenosa Fuckel, Symb. Myc. 321. 1869.
Lachnea arenosa Sacc. Syll. Fung. **8**: 167. 1889.
Scutellinia arenosa Kuntze, Rev. Gen. Pl. **2**: 869. 1891.
?Sepultaria grisea Clements, Bot. Surv. Nebr. **4**: 13. 1896.
?Lachnea grisea Sacc. & Sydow in Sacc. Syll. Fung. **14**: 756. 1899.
Sarcosphaera arenosa Lindau in E. & P. Nat. Pfl. **1**[1]: 182. 1897.

Apothecia scattered, sessile and partially immersed in the ground, finally becoming subsuperficial, cup-shaped or occasionally subscutellate, reaching a diameter of 1 cm., the margin entire or splitting, externally clothed with hairs which are often tufted; hymenium white or whitish; hairs flexuous or crooked, more rarely straight and subrigid on the exposed parts, often swollen at the base, reaching a length of 200 μ and a diameter of 9 μ, the hairs on the lower part of the cup giving rise to rhizoid-like hyphae which extend into the surrounding soil; asci subcylindric above, reaching a length of 200–225 μ and a diameter of 15–18 μ; spores 1-seriate or partially 2-seriate, ellipsoid, the ends slightly narrowed, usually containing two oil-drops, 13–18 × 23–25 μ, smooth; paraphyses gradually enlarged above, where they reach a diameter of 7–9 μ.

On sandy soil.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to Colorado; also in Europe.

ILLUSTRATIONS: Boud. Ic. Myc. *pl. 361;* Cooke, Mycographia *pl. 30, f. 117.*

4. **Sepultaria aurantia** Clements, Bot. Surv. Nebr. **4**: 12. 1896.

?Peziza hybrida Sow. Engl. Fungi 369. 1803.
?Scutellinia hybrida Kuntze, Rev. Gen. Pl. **2**: 869. 1891.
Lachnea aurantia Sacc. & Sydow in Sacc. Syll. Fung. **14**: 757. 1899.

Apothecia scattered or crowded, sessile, concave-hemispheric, subsuperficial, reaching a diameter of 1–2 cm., externally clothed with hairs; hymenium concave, reddish or orange; hairs flexuous near the base, those on the exposed parts rigid, septate, dark-brown; asci cylindric or subcylindric; spores 1-seriate, ellipsoid, containing one or two oil-drops, smooth, 8–10 × 18–25 μ; paraphyses slightly enlarged above, filled with orange granules.

On damp ground.

TYPE LOCALITY: Bellevue, Nebraska.

DISTRIBUTION: Known only from the type locality.

ILLUSTRATIONS: ?Sow. Engl. Fungi *pl. 369, f. 1;* Boud. Ic. Myc. *pl. 356;* Cooke, Mycographia *pl. 32, f. 126, b.*

5. **Sepultaria pellita** (Cooke & Peck) Seaver, comb. nov.

Peziza pellita Cooke & Peck, Grevillea **1**: 5. 1872.
Lachnea pellita Sacc. Syll. Fung. **8**: 169. 1889.
Scutellinia pellita Kuntze, Rev. Gen. Pl. **2**: 869. 1891.

Apothecia scattered or gregarious, sessile and at first entirely immersed in the soil, the margin of the cup appearing above the surface of the soil when expanded, entire or splitting, reaching a diameter of 1–2 cm., externally clothed with flexuous hairs which penetrate into the substratum, binding together the particles of sand and humus on which they grow; hymenium concave, yellowish to pale-orange; hairs brown, sparingly septate, of nearly uniform thickness throughout their entire length, reaching a diameter of 7–8 μ; asci cylindric above, reaching a length of 300 μ and a diameter of 20 μ; spores 1-seriate or occasionally slightly crowded in the upper part of the ascus; ellipsoid, the ends rather strongly narrowed, containing one or two large oil-drops, 12–14 × 25–33 μ; paraphyses rather strongly enlarged above, reaching a diameter of 7–8 μ, filled with yellow granules.

On sandy soil among mosses and humus.

TYPE LOCALITY: Adirondack Mountains.

DISTRIBUTION: New York to Wisconsin.

ILLUSTRATIONS: Grevillea 1: *pl. 1, f. 3;* Cooke, Mycographia *pl. 31, f. 119.*

23. **PSEUDOPITHYELLA** Seaver, Mycologia **19**: 87. 1927.

Apothecia stipitate or subsessile, rather tough, not shrinking much in drying, in general appearance and in habitat resembling a *Pithya;* hymenium scarlet; asci 8-spored, marked near the apex by a prominent thickened ring; spores ellipsoid, hyaline; paraphyses filiform.

Type species, *Sarcoscypha minuscula* Boud. & Torrend.

1. **Pseudopithyella minuscula** (Boud. & Torrend) Seaver, comb. nov.

Sarcoscypha minuscula Boud. & Torrend, Bull. Soc. Myc. Fr. **27**: 128. 1911.

Apothecia minute, not exceeding 1–2 mm. in diameter, usually stipitate, at first nearly globose, becoming discoid or subdiscoid, externally whitish and more or less tomentose; hymenium concave or nearly plane, scarlet, becoming paler in dried plants; stem slender and very variable in length, when springing from the under side of the substratum, short or almost wanting, seldom exceeding 1 mm. in length, gradually expanding above into the apothecium; asci cylindric or subcylindric above, gradually tapering below into a long stem-like base, reaching a length of 600 μ and a diameter of 12–15 μ, marked by two little ear-like protuberances where the thickened ring passes around near the apex, the point of dehiscence about 3 μ inside of the ring; operculum circular, 3–4 μ in diameter; spores smooth, containing two oil-drops, 10–11 $\times$ 15–17 μ; paraphyses filiform, scarcely enlarged above, branched, reaching a diameter of 2 μ at their apices.

On decaying foliage of cedar.

TYPE LOCALITY: Portugal.

DISTRIBUTION: Bermuda Islands; also in Europe.

ILLUSTRATION: Bull. Soc. Myc. Fr. **27**: *pl. 4, f. 3.*

The only specimens of this species known on this side of the Atlantic were collected by the writer, and later by H. H. Whetzel, in Bermuda. Our specimens agree splendidly with the illustration cited.

24. **PERROTIA** Boud. Bull. Soc. Myc. Fr. **17**: 24. 1901.

Apothecia minute, sessile, clothed with hairs, the general appearance being like that of *Lachnella* except that the asci are operculate; asci clavate, 8-spored; spores allantoid, at first simple, later apparently 1-septate, or pseudoseptate; paraphyses present.

Type species, *Peziza flammea* Alb. & Schw.

1. **Perrotia flammea** (Alb. & Schw.) Boud. Hist. Class. Discom. Eu. 66. 1907.

Peziza flammea Alb. & Schw. Consp. Fung. 319. 1805.
?Peziza cinnabarina Schw. Trans. Am. Phil. Soc. II. **4**: 173. 1832.
?Peziza maculincola Schw. Trans. Am. Phil. Soc. II. **4**: 174. 1832.
Lachnella flammea Fries, Summa Veg. Scand. 365. 1849.
Helotium flammeum Karst. Myc. Fenn. **1**: 157. 1871.
?Peziza Meleagris Ellis, Bull. Torrey Club **8**: 123. 1881.
?Lachnella cinnabarina Sacc. Syll. Fung. **8**: 399. 1889.
?Lachnella incarnescens Sacc. Syll. Fung. **8**: 399. 1889.
?Lachnella maculincola Sacc. Syll. Fung. **8**: 400. 1889.
Dasyscypha allantospora Earle in Greene, Pl. Baker. **2**: 5. 1901.
Lachnella rhoina Earle in Greene, Pl. Baker. **2**: 6. 1901.

Apothecia gregarious or crowded, at first subglobose, with the margin strongly incurved, becoming scutellate, externally clothed with cinnabar-red hairs which give to the whole apothecium a brick-red color, varying in old specimens to dark-brown, reaching a diameter of 2 mm.; hymenium concave, reddish, much paler than the outside of the apothecium; hairs flexuous, of nearly uniform thickness throughout their entire length, reaching a diameter of 3–6 μ, sparingly septate, externally densely covered with coarse red granules; asci reaching a length of 100 μ and a diameter of 10–12 μ, gradually tapering below; spores 2-seriate or irregularly crowded, hyaline, reaching a length of 20 μ and a diameter of 3–4 μ; paraphyses rather stout, reaching a thickness of 4 μ.

On dead branches and twigs of deciduous and coniferous trees.

TYPE LOCALITY: Europe.

DISTRIBUTION: Pennsylvania to Montana and Colorado; also in Europe.

ILLUSTRATIONS: Alb. & Schw. Consp. Fung. *pl. 1, f. 6;* Boud. Ic. Myc. *pl. 321;* E. & P. Nat. Pfl. 1^1: 202, *pl. 159, F–G;* Rab. Krypt.-Fl. 1^3: 828, *f. 1–5.*

EXSICCATI: N. Am. Fungi *2323;* Fungi Columb. *1221, 2733.*

25. **LASIOBOLUS** Sacc. Bot. Centr. **18**: 220. 1884.

Apothecia becoming scutellate to discoid, externally clothed with very conspicuous hairs; substance soft and fleshy or waxy; hairs hyaline or subhyaline (white to the naked eye), rigid, non-septate; asci strongly protruding, cylindric to clavate, 8-spored; spores ellipsoid, hyaline; paraphyses filiform to clavate, simple or branched.

Type species, *Peziza papillata* Pers.

Apothecia hyaline to pale-orange; spores 12–14 × 20–24 μ. 1. *L. equinus.*
Apothecia bright-red; spores 12–14 × 24–30 μ. 2. *L. ruber.*

1. **Lasiobolus equinus** (Müll.) Karst. Act. Soc. Fauna Fl. Fenn. **2**: 122. 1885.

Elvela equina Müll. Fl. Dan. **13**: 8. 1778.
Peziza papillata Pers. Tent. Disp. Fung. 34. 1797.
Ascobolus ciliatus Kunze & Schmidt, Myc. Heft. **1**: 90. 1817.
Ascobolus pilosus Fries, Syst. Myc. **2**: 164. 1822.
Peziza cervina Pers. Myc. Eu. **1**: 254. 1822.
Peziza diversicolor Fries, Syst. Myc. **2**: 88. 1822.
Ascobolus papillatus Wallr. Fl. Crypt. Ger. **2**: 514. 1833.
Ascophanus ciliatus Boud. Ann. Sci. Nat. V. **10**: 254. 1869.
Ascobolus diversisporus Fuckel, Symb. Myc. 289. 1869.
Ascophanus pilosus Boud. Ann. Sci. Nat. V. **10**: 254. 1869.
Humaria diversispora Speg. Michelia **1**: 236. 1878.
Lasiobolus pilosus Sacc. Bot. Centr. **18**: 220. 1884.
Ascophanus pilosus ciliatus Phill. Brit. Discom. 312. 1887.
Lasiobolus diversisporus Sacc. Syll. Fung. **8**: 538. 1889.

Apothecia gregarious or crowded, often forming congested masses of considerable extent, at first closed, expanding and becoming discoid or short-cylindric, rarely reaching a diameter of 1 mm., externally clothed with very conspicuous hairs; hymenium concave, plane or slightly convex, roughened by the protruding asci, the color varying from almost hyaline to pale-yellow or orange; hairs swollen near the base, where they reach a diameter of 30–35 μ, tapering above into a sharp-pointed apex, reaching a length of 375 μ, thick-walled, non-septate, hyaline or faintly yellowish; asci cylindric or subcylindric above, tapering rather abruptly below into a stem-like base, reaching a length of 200 μ and a diameter of 20–24 μ; spores 1-seriate or irregularly crowded, smooth, hyaline, or slightly yellowish, ellipsoid, 12–14 × 20–24 μ; paraphyses slender, septate, slightly enlarged above, reaching a diameter of 2–3 μ, often branched.

On dung of various kinds.

TYPE LOCALITY: Europe.

DISTRIBUTION: Massachusetts to South Carolina, North Dakota, Colorado, California, and the Bermuda Islands; also in Europe.

ILLUSTRATIONS: Fl. Dan. *pl. 779, f. 3;* Ann. Sci. Nat. V. **10**: *pl. 12, f. 42–44;* E. & P. Nat. Pfl. 1^1: 189, *f. 152, A–C;* Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 32, f. 1;* Rab. Krypt.-Fl. 1^3: 1081, *f. 1–4;* Pat. Tab. Fung. *pl. 487.*

EXSICCATI: N. Am. Fungi *2334;* Ravenel, Fungi Car. **5**: *37*; Clements, Crypt. Form. Colo. *122.*

2. **Lasiobolus ruber** (Quél.) Sacc. Syll. Fung. **8**: 537. 1889.

Ascophanus ruber Quél. Grevillea **8**: 117. 1880.

Apothecia gregarious or crowded in small congested masses, at first subglobose, becoming discoid or subdiscoid, reaching a diameter of 0.5–1 mm., externally densely clothed with hairs; hymenium concave, plane or convex, bright-red; hairs swollen just above the base and contracted at the point of attachment, tapering above into a bristle-like apex, hyaline or subhyaline, reaching a length of 500 μ and a diameter of 30 μ; asci cylindric to clavate, reaching a length of 200 μ and a diameter of 25–30 μ, tapering rather abruptly below into a stem-like base; spores at first 1-seriate, obliquely arranged with the ends overlapping, often becoming 2-seriate or irregularly crowded, ellipsoid, 12–14 $\times$ 24–30 μ; paraphyses about 3 μ in diameter, enlarged upwards, reaching a diameter of 5–6 μ at their apices.

On cow dung.

TYPE LOCALITY: Jura Mountains.

DISTRIBUTION: Colorado; also in Europe.

ILLUSTRATIONS: Grevillea **8**: *pl. 131, f. 8;* Assoc. Fr. Av. Sci. Compte Rendu **12**: *pl. 7, f. 13;* Boud. Ic. Myc. *pl. 415.*

26. **PATELLA** Weber; Wiggers, Fl. Hols. 106. 1780.

Humaria Fuckel, Symb. Myc. 320. 1869. Not *Humaria* Sacc.

Lachnea Gill. Champ. Fr. Discom. 57. 1879. Not *Lachnaea* L. 1762. (*Lachnea*, 1753.)

Peziza § *Scutellinia* Cooke, Mycographia 260. 1879.

Peziza § *Neottiella* Cooke, Mycographia 261. 1879.

Octospora Hedwig, Descr. **2**: 4, 10. 1788. (in part.)

Ciliaria Quél.; Boud. Bull. Soc. Myc. Fr. **1**: 105. 1885. See Bull. Soc. Bot. Fr. **26**: 235. 1879. Not *Ciliaria* Stackh., 1809, nor *Ciliaria* Haworth, 1821.

Cheilymenia Boud. Bull. Soc. Myc. Fr. **1**: 105. 1885.
Tricharia Boud. Bull. Soc. Myc. Fr. **1**: 104. 1885.
Trichophaea Boud. Bull. Soc. Myc. Fr. **1**: 105. 1885.
Neottiella Sacc. Syll. Fung. **8**: 190. 1889.
Scutellinia Kuntze, Rev. Gen. Pl. **2**: 868. 1891.
Humariella Schröt. Krypt.-Fl. Schles. **3**^{2}: 36. 1893.
Pelodiscus Clements, Bot. Surv. Nebr. **5**: 7. 1901.
Neottiopezis Clements, Bull. Torrey Club **30**: 89. 1903.

Apothecia sessile, scattered, gregarious or congested, superficial, at least when fully matured, scutellate to discoid, more rarely concave-hemispheric, externally setose; hymenium red, yellow to brown, or white, concave or plane; hairs septate, hyaline or colored, flexuous or more often rigid; asci variable, usually 8-spored; spores ellipsoid to fusoid, smooth or rough, hyaline or subhyaline; paraphyses filiform or more commonly strongly clavate.

Type species, *Elvela ciliata* Schaeff.

Hymenium red, yellow, or brown, never white.
  Occurring on soil, rotten wood or decaying plant materials.
    Hairs simple (not forked or branched).
      Hairs well developed, bristly.
        Spores becoming distinctly sculptured.
          Apothecia clothed with brown hairs.
            Spores for a long time smooth, becoming only minutely rough.
              Spores 12–15 × 20–24 μ. — 1. *P. scutellata.*
              Spores 16–18 × 30 μ. — 2. *P. piliseta.*
            Spores becoming strongly roughened.
              Apothecia medium large, 5 mm. or more in diameter.
                Plants tropical; spores about 20 μ long. — 3. *P. cubensis.*
                Plants temperate; spores 23 μ or more long.
                  Occurring on soil among mosses; spores 12–14 × 23–24 μ. — 4. *P. umbrorum.*
                  Occurring on rotten wood; spores 14–16 × 22–26 μ. — 5. *P. Lusatiae.*
              Apothecia small, 1–2 mm. in diameter. — 6. *P. miniata.*
          Apothecia clothed with hyaline hairs. — 7. *P. albocincta.*
        Spores remaining permanently smooth or indistinctly and minutely sculptured.

- Hairs hyaline, white to the naked eye.
  - Spores fusiform, very large. 8. *P. ovilla.*
  - Spores ellipsoid, medium-sized.
    - Spores 12 × 18 μ. 9. *P. Hetieri.*
    - Spores 18 × 24–26 μ. 10. *P. ricciophila.*
- Hairs brown.
  - Spores small, 7–9 × 10–12 μ. 11. *P. punicea.*
  - Spores medium large, 14–22 μ long.
    - Occurring on burnt ground. 12. *P. gilva.*
    - Occurring on old wood.
      - On deciduous wood; spores 11–12 × 20–24 μ. 13. *P. setosa.*
      - On wood of *Sequoia;* spores 12–13 × 17–20 μ. 14. *P. Sequoiae.*
- Hairs poorly developed, blunt, adpressed. 15. *P. melaloma.*

Hairs stellately branched. 16. *P. crucipila.*

Occurring on the dung of animals.

- Spores permanently smooth.
  - Apothecia clothed with stellate hairs. 17. *P. stercorea.*
  - Apothecia clothed with simple hairs.
    - Apothecia medium to large, reaching a diameter of 1 cm.
      - Apothecia bright-yellow.
        - Hairs small, subhyaline. 18. *P. theleboloides.*
        - Hairs very large, dark-brown. 19. *P. coprinaria.*
      - Apothecia ochraceous-brown. 20. *P. maculosa.*
    - Apothecia small, not exceeding 1 mm. in diameter.
      - Apothecia short-cylindric. 21. *P. pulcherrima.*
      - Apothecia discoid or subdiscoid 22. *P. raripila.*
  - Spores becoming minutely sculptured. 23. *P. fimetaria.*

Hymenium white or bluish-white.

- Spores becoming sculptured.
  - Spore-sculpturing consisting of coarse tubercles.
    - Spores 22–25 μ long. 24. *P. paludosa.*
    - Spores 30–35 μ long. 25. *P. tuberculata.*
  - Spore-sculpturing consisting of minute warts or papillae.
    - Apothecia reaching a diameter of 2–3 cm. 26. *P. albida.*
    - Apothecia not exceeding a diameter of 5–10 mm.
      - Spores fusoid, 10–13 × 20–27 μ. 27. *P. gregaria.*
      - Spores ellipsoid, 8 × 15 μ. 28. *P. irregularis.*
- Spores remaining permanently smooth.
  - Apothecia exceeding 1 mm. in diameter.
    - Plants restricted to burned areas. 29. *P. abundans.*
    - Plants growing on damp soil or wood.
      - Occurring on old wood. 30. *P. Erinaceus.*
      - Occurring on damp soil. 31. *P. albospadicea.*
  - Apothecia less than 1 mm. in diameter. 32. *P. pygmaea.*

1. **Patella scutellata** (L.) Morgan, Jour. Myc. **8**: 187. 1902. (PLATE 14, FIG. 2; 45, FIG. 19.)

Peziza scutellata L. Sp. Pl. 1181. 1753.
Elvela ciliata Schaeff. Fung. Bavar. **4**: 112. 1774.
Patella ciliata Weber; Wiggers, Fl. Hols. 106. 1780.
Peziza aurantiaca Bull. Herb. Fr. *pl. 10.* 1780.
Octospora scutellata Hedw. Descr. **2**: 10. 1788.
Peziza ciliata Hoffm. Veg. Crypt. **2**: 25. 1790.
Humaria scutellata Fuckel, Symb. Myc. 321. 1869.
?Peziza laeticolor Karst. Myc. Fenn. **1**: 66. 1871.
Lachnea scutellata Gill. Champ. Fr. Discom. 75. 1879.
Scutellinia scutellata Kuntze, Rev. Gen. Pl. **2**: 869. 1891.
?Scutellinia laeticolor Kuntze, Rev. Gen. Pl. **2**: 869 1891.
Humariella scutellata Schröt. Krypt.-Fl. Schles. 3^2: 37. 1893.
Ciliaria scutellata Boud. Hist. Class. Discom. Eu. 61. 1907.

Apothecia gregarious or occasionally crowded, at first globose, closed and appearing dark-brown by reason of the hairs with which the exterior is covered, expanding and becoming scutellate at maturity, with the margin slightly elevated and fringed with hairs which may be very short or may extend beyond the margin a length of 1 mm.; hymenium bright-red, almost vermillion, fading in dried specimens to pale-yellow or almost white; hairs swollen just above the base and rather abruptly constricted below at the point of attachment, gradually tapering above to a bristly-like apex, thick-walled, several septate, reaching a diameter of 35–40 μ at the thickest point and a length of 1 mm.; asci cylindric or subcylindric, reaching a diameter of 12–15 μ and a length of 200–225 μ; spores 1-seriate or irregularly crowded at first densely filled with oil-drops and smooth, the surface finally becoming delicately sculptured, 12–15 $\times$ 20–24 μ; spore-sculpturing consisting of minute warts; paraphyses strongly enlarged above, reaching a diameter of 7–10 μ at their apices.

On rotten wood or more rarely on soil and leaf-mould.

TYPE LOCALITY: Europe.

DISTRIBUTION: Maine to Washington and Mexico; also in Europe.

ILLUSTRATIONS: Boud. Ic. Myc. *pl. 368;* Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 10, f. 1;* Bull. Herb. Fr. *pl. 10;* Schaeff. Fung. Bavar. *pl. 284;* Cooke, Mycographia *pl. 34, f. 131;* Pat. Tab. Fung. *f. 81;* E. & P. Nat. Pfl. 1^1: 181, *f. 147, A–B;* Massee, Brit. Fungus-Fl. **4**: 290, *f. 20, 21;* Rab. Krypt.-Fl. 1^3: 1034, *f. 1–4;* Sow. Engl. Fungi *pl. 24;* Gill. Champ. Fr. Discom. (suites); Bolton, Geschichte **3**: *pl. 108, f. 1.*

EXSICCATI: Clements, Crypt. Form. Colo. *119;* N. Am. Fungi *1310.*

2. **Patella piliseta** (Clem.) Seaver, comb. nov.

Pelodiscus pilisetus Clements, Bot. Surv. Nebr. **5**: 8. 1901.
Lachnea piliseta Sacc. & Sydow in Sacc. Syll. Fung. **16**: 1147. 1902.

Apothecia at first closed and subglobose, expanding and becoming scutellate to discoid, reaching a diameter of 1 cm.; hymenium plane or concave, yellowish-brown; hairs scattered, rigid, pointed, brown, 2–3-septate, attenuated at the base, reaching a length of 300–400 μ and a diameter of 25 μ; asci cylindric or subcylindric, reaching a length of 250 μ and a diameter of 25 μ; spores obliquely 1-seriate, broad-ellipsoid, densely filled with granules, smooth, becoming indistinctly sculptured, 16–18 $\times$ 30 μ; paraphyses clavate, septate, subhyaline, reaching a diameter of 8 μ at their apices.

On wet wood.

TYPE LOCALITY: Barney Creek, Keya Paha County, Nebraska.

DISTRIBUTION: Nebraska and Colorado.

3. **Patella cubensis** (Berk. & Curt.) Seaver, comb. nov.

Peziza cubensis Berk. & Curt.; Berk. Jour. Linn. Soc. **10**: 366. 1868.
Peziza stictica Berk. & Curt.; Berk. Jour. Linn. Soc. **10**: 367. 1868.
Peziza texensis Berk. & Curt.; Berk. Grevillea **3**: 154. 1875.
Lachnea cubensis Sacc. Syll. Fung. **8**: 176. 1889.
Lachnea stictica Sacc. Syll. Fung. **8**: 177. 1889.
Lachnea texensis Sacc. Syll. Fung. **8**: 182. 1889.
Scutellinia stictica Kuntze, Rev. Gen. Pl. **2**: 869. 1891.
Scutellinia texensis Kuntze, Rev. Gen. Pl. **2**: 869. 1891.

Apothecia sessile, expanding, becoming scutellate, reaching a diameter of 3–6 mm. externally, especially about the margin clothed with long bristly hairs; hymenium slightly concave or nearly plane, bright-red, fading in dried specimens; hairs thick-walled, septate, bristly, tapering above into a bristle-like apex, dark-brown, reaching a length of 1 mm. and a diameter of 25–30 μ, intermixed with much shorter hairs which are occasionally clavate at the apex; asci cylindric, about 14–15 $\times$ 200–220 μ; spores 1-seriate, ellipsoid, hyaline, becoming sculptured, 11–13 $\times$ 16–21 μ; spore-sculpturing consisting of rather coarse warts; paraphyses slender below, strongly enlarged at their apices.

On rotten wood, more rarely on herbaceous stems.

TYPE LOCALITY: Cuba.



1. PATELLA ALBIDA
2. PATELLA SCUTELLATA

DISTRIBUTION: Cuba to Jamaica and Texas; also in Trinidad and South America.

ILLUSTRATIONS: Cooke, Mycographia *pl. 37, f. 144, 145;* Jour. Linn. Soc. **31**: *pl. 18, f. 2.*

4. **Patella umbrorum** (Fries) Seaver, comb. nov.

Peziza umbrosa Fries, Syst. Myc. **2**: 85. 1822. Not *P. umbrosa* Schr.
Peziza umbrorum Fries, Syst. Myc. **2**: 612. 1823.
Humaria umbrorum Fuckel, Symb. Myc. 323. 1869.
Lachnea umbrorum Gill. Champ. Fr. Discom. 209. 1886.
Scutellinia umbrarum Kuntze, Rev. Gen. Pl. **2**: 869. 1891.
?Sepultaria rubropurpurea Clements, Bot. Surv. Nebr. **4**: 15. 1896.
?Lachnea rubropurpurea Sacc. & Sydow in Sacc. Syll. Fung. **14**: 755. 1899.
?Scutellinia heterospora Clements, Bull. Torrey Club **30**: 90. 1903.
?Scutellinia dispora Clements, Bull. Torrey Club **30**: 90. 1903.
?Lachnea heterospora Sacc. & D. Sacc. in Sacc. Syll. Fung. **18**: 36. 1906.
?Lachnea dispora Sacc. & D. Sacc. in Sacc. Syll. Fung. **18**: 37. 1906.
Ciliaria umbrorum Boud. Hist. Class. Discom. Eu. 61. 1907.

Apothecia gregarious or densely crowded, at first closed and globose, at maturity scutellate, reaching a diameter of 5 mm. to 1 cm., externally appearing brown from the dark-colored hairs; hymenium slightly concave, almost plane or occasionally slightly convex, bright-red, almost scarlet; excipulum composed of large roundish or slightly angular cells which form a palisade near the margin, the loose ends of which project, giving rise to a fringe-like border of club-shaped hyphae; hairs ranging from a few pale cells when young to long rigid setae at maturity, dark reddish-brown, reaching a length of 1 mm. and a diameter of 35 μ at the broadest point, closely septate, the walls 3–4 μ thick, constricted below near the point of attachment; asci cylindric above, gradually tapering below into a stem-like base, reaching a length of 325–350 μ and a diameter of 26–27 μ, 8-spored but part of the spores often remaining undeveloped; spores 1-seriate, usually obliquely arranged in the ascus, at first smooth and usually containing one large oil-drop, 12–14 $\times$ 23–24 μ, hyaline, at maturity becoming sculptured; spore-sculpturing consisting of rather coarse warts; warts about 2 μ in diameter; paraphyses rather stout, about 3–4 μ in diameter below, strongly and rather abruptly enlarged above, reaching a diameter of 10 μ at their apices, densely filled with minute orange granules.

On very damp soil, more rarely on rotten wood.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to Colorado, New Mexico and West Virginia; also in Europe.

ILLUSTRATIONS: Cooke, Mycographia *pl. 35, f. 138;* Boud. Ic. Myc. *369;* Gill. Champ. Fr. Discom. *pl. 68.*

EXSICCATI: N. Am. Fungi *2321, 2911.*

5. **Patella Lusatiae** (Cooke) Seaver, comb. nov.

Peziza Lusatiae Cooke, Mycographia 80. 1875.
?Peziza badioberbis Berk.; Cooke, Grevillea **8**: 61. 1879.
?Lachnea badioberbis Sacc. Syll. Fung. **8**: 173. 1889.
Lachnea Lusatiae Sacc. Syll. Fung. **8**: 178. 1889.
Scutellinia Lusatiae Kuntze, Rev. Gen. Pl. **2**: 869. 1891.
Scutellinia badeoberbis Kuntze, Rev. Gen. Pl. **2**: 869. 1891.
Ciliaria Lusatiae Boud. Hist. Class. Discom. Eu. 62. 1907.

Apothecia gregarious, at first globose, gradually expanding and at maturity scutellate, reaching a diameter of 5 mm., externally densely clothed with hairs; hymenium plane or slightly concave, bright-red; hairs very long and forming a conspicuous fringe about the border of the apothecium, dark-colored, septate, swollen near the base and often forked below, gradually tapering above to a bristle-like apex, often reaching a length of more than 1 mm.; asci cylindric above, reaching a diameter of 15–18 μ; spores 1-seriate, broad-ellipsoid or slightly narrowed toward the ends, containing one or two oil-drops, at first smooth, becoming sculptured, hyaline or faintly yellowish, 14–16 $\times$ 22–26 μ; spore-sculpturing consisting of rather prominent warts; paraphyses strongly enlarged above, reaching a diameter of 8–9 μ at their apices.

On rotten wood.

TYPE LOCALITY: Germany.

DISTRIBUTION: New York; also in Europe and New Zealand.

ILLUSTRATION: Cooke, Mycographia *pl. 37, f. 146.*

6. **Patella miniata** (Clem.) Seaver, comb. nov.

Pelodiscus miniatus Clements, Bot. Surv. Nebr. **5**: 8. 1901.
Lachnea Clementsi Sacc. & Sydow in Sacc. Syll. Fung. **16**: 1147. 1902.

Apothecia occurring singly or in groups of two or three, reaching a diameter of 1–2 mm. and 0.5 mm. high, margin naked, externally except the base clothed with very short hairs; hymenium concave or nearly plane, circular in form, red; hairs short, continuous, flexuous, pale-yellow; spores 1-seriate, parallel with the ascus or obliquely arranged, ellipsoid, often only four

developed, the other four remaining abortive, 12–14 × 26–28 μ, hyaline, at maturity beautifully sculptured; spore-sculpturing consisting of papillae 2–3 μ high; paraphyses septate, enlarged above, reaching a diameter of 9–12 μ at their apices.

On stercoriated mud.

TYPE LOCALITY: Beal's slough, Otawaine woods, Nebraska.

DISTRIBUTION: Known only from the type locality.

7. **Patella albocincta** (Berk. & Curt.) Seaver, comb. nov. (PLATE 45, FIG. 11, 12.)

?Peziza Polytrichi Schum. Enum. Pl. Saell. 423. 1803.
Peziza albocincta Berk. & Curt. Grevillea **3**: 154. 1875.
Neottiella albocincta Sacc. Syll. Fung. **8**: 190. 1889.
Neottiella ovilla flavodisca Cooke & Massee, Grevillea **21**: 71. 1895.
Neottiella Polytrichi Massee, Brit. Fungus-Fl. **4**: 370. 1895.

Apothecia gregarious or scattered, at first closed and subglobose, gradually expanding, at maturity turbinate to subscutellate, sessile or short-stipitate, reaching a diameter of 10–12 mm. but often much smaller, externally clothed with hyaline (white to the naked eye) hairs which form a delicate fringe about the margin of the apothecium; marginal hairs often in dense fascicles, individual hairs 4–5 μ in diameter at the base, gradually tapering toward the apex, which is blunt and about 2 μ in diameter, reaching a length of 275 μ, hyaline; asci cylindric or subcylindric, reaching a length of 200 μ and a diameter of 14–16 μ; spores 1-seriate at maturity or with the ends slightly overlapping, ellipsoid, with the ends strongly narrowed, containing one large oil-drop, finally becoming sculptured, 10–12 × 24–28 μ; spore-sculpturing consisting of minute warts; paraphyses slightly enlarged above, reaching a diameter of 4–5 μ at their apices.

On the ground in woods, often among mosses.

TYPE LOCALITY: South Carolina.

DISTRIBUTION: New York to South Carolina; also in Europe.

ILLUSTRATIONS: Massee, Brit. Fungus-Fl. **4**: 371, *f. 1–4;* Grevillea **22**: 109, *f. 1–5;* Cooke, Mycographia *pl. 40, f. 157.*

8. **Patella ovilla** (Peck) Seaver, comb. nov.

Peziza ovilla Peck, Ann. Rep. N. Y. State Mus. **28**: 66. 1879.
Neottiella ovilla Sacc. Syll. Fung. **8**: 194. 1889.
Humaria rhodoleuca Bres. Fungi Trid. **2**: 79. 1898.
Neottiopezis macrospora Clements, Bull. Torrey Club **30**: 89. 1903.
Neottiella macrospora Sacc. & D. Sacc. in Sacc. Syll. Fung. **18**: 39. 1906.

Apothecia subglobose or hemispheric, slightly attenuated below, the margin crenate, reaching a diameter of 5 mm. to 1 cm., externally whitish and clothed with tomentose hairs; the excipular cells reaching a diameter of 20 μ; hairs blunt, rather thin-walled, septate and slightly constricted at the septa, reaching a diameter of 15 μ; asci cylindric above, tapering below into a stem-like base, reaching a length of 280–350 μ and a diameter of 14–16 μ, 8-spored; spores fusoid, containing one to three oil-drops, 10–14 $\times$ 32–40 μ, smooth or very minutely sculptured; spore-sculpturing consisting of minute warts; paraphyses branching, septate, 3.5–4 μ in diameter, enlarged above, where they reach a diameter of 4–7 μ.

On the ground in woods.

TYPE LOCALITY: Sageville, New York.

DISTRIBUTION: New York to Minnesota and Nebraska; also in Europe.

ILLUSTRATION: Bres. Fungi Trid. **2**: *pl. 193, f. 2.*

9. **Patella Hetieri** (Boud.) Seaver, comb. nov.

Neottiella Hetieri Boud. Bull. Soc. Myc. Fr. **12**: 12. 1896.

Apothecia gregarious or scattered, at first subglobose, opening and gradually expanding, finally becoming scutellate or subdiscoid, reaching a diameter of 2 mm., externally pale-orange and clothed with delicate white hairs; hymenium slightly concave or nearly plane, bright orange-red; hairs fairly abundant but not conspicuous to the unaided eye, rigid, blunt when young, sharp-pointed when mature, hyaline, septate, reaching a length of 150–200 μ and a diameter of 4–6 μ; asci cylindric above, gradually tapering below into a short stem-like base, reaching a length of 200–225 μ and a diameter of 16–20 μ, 8-spored; spores 1-seriate, ellipsoid or more often ovoid, containing one large oil-drop or very rarely two, hyaline, about 12 $\times$ 18 μ; paraphyses stout, rather freely branched, septate, often swollen near the septa, filled with orange granules, reaching a diameter of 8 μ at their apices.

On old burnt place which has been overrun with mosses, or occasionally where there is no trace of fire.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York and Connecticut; also in Europe.

ILLUSTRATION: Bull. Soc. Myc. Fr. **12**: *pl. 3, f. 2.*

10. **Patella ricciophila** Seaver, sp. nov. (PLATE 45, FIG. 15, 16.)

Apothecia gregarious or scattered, at first globose, becoming expanded and subdiscoid, reaching a diameter of 1.5 mm., pale-orange, surrounded by a pale-orange or whitish border, and rather densely clothed with white hairs; hymenium plane, slightly concave, or convex, a little brighter in color than the outside of the apothecium, becoming roughened or pitted; hairs perfectly hyaline, thick-walled, tapering into an acute apex, reaching a length of 150–200 μ and a diameter of 16–18 μ; asci cylindric or subcylindric, with a short stem-like base, reaching a length of 180–200 μ and a diameter of 20 μ, 8-spored; spores 1-seriate or rarely crowded, decidedly ovoid, the small end directed either upward or downward, usually containing one large oil-drop, hyaline, smooth, about 18 $\times$ 24–26 μ; paraphyses stout, slightly enlarged above, reaching a diameter of 5–6 μ, filled with orange granules.

Closely associated with and apparently growing on the living thallus of *Riccia nigrella* brought from Texas and grown in the propagating house of The New York Botanical Garden, December 6, 1915.

DISTRIBUTION: Known only from the type collection.

11. **Patella punicea** (Clem.) Seaver, comb. nov.

Sepultaria punicea Clements, Bot. Surv. Nebr. **4**: 14. 1896.
Lachnea punicea Sacc. & Sydow in Sacc. Syll. Fung. **14**: 759. 1899.

Apothecia subgregarious, superficial or nearly so, at first subglobose, expanding and becoming scutellate to discoid, reaching a diameter of 2 mm., externally densely clothed with hairs; hymenium bright-red, concave; hairs brown, septate, sharp-pointed, reaching a length of 300–400 μ and a diameter of 16–20 μ; asci cylindric or subcylindric, reaching a length of 225–230 μ and a diameter of 10–14 μ; spores 1-seriate, ellipsoid, smooth, hyaline, 7–9 $\times$ 10–12 μ; paraphyses filiform, clavate, densely filled with orange granules.

On wet, decaying wood.

TYPE LOCALITY: Bluffs of the Missouri River, Otoe County, Nebraska.

DISTRIBUTION: Known only from the type locality.

12. **Patella gilva** (Boud.) Seaver, comb. nov.

Peziza gilva Boud.; Cooke, Mycographia 240. 1879.
Tricharia gilva Boud. Hist. Class. Discom. Eu. 57. 1907.
Lachnea gilva Sacc. Syll. Fung. **8**: 184. 1889.
Scutellinia gilva Kuntze, Rev. Gen. Pl. **2**: 869. 1891.

Apothecia scattered or occasionally closely crowded, at first globose or subglobose, gradually expanding and becoming shallow cup-shaped, reaching a diameter of 5–6 mm., externally ochraceous-brown and clothed with rather inconspicuous hairs; hymenium concave, ochraceous or brownish; hairs blunt when young, sharp-pointed when mature, septate, pale-yellowish, reaching a length of 200 μ and a diameter of 8 μ; asci cylindric above, gradually tapering below into a stem-like base; spores 1-seriate, ellipsoid, hyaline, containing no oil-drops, about 8–10 $\times$ 14–18 μ, smooth; paraphyses rather stout, gradually enlarged above, reaching a diameter of 4–5 μ at their apices.

On an old burnt place which had been overrun with mosses.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York; also in Europe.

ILLUSTRATIONS: Cooke, Mycographia *pl. 113, f. 406;* Boud. Ic. Myc. *pl. 347.*

13. **Patella setosa** (Nees) Seaver; Brenckle, Fungi Dakot. *458.* 1920.

Peziza setosa Nees, Syst. Pilze 260. 1817.
Humaria setosa Fuckel, Symb. Myc. 321. 1869.
Lachnea setosa Gill. Champ. Fr. Discom. 75. 1880.
Scutellinia setosa Kuntze, Rev. Gen. Pl. **2**: 869. 1891.
Ciliaria setosa Boud. Hist. Class. Discom. Eu. 62. 1907.
?Lachnea subcrinita Rehm, Ann. Myc. **7**: 535. 1909.

Apothecia thickly gregarious but seldom crowded, at first globose or subglobose and bristling with a dense covering of dark-colored setae, expanding and becoming discoid or subdiscoid, reaching a diameter of 2–3 mm.; hymenium plane or nearly so, dull-red; hairs reaching a length of 500 μ and a diameter of 25 μ at the broadest point near the base, forked and rooting at the extreme base, thick-walled, dark-brown, subacute at their apices; asci cylindric or subcylindric, reaching a length of 300–325 μ and a diameter of 12–15 μ; spores 1-seriate, parallel with the ascus or slightly oblique, ellipsoid, smooth, filled with numerous oil-drops which become more or less broken up in dried plants, 11–13 $\times$ 20–22 μ; paraphyses enlarged at their apices, where they reach a diameter of 4–5 μ.

On dead and often partially rotted wood of deciduous trees.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to North Dakota, Colorado and Nicaragua; also in Europe.

ILLUSTRATIONS: Boud. Ic. Myc. *pl. 370;* Cooke, Mycographia *pl. 34, f. 133;* Nees, Syst. Pilze *pl. 37, f. 275;* Gill. Champ. Fr. Discom. (suites); Proc. Ohio Acad. Sci. **5**: *pl. 1, f. 1–7.*

EXSICCATI: N. Am. Fungi *2033, 2910* (as *Peziza vitellina* Pers.); Seaver, North Dakota Fungi *27;* Brenckle, Fungi Dakot. *458.*

14. **Patella Sequoiae** (Phill.) Seaver, comb. nov.

Peziza Sequoiae Phill. Grevillea **7**: 22. 1878.
Lachnea Sequoiae Sacc. Syll. Fung. **8**: 176. 1889.
Scutellinia Sequoiae Kuntze, Rev. Gen. Pl. **2**: 869. 1891.

Apothecia gregarious, sessile, reaching a diameter of 3–5 mm., externally clothed with hairs; hymenium concave, reddish or flesh-colored; hairs short near the base, the marginal ones reaching a length of 600 μ long, straight and subacute, sparingly septate; asci cylindric or subcylindric; spores 1-seriate, ellipsoid, smooth, hyaline, 12–13 $\times$ 17–20 μ; paraphyses filiform, slightly enlarged above.

On dead twigs of *Sequoia gigantea.*

TYPE LOCALITY: California.

DISTRIBUTION: Known only from the type locality.

ILLUSTRATION: Cooke, Mycographia *pl. 113, f. 403.*

15. **Patella melaloma** (Alb. & Schw.) Seaver, comb. nov. (PLATE 15, FIG. 2.)

Humaria melaloma Massee, Brit. Fungus-Fl. **4**: 411. 1895.
Peziza melaloma Alb. & Schw. Consp. Fung. 336. 1805.
Peziza intermixta Karst. Not. Fauna Fl. Fenn. **10**: 119. 1869.
Pyronema melalomum Fuckel, Symb. Myc. 319. 1869.
Peziza adusta Cooke & Peck; Peck, Ann. Rep. N. Y. State Mus. **27**: 107. 1875.
Aleuria melaloma Gill. Champ. Fr. Discom. 54. 1879.
Humaria melalomoides Rehm in Sydow, Mycoth. March. *778.* 1884.
Lachnea melaloma Sacc. Syll. Fung. **8**: 181. 1889.
?Humaria macrocystis Sacc. Syll. Fung. **8**: 122. 1889.
?Pyronema subhirsutum Sacc. Syll. Fung. **8**: 108. 1889.
Scutellinia melaloma Kuntze, Rev. Gen. Pl. **2**: 869. 1891.
Humariella melaloma Schröt. Krypt.-Fl. 3^2: 37. 1893.
Lachnea intermixta Rehm in Rab. Krypt.-Fl. 1^3: 1047. 1896.
Humaria adusta Sacc. Syll. Fung. **8**: 141. 1889.

Humaria flavoaurantiaca Rehm, Ann. Myc. **2**: 35. 1904.
Anthracobia melaloma Boud. Hist. Class. Discom. Eu. 65. 1907.

Apothecia gregarious or more often densely crowded, forming continuous masses often several cm. in diameter, at maturity scutellate with the margin slightly elevated, externally dark reddish-brown and clothed with hairs which are closely adpressed and not extending above the margin of the apothecium, reaching a diameter of 3–5 mm. or rarely 1 cm.; hymenium slightly concave, plane or convex, very variable in color, ranging from pale- to bright-orange or dark-brown; hairs flexuous, brown, sparingly septate, the ends blunt, of variable length; excipular cells large, rounded or angular, with brown cell walls, reaching a diameter of 35 μ; asci cylindric or subcylindric, reaching a length of 150–200 μ and a diameter of 12–14 μ, rather abruptly narrowed below; spores 1-seriate, parallel with the ascus or slightly oblique, or occasionally irregularly disposed, containing one or more, commonly two oil-drops, ellipsoid, smooth, hyaline, 7–10 $\times$ 16–21 μ; paraphyses enlarged upwards, reaching a diameter of 4–6 μ at their apices, reddish.

On burnt ground and charcoal beds.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to Iowa, Texas, California and Jamaica; also in Europe.

ILLUSTRATIONS: Alb. & Schw. Consp. Fung. *pl. 2, f. 5;* Boud. Ic. Myc. *pl. 307;* Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 10, f. 2;* Gill. Champ. Fr. Discom. *pl. 55, f. 2;* Pat. Tab. Fung. *f. 275;* Cooke, Mycographia *pl. 17, f. 67;* Rab. Krypt.-Fl. 1^3: 1032, *f. 1–5.*

16. **Patella crucipila** (Cooke & Phill.) Seaver, comb. nov.

Peziza crucipila Cooke & Phill.; Cooke, Mycographia 136. 1876.
Lachnea crucipila Phill. Brit. Discom. 229. 1887.
Neottiella crucipila Sacc. Syll. Fung. **8**: 192. 1889.

Apothecia gregarious or crowded, at first subglobose, expanding and becoming scutellate to discoid, the margin of the apothecium even or irregularly wavy, externally pale-yellow and clothed with a mass of rather pale hairs, attached to the substratum by a mass of coarse, white mycelium, reaching a diameter of about 5 mm.; hymenium concave or plane, bright-orange, brighter than the outside of the apothecium (the type described as pale-scarlet), becoming paler in dried specimens; excipular cells large and irregularly polygonal, reaching a diameter of

50–75 μ, the walls hyaline or very faintly yellowish; hairs 2–4-pronged or more rarely with five or six rays springing from nearly the same point, the basal cell often enlarged and irregularly rectangular, the rays reaching a diameter of 20 μ at their bases and a length of 500–600 μ, sharp-pointed, pale-brown; asci cylindric or subcylindric, reaching a length of 200–225 μ and a diameter of 12–14 μ; spores 1-seriate, ellipsoid, smooth, thin-walled and easily collapsing, 9 × 15–20 μ; paraphyses clavate, filled with orange granules.

On rich sandy soil in coniferous woods, occasionally on bits of wood.

TYPE LOCALITY: Great Britain.

DISTRIBUTION: Washington; also in Europe.

ILLUSTRATIONS: Cooke, Mycographia *pl. 60, f. 237;* Gill. Champ. Fr. Discom. *pl. 71;* Pat. Tab. Fung. *f. 481.*

17. **Patella stercorea** (Pers.) Weber; Wiggers, Fl. Hols. 106. 1780.

?Elvela lutea Scop. Fl. Carn. **2**: 481. 1772.
Peziza stercorea Pers. Obs. Myc. **2**: 89. 1799.
Peziza ciliata Bull. Champ. Fr. **1**: 257. 1809.
Humaria stercorea Fuckel, Symb. Myc. 321. 1869.
?Humaria alpina Fuckel, Symb. Myc. Nacht. **3**: 32. 1875.
Lachnea stercorea Gill. Champ. Fr. Discom. 76. 1880.
Lasiobolus stercoreus Karst. Act. Fauna Fl. Fenn. **2**: 122. 1885.
?Lachnea alpina Sacc. Syll. Fung. **8**: 180. 1889.
?Scutellinia alpina Kuntze, Rev. Gen. Pl. **2**: 869. 1891.
Scutellinia stercorea Kuntze, Rev. Gen. Pl. **2**: 869. 1891.
Humariella stercorea Schröt. Krypt.-Fl. Schles. **3²**: 37. 1893.
Patella lutea Morgan, Jour. Myc. **8**: 188. 1902.
Cheilymenia stercorea Boud. Hist. Class. Discom. Eu. 63. 1907.

Apothecia gregarious or crowded, expanding and becoming scutellate to discoid, externally pale-yellowish, clothed with hairs of two kinds, simple and compound, reaching a diameter of 2 mm.; hymenium plane or concave, bright-orange; the simple hairs confined mainly to the margin of the apothecium, giving rise to a conspicuous fringe-like border, reaching a length of 1 mm. and a diameter of 25–50 μ, 2–5-septate, thick-walled, acute at the apex and with an enlarged and often forked base, dark-brown; stellate hairs usually not extending above the margin of the apothecium, consisting of a triangular or rectangular basal portion with 2–5 rays which are straight, acute at their apices, and reach a maximum length of 200 μ, the rays

similar in color and appearance to the simple hairs but smaller; asci cylindric or subcylindric, reaching a length of 225–300 μ and a diameter of 15–20 μ; spores 1-seriate, parallel with the ascus or oblique, ellipsoid, smooth, pale-yellow, 9–14 × 18–24 μ; paraphyses clavate, reaching a diameter of 7–8 μ at their apices.

On cow dung.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to California; also in Europe.

ILLUSTRATIONS: Boud. Ic. Myc. *pl. 384;* Cooke, Mycographia *pl. 38, f. 147, 148;* Gill. Champ. Fr. Discom. *pl. 63, f. 1;* Bull. Herb. Fr. *pl. 438, f. 2;* Sow. Engl. Fungi *pl. 352.*

18. **Patella theleboloides** (Alb. & Schw.) Seaver, comb. nov.

?Peziza fimetaria Schum. Pl. Saell. **2**: 426. 1803.
Peziza theleboloides Alb. & Schw. Consp. Fung. 321. 1805.
?Pyronema subhirsutum Fuckel, Symb. Myc. 320. 1869. Excl. syn.
Peziza dalmeniensis Cooke, Grevillea **3**: 66. 1874.
Aleuria subhirsuta Gill. Champ. Fr. Discom. 49. 1879.
Lachnea theleboloides Gill. Champ. Fr. Discom. 74. 1879.
Humaria theleboloides Quél. Ench. Fung. 285. 1886.
Lachnea dalmeniensis Sacc. Syll. Fung. **8**: 180. 1889.
Scutellinia theleboloides Kuntze, Rev. Gen. Pl. **2**: 869. 1891.
Scutellinia dalmeniensis Kuntze, Rev. Gen. Pl. **2**: 869. 1891.
Lachnea Lojkaena Rehm in Rab. Krypt.-Fl. **1**^{3}: 1045. 1896.
Lachnea diplotricha Rehm, Ann. Myc. **2**: 34. 1904.
Cheilymenia theleboloides Boud. Hist. Class. Discom. Eu. 62. 1907.
Cheilymenia subhirsuta Boud. Hist. Class. Discom. Eu. 62. 1907.
Cheilymenia fimetaria Boud. Hist. Class. Discom. Eu. 63. 1907.
Cheilymenia dalmeniensis Boud. Hist. Class. Discom. Eu. 63. 1907.

Apothecia gregarious or closely crowded, forming congested masses, at first subglobose or short-cylindric, expanding and becoming scutellate to discoid, reaching a diameter of 3–5 mm. or rarely as large as 1 cm., externally pale-yellow, sparingly clothed with hairs; hymenium concave or plane, bright-yellow; the excipular cells rounded or angular, reaching a diameter of 35 μ, the cell walls pale-yellow; hairs rigid, hyaline or very pale-yellow, enlarged at the base, tapering into a bristle-like apex, 2–4-septate, reaching a length of 200–500 μ and a diameter of 10 μ at their bases, very rarely branched; asci cylindric above, abruptly tapering at the base, reaching a length of 225–275 μ and a diameter of 12–14 μ; spores 1-seriate, parallel with the ascus or oblique, ellipsoid, smooth, hyaline or faintly yellowish, thin-walled, often collapsing on one side, 7–10 × 14–20 μ; pa-

raphyses slightly enlarged at their apices, reaching a diameter of 3–4 μ, filled with yellow granules.

On cow dung or occasionally on rich soil, rubbish, paper, cloth, etc.

TYPE LOCALITY: Europe.

DISTRIBUTION: Connecticut to Delaware, North Dakota, California and Bermuda; also in Europe.

ILLUSTRATIONS: Alb. & Schw. Consp. Fung. *pl. 12, f. 4;* Boud. Ic. Myc. *pl. 380, 381;* Gill. Champ. Fr. Discom. *pl. 50, f. 1;* Cooke, Mycographia *pl. 39, f. 151, 153;* Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 8, f. 2.*

EXSICCATI: N. Am. Fungi *2035;* Brenckle, Fungi Dakotenses *141* (as *Humaria subhirsuta* (Schum.) Karst.).

19. **Patella coprinaria** (Cooke) Seaver, comb. nov. (PLATE 45, FIG. 18.)

Peziza coprinaria Cooke, Grevillea **4**: 91. 1875.
Peziza scubalonta Cooke & Gerard; Cooke, Grevillea **4**: 92. 1875.
Lachnea coprinaria Phill. Brit. Discom. 224. 1887.
Lachnea scubalonta Sacc. Syll. Fung. **8**: 179. 1889.
Scutellinia coprinaria Kuntze, Rev. Gen. Pl. **2**: 869. 1891.
Sepultaria pediseta Clements, Bot. Surv. Nebr. **4**: 13. 1896.
Scutellinia scubalonta Kuntze, Rev. Gen. Pl. **2**: 869. 1891.
Lachnea pediseta Sacc. & Sydow in Sacc. Syll. Fung. **14**: 758. 1899.
Cheilymenia coprinaria Boud. Hist. Class. Discom. Eu. 63. 1907.

Apothecia gregarious or scattered, at first closed, expanding and becoming scutellate to discoid, the margin slightly wavy, pale-yellow, sparingly clothed with hairs, reaching a diameter of 3 mm. to 1 cm.; hymenium plane or concave, bright-yellow to pale-orange; the excipular cells very large, reaching a diameter of 75–80 μ, irregularly polygonal, thin-walled, the walls subhyaline; hairs conspicuous, those about the margin projecting into a sparse fringe-like border, pale-brown, sparingly septate, the base often once or several times divided, giving rise to root-like branches, tapering above into a bristle-like apex, reaching a length of 600 μ and a diameter of 20–30 μ at the base; asci cylindric or subcylindric, reaching a length of 200 μ and a diameter of 12–14 μ; spores 1-seriate, ellipsoid, smooth, hyaline, 8–10 × 17–20 μ; paraphyses clavate, reaching a diameter of 5–7 μ at their apices.

On dung of various kinds.

TYPE LOCALITY: Rannock, Scotland.

DISTRIBUTION: New York to Nebraska and California; also in Europe.

ILLUSTRATIONS: Cooke, Mycographia *pl. 38, f. 149, 150;* Boud. Ic. Myc. *pl. 383.*

20. **Patella maculosa** (Phill.) Seaver, comb. nov.

Peziza maculosa Phill. Grevillea **5**: 116. 1877.
Lachnea maculosa Sacc. Syll. Fung. **8**: 171. 1889.

Apothecia sessile, superficial or semi-immersed, fleshy, brown, reaching a diameter of 1 cm., externally marked with brown spots and clothed with flexuous hairs; hymenium slightly concave, brownish; hairs slender, septate, blunt, pale-brown, minutely sculptured; asci cylindric, 8-spored; spores ellipsoid, smooth, hyaline, 7 × 14 μ; paraphyses filiform, septate.

On cow dung.

TYPE LOCALITY: Blue Canyon, Sierra Nevada Mountains, California.

DISTRIBUTION: Known only from the type locality.

ILLUSTRATIONS: Grevillea **5**: *pl. 89, f. 11;* Cooke, Mycographia *pl. 99, f. 358.*

21. **Patella pulcherrima** (Crouan) Seaver, comb. nov.

Ascobolus pulcherrimus Crouan, Ann. Sci. Nat. IV. **10**: 196. 1858.
Peziza pulcherrima Cooke, Mycographia 84. 1876.
Humaria pulcherrima Speg. Michelia **1**: 237. 1878.
Lachnea pulcherrima Gill. Champ. Fr. Discom. 76. 1880.
Cheilymenia pulcherrima Boud. Hist. Class. Discom. Eu. 63. 1907.
Scutellinia pulcherrima Kuntze, Rev. Gen. Pl. **2**: 869. 1891.
Lasiobolus pulcherrimus Schröt. Krypt.-Fl. Schles. **3²**: 54. 1893.

Apothecia gregarious or congested, often forming dense masses several cm. in diameter, short-cylindric, scarcely exceeding 1 mm. in diameter and reaching a height of 1 mm., externally pale-yellow and sparingly clothed with hairs; hymenium plane or concave, bright-yellow; hairs small, rigid, enlarged at their bases, acute at their apices, 2–4-septate, pale-brown, reaching a length of 100–200 μ and a diameter of 15 μ at their bases; asci cylindric or subcylindric, reaching a length of 200–235 μ and a diameter of 20 μ; spores 1-seriate, ellipsoid, hyaline or very slightly yellowish, smooth, thin-walled, often collapsing on one side, 10–12 × 20–22 μ; paraphyses filiform, slightly enlarged above, yellow.

On cow dung.

TYPE LOCALITY: France.

DISTRIBUTION: Bermuda Islands and reported from Nebraska; also in Europe.

ILLUSTRATIONS: Ann. Sci. Nat. IV. **10**: *pl. 13, f. 32–37;* Cooke, Mycographia *pl. 39, f. 154;* Gill. Champ. Fr. Discom. *pl. 63, f. 2.*

22. **Patella raripila** (Phill.) Seaver, comb. nov.

Ascobolus raripilus Phill. Grevillea **7**: 23. 1878.
Lasiobolus raripilus Sacc. Syll. Fung. **8**: 537. 1889.

Apothecia gregarious or closely crowded, often forming congested masses of considerable extent, reaching a diameter of 1–2 mm., pale-yellow, externally scantily clothed with hairs; hymenium at first concave, becoming plane or a little convex, bright-yellow, a little brighter than the outside of the apothecium; hairs pale-yellow or subhyaline, rigid, tapering into a bristle-like apex; asci broad-clavate; spores 1-seriate or irregularly disposed, ellipsoid, hyaline, 10–14 × 25–30 μ; paraphyses slender, slightly enlarged above.

On the dung of animals and on the surrounding soil.

TYPE LOCALITY: California.

DISTRIBUTION: California and Iowa.

ILLUSTRATION: Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 32, f. 2.*

23. **Patella fimetaria** Seaver, sp. nov.

Apothecia gregarious or scattered, at first globose, expanding and becoming discoid, reaching a diameter of 5 mm., externally densely clothed with hairs; hymenium plane or slightly concave, yellowish to orange; hairs rigid, reaching a length of 1 mm. and a diameter of 30 μ, thick-walled, septate, attenuated above into a very sharp-pointed apex, the base forked and root-like, dark-brown; asci cylindric or subcylindric, reaching a length of 200–225 μ and a diameter of 15 μ; spores 1-seriate, ellipsoid, hyaline, containing two large oil-drops, becoming minutely sculptured, about 12 × 20 μ; spore-sculpturing consisting of minute papillae or warts; paraphyses strongly enlarged above, reaching a diameter of 7–8 μ at their apices, nearly hyaline.

On cow dung.

Type collected east of Tripoli along Halifax Brook, Wash-

ington County, New York, by Stewart H. Burnham (Herb. The New York Botanical Garden).

DISTRIBUTION: Known only from the type locality.

24. **Patella paludosa** (Boud.) Seaver, comb. nov.

Ciliaria paludosa Boud. Bull. Soc. Myc. Fr. **10**: 65. 1894.
Lachnea paludosa Sacc. Syll. Fung. **11**: 400. 1895.
Lachnea gregaria pseudogregaria Rick in Oester. Bot. Zeitschr. **48**: 62. 1898.
Lachnea pseudogregaria Rehm, Ascom. *1268;* Hedwigia **38**: (243). 1899.
Trichophaea pseudogregaria Boud. Hist. Class. Discom. Eu. 60. 1907.
Trichophaea paludosa Boud. Hist. Class. Discom. Eu. 60. 1907.

Apothecia closely gregarious, at first subglobose, finally becoming scutellate to discoid, reaching a diameter of 1–2 mm., externally clothed with dark-colored hairs which project about the margin of the apothecium giving rise to a fringe-like border; hymenium concave or nearly plane, whitish or bluish-white; hairs rigid, sharp-pointed, sparingly septate, but the septa often indistinct, so that the hairs often appear to be non-septate, reddish-brown, reaching a length of 500 μ and a diameter of 20 μ at the broadest point; asci cylindric or subcylindric above, gradually tapering below, reaching a length of 300 μ and a diameter of 15–17 μ; spores 1-seriate, ellipsoid, containing two oil-drops which become more or less obscured at maturity, becoming sculptured, reaching a length of 22–25 μ and a diameter of 15–17 μ; spore-sculpturing taking the form of tubercles 2–3 μ in diameter, giving to the spore a scalloped appearance; paraphyses slender, slightly enlarged above, hyaline.

On soil in swampy places.

TYPE LOCALITY: Montmorency, France.

DISTRIBUTION: New York to Iowa; also in Europe.

ILLUSTRATIONS: Bull. Soc. Myc. Fr. **10**: *pl. 2, f. 4;* Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 11, f. 1.*

25. **Patella tuberculata** Seaver, sp. nov.

Apothecia gregarious but not usually crowded, at first globose and closed, the light-colored exterior concealed by the dense covering of dark-brown hairs, expanding and becoming scutellate to discoid, reaching a diameter of 1–2 mm.; hymenium white or whitish, plane or very slightly concave; hairs reddish-brown, sparingly septate, straight or nearly so, thick-walled, tapering into a bristle-like apex, varying in length according to age, but reaching an extreme of 500 μ and a diameter of 20 μ; asci cylindric

or subcylindric above, tapering into a stem-like base, reaching a length of 300–500 μ and a diameter of 25–27 μ; spores 1-seriate, at first smooth, containing two oil-drops, soon becoming sculptured, reaching a length of 30–35 μ and a diameter of 20–22 μ, hyaline; spore-sculpturing taking the form of numerous tubercles which reach a diameter of 3 μ or rarely as large as 4–5 μ, the tubercles rounded and giving the spore a scalloped appearance; paraphyses slender below, enlarged above, reaching a diameter of 7–9 μ at their apices.

On damp sandy soil among mosses.

TYPE LOCALITY: The New York Botanical Garden.

DISTRIBUTION: Known only from the type locality.

Distinguished from the preceding by the larger spores and coarser tubercles.

26. **Patella albida** (Schaeff.) Seaver; Brenckle, Fungi Dakot. *407*. 1916. (PLATE 14, FIG. 1.)

Elvela albida Schaeff. Fung. Bavar. **4**: 101. 1774.
Peziza hemisphaerica Weber; Wiggers, Fl. Hols. 105. 1780.
Peziza labellum Bull. Champ. Fr. *pl. 204*. 1784.
Octospora fasciculata Hedw. Descr. **2**: 14. 1788.
Peziza hispida Sow. Engl. Fungi *pl. 147*. 1798.
Humaria hemisphaerica Fuckel, Symb. Myc. 322. 1869.
Lachnea hemispherica Gill. Champ. Fr. Discom. 73. 1879.
Scutellinia hemisphaerica Kuntze, Rev. Gen. Pl. **2**: 869. 1891.
Sepultaria albida Morgan, Jour. Myc. **8**: 188. 1902.
?Scutellinia chaetoloma Clements, Bull. Torrey Club **30**: 89. 1903.
?Lachnea chaetoloma Sacc. & D. Sacc. in Sacc. Syll. Fung. **18**: 34. 1906.

Apothecia gregarious or scattered, at first globose and closed, gradually expanding and becoming concave-hemispheric, or occasionally shallow cup-shaped, reaching a diameter of 2–3 cm. and about one-half or one-third as deep, the margin wavy or regular, externally entirely clothed with tufted hairs which project and form a fringe about the margin of the apothecium; hymenium white or whitish, becoming yellowish or yellowish-brown in dried specimens; hairs enlarged at the base, tapering above into a bristle-like apex, reaching a diameter of 15–20 μ at the base and a length of 400–500 μ, dark-brown, septate, the walls about 3 μ thick; asci cylindric or subcylindric above, tapering rather abruptly below into a stem-like base, reaching a length of 325 μ and a diameter of 15–18 μ; spores 1-seriate, ellipsoid, 12–15 × 25–27 μ, hyaline or subhyaline, usually containing two large oil-drops which become more or less broken

up in dried specimens, becoming delicately sculptured; spore-sculpturing consisting of minute papillae often indistinct in dried specimens; paraphyses slender, rather strongly enlarged at their apices, reaching a diameter of 7–8 μ.

On soil and more rarely on rotten wood.

TYPE LOCALITY: Europe.

DISTRIBUTION: Maine to Winnipeg, Kansas and North Carolina; also in Europe.

ILLUSTRATIONS: Bull. Herb. Fr. *pl. 204;* Hedw. Descr. **2**: *pl. 4, f. B;* Gill. Champ. Fr. Discom. *pl. 61, f. 1;* Rab. Krypt.-Fl. 1^3: 1033, *f. 1–4;* Sow. Engl. Fungi *pl. 147;* E. & P. Nat. Pfl. 1^1: 181, *f. 147, C–D;* Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 9, f. 1;* Pat. Tab. Fung. *f. 82;* Cooke, Mycographia *pl. 30, f. 115;* Massee, Brit. Fungus-Fl. **4**: 290, *f. 5–9;* Schaeff. Fung. Bavar. **2**: *pl. 151.*

EXSICCATI: N. Am. Fungi *837;* Brenckle, Fungi Dakotenses *407;* Clements, Crypt. Form. Colo. *118.*

27. **Patella gregaria** (Rehm) Seaver, comb. nov. (PLATE 15, FIG. 1; 45, FIG. 13, 14.)

?Peziza proximella Karst. Not. Fauna Fl. Fenn. **10**: 125. 1869.
Humaria gregaria Rehm; Winter, Flora **55**: 508. 1872.
Lachnea gregaria Phill. Brit. Discom. 214. 1887.
Scutellinia gregaria Kuntze, Rev. Gen. Pl. **2**: 869. 1891.
Lachnea amphidoxa Rehm in Rab. Krypt.-Fl. 1^3: 1048. 1896.
Trichophaea gregaria Boud. Hist. Class. Discom. Eu. 60. 1907.

Apothecia gregarious or more densely crowded, at first globose, becoming concave-hemispheric and resembling in gross characters *P. albida* but much smaller, occasionally becoming expanded and scutellate, reaching a diameter of 4–5 mm.; hymenium concave, becoming nearly plane, white or whitish; the excipular cells rounded or angular, giving rise to hairs which are usually borne in clusters; hairs closely adpressed to the sides of the apothecium, the marginal ones projecting, giving rise to a delicate fringe-like border, septate, blunt at their apices when young, becoming sharp-pointed as they mature, reaching a length of 150–300 μ and a diameter of 10–12 μ, pale-brown; asci cylindric or subcylindric above, reaching a length of 300 μ and a diameter of 17 μ; spores 1-seriate, the ends overlapping, ellipsoid to fusoid, filled with oil-drops, becoming sculptured, reaching a length of 20–27 μ and a diameter of 10–13 μ; spore-



1. PATELLA GREGARIA
2. PATELLA MELALOMA

sculpturing taking the form of minute warts; paraphyses slender, slightly enlarged above, where they reach a diameter of 6 μ.

On rather sandy soil in woods and open places, often under pines.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to Delaware; also in Europe.

ILLUSTRATIONS: Cooke, Mycographia *pl. 32, f. 123;* Boud. Ic. Myc. *pl. 364;* Grevillea **3**: *pl. 32, f. 98.*

28. **Patella irregularis** (Clements) Seaver, comb. nov.

Scutellinia irregularis Clements, Bull. Torrey Club **30**: 90. 1903.
Lachnea irregularis Sacc. & D. Sacc. in Sacc. Syll. Fung. **18**: 36. 1906.

Apothecia gregarious, sessile, irregularly cup-shaped, finally subscutellate, reaching a diameter of 4–10 mm., externally whitish but appearing dark on account of the brown hairs with which they are clothed; hymenium concave, whitish; hairs septate, rigid, reaching a length of 325 μ and a diameter of 8 μ; asci cylindric or subcylindric, reaching a length of 200 μ and a diameter of 16 μ; spores 1-seriate, ellipsoid, containing two oil-drops, becoming rough, 8 × 15 μ; paraphyses clavate, slightly enlarged above, where they reach a diameter of 3 μ.

On decaying *Picea*.

TYPE LOCALITY: Jack Brook, Minnehaha, Colorado.

DISTRIBUTION: Known only from the type locality.

29. **Patella abundans** (Karst.) Seaver, comb. nov.

Peziza abundans Karst. Not. Fauna Fl. Fenn. **10**: 124. 1869.
?Peziza cretea Cooke, Trans. Bot. Soc. Edinburgh **13**: 46. 1877.
?Lachnea cretea Sacc. Syll. Fung. **8**: 184. 1889.
Lachnea abundans Sacc. Syll. Fung. **8**: 186. 1889.
?Scutellinia cretea Kuntze, Rev. Gen. Pl. **2**: 869. 1891.
Scutellinia abundans Kuntze, Rev. Gen. Pl. **2**: 869. 1891.
Trichophaea abundans Boud. Hist. Class. Discom. Eu. 61. 1907.

Apothecia gregarious or more often densely crowded, at first globose or subglobose, expanding and becoming discoid or subdiscoid, reaching a diameter of 1–3 mm., externally clothed with pale-brown hairs; hymenium concave or plane, whitish, becoming yellowish in dried specimens; hairs sparingly septate, usually subglobose at the base, tapering above into a bristle-like apex, reaching a length of 100–150 μ and a diameter of 7 μ, 1–3-septate when mature, subhyaline to pale-brown; asci cylindric or subcylindric above, abruptly tapering below, reaching a length of

150 μ and a diameter of 10 μ; spores 1-seriate, with the ends slightly overlapping, ellipsoid, the ends strongly narrowed, containing one or two conspicuous oil-drops, hyaline, 8–9 × 15–17 μ; paraphyses rather strongly enlarged above, reaching a diameter of 5–6 μ at their apices.

On ashes in burned places and on plaster walls.

TYPE LOCALITY: Finland.

DISTRIBUTION: New York to Iowa; also in Europe.

ILLUSTRATIONS: Trans. Bot. Soc. Edinburgh **13**: *pl. 3, f. k–o;* Cooke, Mycographia *pl. 100, f. 362;* Grevillea **6**: *pl. 97, f. 4–7;* Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 11, f. 3.*

30. **Patella Erinaceus** (Schw.) Morgan, Jour. Myc. **8**: 188. 1902.

Peziza Erinaceus Schw. Schr. Nat. Ges. Leipzig **1**: 119. 1822.
Lachnea Erinaceus Sacc. Syll. Fung. **8**: 182. 1889.
Scutellinia Erinaceus Kuntze, Rev. Gen. Pl. **2**: 869. 1891.
?Lachnea chrysotricha Rehm, Ann. Myc. **5**: 520. 1907.

Apothecia gregarious, at first subglobose, expanding and becoming scutellate to discoid, reaching a diameter of 2–5 mm., externally densely clothed with hairs; hymenium concave or nearly plane, whitish; hairs long, rigid, brown, reaching a length of 300–500 μ and a diameter of 14–18 μ, tapering rather abruptly into a bristle-like apex; asci cylindric or subcylindric above; spores 1-seriate, ellipsoid, smooth, hyaline, 10–11 × 18–20 μ; paraphyses slender, enlarged at their apices.

On decaying wood of deciduous trees.

TYPE LOCALITY: Pennsylvania.

DISTRIBUTION: New York to Ohio and ? Michigan.

ILLUSTRATION: Cooke, Mycographia *pl. 36, f. 140.*

31. **Patella albospadicea** (Grev.) Seaver, comb. nov.

Peziza albospadicea Grev. Fl. Edin. 420. 1824.
Peziza Woolhopeia Cooke & Phill. Grevillea **6**: 75. 1877.
Lachnea Woolhopeia Sacc. Syll. Fung. **8**: 185. 1889.
Scutellinia albospadicea Kuntze, Rev. Gen. Pl. **2**: 869. 1891.
Scutellinia Woolhopeia Kuntze, Rev. Gen. Pl. **2**: 869. 1891.
Sepultaria bryophila Clements, Bot. Surv. Nebr. **4**: 13. 1896.
Sepultaria pseudocrenulata Clements, Bot. Surv. Nebr. **4**: 14. 1896.
Lachnea pseudocrenulata Sacc. & D. Sacc. in Sacc. Syll. Fung. **14**: 755. 1899.
Lachnea bryophila Sacc. & D. Sacc. in Sacc. Syll. Fung. **14**: 757. 1899.
Trichophaea albospadicea Boud. Hist. Class. Discom. Eu. 61. 1907.

Apothecia gregarious, sessile, at first globose, becoming concave-hemispheric, finally almost entirely discoid, reaching a diameter of 6 mm., externally clothed with dense covering of hairs when young, the hairs appearing more scant at maturity; hymenium concave or plane, whitish or grayish; hairs often swollen at the base, tapering into a sharp-pointed apex, rather closely septate, pale-brown, reaching a length of 350 μ but often much shorter, and a diameter of 12 μ; asci cylindric or subcylindric, gradually tapering below into a stem-like base, reaching a length of 235 μ and a diameter of 10 μ; spores 1-seriate, ellipsoid, the ends slightly narrowed, containing one or two oil-drops, 9–12 $\times$ 14–20 μ, smooth, hyaline; paraphyses straight, very slightly enlarged at their apices, reaching a diameter of 3 μ.

On the ground in shady places, often among mosses.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to Nebraska; also in Europe.

ILLUSTRATIONS: Cooke, Mycographia *pl. 36, f. 141* and *pl. 113, f. 404;* Boud. Ic. Myc. *pl. 365;* Gill. Champ. Fr. Discom. (suites); Pat. Tab. Fung. *f. 171.*

32. **Patella pygmaea** (Clements) Seaver, comb. nov.

Sepultaria pygmaea Clements, Bot. Surv. Nebr. **4**: 14. 1896.
Lachnea pygmaea Sacc. & Sydow in Sacc. Syll. Fung. **14**: 755. 1899.

Apothecia gregarious, at first closed and subglobose, becoming scutellate to subdiscoid, reaching a diameter of 0.5–0.75 mm. or rarely as large as 1 mm., externally clothed with hairs; hymenium plane or concave, gray, becoming slightly brownish when old; hairs rigid, the ends blunt or becoming acute, septate, brown, reaching a length of 175–250 μ and a diameter of 12.5 μ; asci cylindric or subcylindric; spores ellipsoid, smooth, hyaline, 7 $\times$ 12 μ; paraphyses numerous, filiform.

On rich ground among filaments of *Lyngbya.*

TYPE LOCALITY: Otawaine Woods, Lancaster county, Nebraska.

DISTRIBUTION: Known only from the type locality.

DOUBTFUL AND EXCLUDED SPECIES

Lachnea vitellina (Pers.) Sacc. Syll. Fung. **8**: 179. 1889; *Peziza vitellina* Pers. Myc. Eu. **1**: 257. 1822. Specimens distributed under this name in N. Am. Fungi *2910* are *Patella setosa* (Nees) Seaver.

Lachnea crispata (Berk. & Curt.) Sacc. Syll. Fung. **8**: 169. 1889; *Peziza crispata* Berk. & Curt.; Berk. Jour. Linn. Soc. **10**: 367. 1868. No specimen of this species has been and its identity is uncertain.

Neottiella Spraguei (Berk. & Curt.) Sacc. Syll. Fung. **8**: 190. 1889; *Peziza Spraguei* Berk. & Curt.; Cooke, Bull. Buffalo Acad. Sci. **2**: 289. 1875. Examination of a specimen of this species in the herbarium of The New York Botanical Garden shows it to be a small specimen of *Peziza* but its exact identity is uncertain.

Lachnea livida (Schum.) Gill. Champ. Fr. Discom. 73. 1880; *Peziza livida* Schum. Enum. Pl. Saell. **2**: 422. 1803. American specimens referred to this name do not differ from *Patella Erinaceus* (Schw.) Seaver.

Neottiella albotecta (Berk. & Curt.) Sacc. Syll. Fung. **8**: 191. 1889; *Peziza albotecta* Berk. & Curt.; Berk. Jour. Linn. Soc. **10**: 367. 1868. No specimen of this species has been seen and its identity is uncertain.

Lachnea hirta (Schum.) Gill. Champ. Fr. Discom. 75. 1880; *Peziza hirta* Schum. Enum. Pl. Saell. **2**: 422. 1803. Most of the specimens examined under this name have been found to be either *Patella umbrorum* or *Patella scutellata*. In 1881, Rehm described *Humaria hirtella* as intermediate between *Peziza hirta* and *P. scutellata*. Later this species was made a synonym of *Lachnea hirta* (Schum.) Gill. We have no means of knowing just what Scumacher's species really was and since in current literature the name has been interpreted differently no species can be found to which the name can be applied with certainty.

Neottiella sericeovillosa Rehm; Kellerm. Jour. Myc. **14**: 6. 1908. No specimen of this species has been seen and its identity is in doubt. The species was described from material collected in Guatemala.

Lachnea fissilis (Sacc. & Cooke) Sacc. Syll. Fung. **8**: 169. 1889; *Peziza fissilis* Sacc. & Cooke, Michelia **2**: 257. 1881. The specie: was reported from Nebraska by Clements, placed in the subgenus *Sepultaria* and said to be closely related to *S. Geaster*, *S. tenuis* and *S. arenosa*.

Neottiella callichroa (Boud.) Sacc. Syll. Fung. **8**: 190. 1889; *Humaria callichroa* Boud. Bull. Soc. Bot. Fr. **28**: 93. 1881. Reported from Nebraska by Clements. No specimen has been seen.

Neottiopezis macrospora Clements, Bull. Torrey Club **30**: 89. 1903 Apothecia described as 3–6 mm. in diameter and spores 16–18 × 40–42 μ. No specimen has been seen.

27. **WYNNEA** Berk. & Curt. Jour. Linn. Soc. **9**: 424. 1867.

Apothecia usually cespitose and branched at the base, springing from a sclerotium buried in the substratum, thick, firm, subcartilaginous, tough and subcoriaceous in drying, erect, elongate ear-shaped or spoon-shaped, simple or proliferous; asci cylindric or subcylindric, 8-spored; spores ellipsoid to fusoid, unequal-sided; paraphyses stout, simple or branched.

Type species, *Wynnea gigantea* Berk. & Curt.

Spores 12 × 25–30 μ. 1. *W. gigantea.*
Spores 15–16 × 32–40 μ. 2. *W. americana.*

WYNNEA AMERICANA

1. **Wynnea gigantea** Berk. & Curt. Jour. Linn. Soc. **9**: 424. 1867.

?Peziza macrotis Berk. Jour. Bot. and Kew Misc. **3**: 203. 1851.
?Wynnea macrotis Berk. Jour. Linn. Soc. **9**: 424. 1867.
?Midotis macrotis Sacc. Syll. Fung. **8**: 547. 1889.
Midotis gigantea Sacc. Syll. Fung. **8**: 547. 1889.

Apothecia cespitose and springing from a common stem which is about 6–7 cm. long and 2 cm. thick, deeply rugose and cracked, the individual apothecia smooth, externally becoming wrinkled when dry but not cracked like the stem, reaching a length of 5–7 cm. with the margins incurved, simple or occasionally proliferous; asci cylindric or subcylindric; spores subcymbiform, 12 × 25–30 μ; paraphyses slightly enlarged above.

Habitat unknown, but probably on the ground.

TYPE LOCALITY: Orizaba, Mexico.

DISTRIBUTION: Mexico and (India?).

ILLUSTRATIONS: Jour. Linn. Soc. **9**: *pl. 12;* Cooke, Mycographia *pl. 24, f. 93.*

2. **Wynnea americana** Thaxter, Bot. Gaz. **39**: 246. 1905. (PLATE 16.)

Sclerotium tough, subgelatinous, coriaceous on drying, irregularly lobed, variable in size, reaching a diameter of 4–5 cm., brown, the main axis becoming variously divided almost immediately after emerging from the ground, the divisions giving rise to clusters of apothecia of variable size and number; apothecia from a few to twenty-five from a single sclerotium, simple or more rarely proliferous, erect, elongate ear-shaped, very variable in size, reaching an extreme length of 6–13 cm., the average about 8 cm., the margins somewhat incurved on drying, the external surface rich blackish-brown, finely verrucose, the roughness due to projecting cells which become hair-like near the margin; hymenium even, dark purplish-red or brown; asci elongate, reaching a length of 500–540 μ and a diameter of 18 μ; spores subcymbiform, 15–16 × 32–40 μ, with the extremities apiculate, striately marked, the striations consisting of about 8 alternately light and dark bands extending the length of the spore, the entire spore when fresh filled with oil-drops; paraphyses septate, enlarged above and pale-brown.

On the ground in rich woods.

TYPE LOCALITY: Burbank, East Tennessee.

DISTRIBUTION: North Carolina to Tennessee and Ohio.

ILLUSTRATION: Thaxter, Bot. Gaz. **39**: *pl. 4* and *5* (opposite p. 246).

28. **PHILLIPSIA** Berk. Jour. Linn. Soc. **18**: 388. 1881.

Apothecia usually eccentric, attached by a very broad base which is often extended into a rather long, thick stem; hymenium concave or plane, bright-colored, the substance not shrinking much in drying; asci cylindric or subcylindric, 8-spored; spores usually striate, subhyaline; paraphyses very slender.

Type species, *Peziza domingensis* Berk.

Apothecia cup-shaped, reddish-purple.	
Apothecia not exceeding a diameter of 2 cm.	1. *P. domingensis.*
Apothecia reaching a diameter of 4–6 cm.	
Cup-shaped; substance thick.	2. *P. gigantea.*
Orbicular; substance thin.	3. *P. Chardoniana.*
Apothecia spoon-shaped, yellowish-brown.	4. *P. dochmia.*

1. **Phillipsia domingensis** Berk. Jour. Linn. Soc. **18**: 388. 1881. (PLATE 17, FIG. 2.)

Peziza domingensis Berk. Ann. Mag. Nat. Hist. II. **9**: 201. 1852.
Helotium purpuratum Kalchbr. in Thüm. Myc. Univ. *1614*. 1880.
?Peziza Harmoge Berk. & Br. Jour. Linn. Soc. **14**: 104. 1875.
?Peziza cordovensis Cooke, Hedwigia **14**: 81. 1875.
Phillipsia kermesina Kalchbr. & Cooke, Grevillea **9**: 25. 1880.
Phillipsia subpurpurea Berk. & Br. Trans. Linn. Soc. II. **2**: 69. 1883.
Lachnea crispata Sacc. Syll. Fung. **8**: 169. 1889.
Otidea domingensis Sacc. Syll. Fung. **8**: 97. 1889.

Apothecia shallow cup-shaped or subdiscoid usually unequal-sided or occasionally regular in form, attached by a broad base which is often extended into a thick stem, externally nearly white or with a pinkish tinge, fading when dry, more or less tomentose, especially near the base, reaching a diameter of 1–2 cm. the substance tough and corky when dry; hymenium concave or nearly plane, dark reddish-purple, becoming pitted as a result of shrinkage in drying which gives it the appearance of a resupinate polypore; stem often reaching a length of 1 cm. and several mm. thick, or nearly wanting; asci cylindric above, reaching a length of 300–360 μ and a diameter of 15–20 μ, gradually tapering below into a stem-like base; spores 1-seriate or with the ends slightly overlapping, ellipsoid, unequal-sided, the ends blunt or rarely narrowed, striated, the striations consisting of a few bands extending the length of the spore, containing one or two large oil-drops and often several smaller ones; paraphyses filiform, slightly enlarged above.

On old wood and bark.

TYPE LOCALITY: Santo Domingo.

DISTRIBUTION: West Indies and (Mexico?); also in Africa, Australia and Japan.

ILLUSTRATIONS: Cooke, Austr. Fungi, *pl. 19, f. 151;* Grevillea **9**: *pl. 136, f. 21;* Jour. Linn. Soc. **31**: *pl. 16, f. 7–9;* Mycologia **5**: *pl. 89* (lower figure) and *pl. 90, f. 7.*

2. **Phillipsia gigantea** Seaver, sp. nov. (PLATE 17, FIG. 1.)

Apothecia occurring singly or in clusters of 3 or 4 each with their bases more or less fused together, the individual apothecia infundibuliform, tapering below into a rather thick, stem-like base, elongated on one side or nearly symmetrical, the margin often more or less lobed, reaching a diameter of 4–6 cm. and a depth of 3–4 cm., externally nearly white or with a pinkish tinge which is somewhat darker below, delicately tomentose, the substance tough so that the apothecia maintain their form fairly well in drying; hymenium bright-purple or purplish-red, splitting into numerous vein-like markings when dry; stem thick, often reaching a length of 1–2 cm. but not clearly defined; asci cylindric above, rounded at the apex, with the operculum opening a little to one side of the center, gradually tapering below into a stem-like base, reaching a length of 500 μ and a diameter of 17–18 μ at the broadest point, 8-spored, but often with only four or five of the spores developed; spores 1-seriate, ellipsoid, unequal-sided with the ends bluntly attenuated, containing two oil-drops which often merge into one, longitudinally marked with light and dark bands, 12–14 $\times$ 25–27 μ, hyaline; paraphyses very slender, reaching a diameter of 2–3 μ at their apices.

On decaying wood of some monocotyledon.

TYPE LOCALITY: Castleton Gardens, Jamaica.

DISTRIBUTION: Known only from the type locality.

Differing from the preceding mainly in its larger size.

3. **Phillipsia Chardoniana** Seaver, Mycologia **17**: 48. 1925. (FRONTISPIECE.)

Apothecia orbicular or nearly so, subsessile, reaching a diameter of 3–5 cm., eccentrically attached, the hymenium brick-reddish, the substance thin (about 1–2 mm., not thick or corky as in other species); asci cylindric or subcylindric, reaching a length of 300 μ and a diameter of 14 μ, 8-spored; spores ellipsoid,

unequal-sided, usually containing two large oil-drops, longitudinally striated with light and dark bands, about 12 × 26 μ.

On decaying wood.

TYPE LOCALITY: Ajuntas, Porto Rico.

DISTRIBUTION: Known only from the type locality.

ILLUSTRATION: Mycologia **17**: *pl. 4.*

4. **Phillipsia dochmia** (Berk. & Curt.) Seaver, comb. nov.

Peziza dochmia Berk. & Curt.; Berk. Jour. Linn. Soc. **10**: 364. 1869.
?Peziza hirneoloides Berk. & Curt.; Berk. Jour. Linn. Soc. **10**: 365. 1869.
Otidea dochmia Sacc. Syll. Fung. **8**: 95. 1889.
?Otidea hirneoloides Sacc. Syll. Fung. **8**: 96. 1889.

Apothecia subsessile and attached by a rather thick, stem-like base, at maturity decidedly spoon-shaped, yellowish to yellowish-brown, reaching a diameter of 2–4 cm.; hymenium a little darker than the outside of the apothecium; asci cylindric above, tapering below; spores 1-seriate, or with the ends slightly overlapping, ellipsoid, strongly unequal-sided, usually containing two large oil-drops and numerous smaller ones, the striations consisting of rather broad light and dark bands extending the length of the spore; 12–15 × 25–33 μ; paraphyses slender, slightly enlarged above, adhering closely together.

On rotten wood.

TYPE LOCALITY: Cuba.

DISTRIBUTION: Cuba.

ILLUSTRATION: Cooke, Mycographia *pl. 56, f. 220, 221.*

29. **SCODELLINA** S. F. Gray, Nat. Arrang. Brit. Pl. **1**: 668. 1821.

Otidea Fuckel, Symb. Myc. 329. 1869.
Wynnella Boud. Bull. Soc. Myc. Fr. **1**: 102. 1885.

Apothecia scattered or crowded, often cespitose and occasionally fused together at the base or branched, usually large or medium sized, substipitate, rarely with a well-developed stem, usually unsymmetrical, split on one side or elongated and inclined to ear-shaped, externally usually furfuraceous; asci typically cylindric, 8-spored; spores smooth, usually containing one or two oil-drops; paraphyses filiform or very slightly enlarged above, simple or branched, straight or more often hooked at their apices.

Type species, *Peziza leporina* Batsch.

1. PHILLIPSIA GIGANTEA
2. PHILLIPSIA DOMINGENSIS
3. COOKEINA COLENSOI

Spores remaining permanently smooth.
  Spores 7–8 × 12–14 μ, containing two oil-drops. 1. *S. leporina.*
  Spores 12–13 × 20–24 μ, containing one large oil-drop. 2. *S. auricula.*
Spores becoming minutely rough. 3. *S. grandis.*

1. **Scodellina leporina** (Batsch) S. F. Gray, Nat. Arrang. Brit. Pl. **1**: 668. 1821.

Peziza leporina Batsch, Elench. Fung. 117. 1783.
Peziza onotica Pers. Syn. Fung. 637. 1801.
Scodellina onotica S. F. Gray, Nat. Arrang. Brit. Pl. **1**: 668. 1821.
Peziza onotica ochracea Fries, Syst. Myc. **2**: 48. 1822.
Otidea onotica Fuckel, Symb. Myc. 329. 1869.
Otidea leporina Fuckel, Symb. Myc. 329. 1869.
Peziza unicisa Peck, Ann. Rep. N. Y. State Mus. **26**: 81. 1874.
Aleuria leporina Gill. Champ. Fr. Discom. 40. 1879.
Aleuria onotica Gill. Champ. Fr. Discom. 40. 1879.
Otidea ochracea Seaver, Bull. Lab. Nat. Hist. State Univ. Iowa **5**: 45. 1904.

Apothecia gregarious or scattered, strongly elongated on one side and split on the opposite side to the base, externally rather bright-yellow to yellowish-brown, often becoming dull with age or when dry, reaching a height of 4–5 cm. and about half as broad; hymenium similar in color to the outside of the apothecium; asci cylindric or subcylindric, reaching a length of 200 μ and a diameter of 10–12 μ; spores 1-seriate, narrow-ellipsoid, each containing two small oil-drops, 7–8 × 12–14 μ; paraphyses slender, usually strongly curved or hooked at their apices.

On soil in woods.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to Washington and California; also in Europe.

ILLUSTRATIONS: Schaeff. Fung. Bavar. *pl. 156* (in part); Sow. Engl. Fungi *pl. 79;* Bres. Fungi Trid. *pl. 182;* Cooke, Mycographia *pl. 53, f. 210* and *pl. 54, f. 211;* Fl. Dan. *pl. 1077, f. 2;* Gonnerm. & Rab. Myc. Eu. **3**: *pl. 2, f. 4;* Nees, Syst. Pilze *pl. 37, f. 278;* Rab. Krypt.-Fl. **1**[3]: 1022, *f. 1–4;* Bull. Lab. Nat. Hist. State Univ. Iowa **5**: *pl. 10.*

2. **Scodellina auricula** (Schaeff.) Seaver, comb. nov.

Elvela auricula Schaeff. Fung. Bavar. Ind. 103. 1774.
Peziza auricula Cooke, Mycographia 124. 1876.
Aleuria auricula Gill. Champ. Fr. Discom. 40. 1879.
Otidea auricula Rehm, Hedwigia **22**: 34. 1883.

Wynnella auricula Boud. Hist. Class. Discom. Eu. 51. 1907.
Otidea neglecta Massee, Grevillea **22**: 66. 1893.

Apothecia gregarious or scattered, spoon-shaped or ear-shaped, elongated on one side and entirely split to the base of the opposite side, the margin even or more or less crenate, externally bright-yellow or becoming yellowish-brown, remaining lighter toward the base, reaching a height of 4–5 cm. and half as broad; hymenium a little darker than the outside of the apothecium, often becoming brownish-black with age; asci cylindric or subcylindric, attenuated below into a stem-like base, reaching a length of 250 μ and a diameter of 20 μ; spores 1-seriate, broad-ellipsoid, each containing one large oil-drop and numerous small ones, 12–13 $\times$ 20–24 μ, paraphyses slender, strongly enlarged above where they reach a diameter of 10 μ, filled with yellowish granules.

On the ground in woods.

TYPE LOCALITY: Europe.

DISTRIBUTION: Florida to Wisconsin and Alberta; also in Europe.

ILLUSTRATIONS: Schaeff. Fung. Bavar. *pl. 156* (in part); Bres. Fungi Trid. *pl. 73;* Boud. Ic. Myc. *pl. 250;* Cooke, Mycographia *pl. 54, f. 213.*

3. **Scodellina grandis** (Pers.) Seaver, comb. nov.

Peziza grandis Pers. Obs. Myc. **1**: 27. 1796.
Otidea grandis Massee, Brit. Fungus-Fl. **4**: 446. 1895.
?Otidea Harperiana Rehm, Ann. Myc. **2**: 34. 1904.

Apothecia cespitose or solitary, sessile or substipitate, shallow cup-shaped, the margin incurved, usually split on one side to near the base and with the split edge rolled inward, externally yellowish to dark-brownish, with a tinge of olive, pruinose above and whitish tomentose toward the base, reaching a diameter of 5–6 cm. and about half as deep; hymenium pale-yellow, lighter than the outside of the apothecium; asci cylindric above, gradually tapering below, reaching a length of 280 μ and a diameter of 12–13 μ at the widest point, 8-spored; spores 1-seriate, parallel or obliquely disposed, ellipsoid, with the ends quite strongly narrowed, containing two oil-drops, hyaline, smooth, or at maturity very minutely roughened, about 8 $\times$ 15–18 μ; paraphyses rather stout, reaching a diameter of 7–8 μ, strongly curved or hooked at their apices and densely filled with minute granules.

On soil in coniferous or mixed woods.

TYPE LOCALITY: Europe.

DISTRIBUTION: Connecticut to Pennsylvania and Wisconsin; also in Europe.

ILLUSTRATIONS: Boud. Ic. Myc. *pl. 328;* Cooke, Mycographia *pl. 105, f. 376;* Pat. Tab. Fung. *f. 480.*

DOUBTFUL SPECIES

Otidea obtecta (Schw.) Sacc. Syll. Fung. **8**: 98. 1889; *Peziza obtecta* Schw. Trans. Am. Phil. Soc. II. **4**: 170. 1832. The identity of this species is uncertain.

30. **COOKEINA** Kuntze, Rev. Gen. Pl. **2**: 849. 1891.

Peziza § *Trichoscypha* Cooke, Mycographia 252. 1879.
Trichoscypha Sacc. Syll. Fung. **8**: 160. 1889. Not *Trichoscypha* Hooker, 1862.
Pilocratera P. Henn. in Engler, Bot. Jahr. **14**: 363. 1892.

Apothecia stipitate or substipitate, bright-colored, some shade of red or yellow (especially the hymenium), externally hairy or tomentose or occasionally pruinose, the substance tough, not shrinking much in drying; hairs when present fasciculate; asci cylindric or subcylindric, usually with an abrupt stem at the base, 4–8-spored; spores hyaline or subhyaline, ellipsoid to fusoid, usually striate, the striations consisting of light and dark bands extending the length of the spore; paraphyses usually very slender.

Type species, *Peziza Tricholoma* Mont.

Apothecia clothed with well-developed hairs.
  Hairs long and conspicuous, covering the outside of the apothecium. 1. *C. Tricholoma.*
  Hairs short and inconspicuous, mostly near the margin.
    Asci 8-spored. 2. *C. sulcipes.*
    Asci 4-spored. 3. *C. tetraspora.*
Apothecia pruinose but with no well-developed hairs. 4. *C. Colensoi.*

1. **Cookeina Tricholoma** (Mont.) Kuntze, Rev. Gen. Pl. **2**: 849. 1891. (PLATE 18, FIG. 2.)

Peziza Tricholoma Mont. Ann. Sci. Nat. II. **2**: 77. 1834.
Peziza Hystrix Berk. Ann. Mag. Nat. Hist. II. **9**: 201. 1852.
Trichoscypha Tricholoma Sacc. Syll. Fung. **8**: 160. 1889.
Pilocratera Tricholoma P. Henn. in Engler, Bot. Jahr. **14**: 364. 1892.
Peziza striispora Ellis & Ev. Bull. Lab. Nat. Hist. State Univ. Iowa **4**: 69. 1896.
Sarcoscypha striispora Sacc. Syll. Fung. **14**: 754. 1899.

Apothecia stipitate or substipitate, cup-shaped with the margin slightly incurved, reaching a diameter of 1–1.5 cm. and a depth of 1 cm., externally clothed with long hairs which are more numerous around the margin, forming an incurved fringe, deep-red, fading in drying; hymenium bright-red, almost scarlet; stem often so short that the apothecia appear to be sessile and again, often 2–3 cm. long and about 2 mm. in diameter; hairs often 2–3 mm. long and 100–175 μ in diameter at the base, gradually tapering upward, whitish or pale-brown, composed of a dense fascicle of mycelial threads; asci cylindric above, reaching a length of 350–374 μ and a diameter of 20 μ, abruptly narrowed below into a short, appendage-like base; spores ellipsoid to fusoid, 12–14 $\times$ 27–33 μ, hyaline or subhyaline, containing one or two large oil-drops, usually marked with delicate, longitudinal striations; paraphyses filiform, slender, slightly enlarged upwards.

On old wood and bark.

TYPE LOCALITY: Central America.

DISTRIBUTION: West Indies, Mexico, Central America; also in South America, Australia and the Philippine Islands.

ILLUSTRATIONS: Ann. Sci. Nat. II. **2**: *pl. 4, f. 2;* Cooke, Mycographia *pl. 51, f. 202;* E. & P. Nat. Pfl. 1^1: 195, *f. 155, C–E;* Mycologia **5**: *pl. 88* (lower figure) and *pl. 90, f. 1–2.*

2. **Cookeina sulcipes** (Berk.) Kuntze, Rev. Gen. Pl. **2**: 849. 1891. (PLATE 18, FIG. 1.)

Peziza sulcipes Berk. London Jour. Bot. **1**: 141. 1842.
Peziza Hindsii Berk. London Jour. Bot. **1**: 456. 1842.
?Peziza Afzelii Fries, Nov. Acta Reg. Soc. Scient. Upsal. III. **1**: 121. 1855.
Trichoscypha Hindsii Sacc. Syll. Fung. **8**: 161. 1889.
Trichoscypha sulcipes Sacc. Syll. Fung. **8**: 161. 1889.
?Trichoscypha Afzelii Sacc. Syll. Fung. **8**: 161. 1889.
Cookeina Hindsii Kuntze, Rev. Gen. Pl. **2**: 849. 1891.
?Cookeina Afzelii Kuntze, Rev. Gen. Pl. **2**: 849. 1891.
?Pilocratera Engleriana P. Henn. in Engler, Bot. Jahr. **14**: 363. 1892.
Pilocratera Hindsii Lindau in E. & P. Nat. Pfl. **1**: 195. 1897.
Geopyxis elata Massee, Bull. Royal Gardens Kew, **1898**: 123. 1898.

Apothecia stipitate or substipitate, cup-shaped, reaching a diameter of 1–2 cm. and a depth of 1 cm., or sometimes more shallow, externally often marked with several concentric rings near the margin which is fringed with very short hairs, yellowish; hymenium deep-orange to nearly scarlet, faded in dried specimens; stems often so short that the cups appear to be sessile,



1. COOKEINA SULCIPES
2. COOKEINA TRICHOLOMA

but occasionally reaching a length of 3 cm., and about 2 mm. in diameter; hairs comparatively short, subconical in form, reaching a length of 400–500 μ and a diameter of 75–100 μ at the base, gradually tapering upwards, composed of a fascicle of mycelial threads, pale-yellow; asci cylindric, reaching a length of 300–350 μ and a diameter of 20 μ, with a short appendage-like base; spores ellipsoid, with the ends slightly narrowed, hyaline or subhyaline, containing one or two oil-drops, 14–18 × 27–33 μ, marked with delicate, longitudinal striations; paraphyses filiform, slightly enlarged above.

On old wood and bark.

TYPE LOCALITY: Surinam, South America.

DISTRIBUTION: West Indies and New Mexico; also reported from South America and Australia.

ILLUSTRATIONS: London Jour. Bot. **1**: *pl. 15* (in part); Cooke, Mycographia *pl. 51, f. 199*, 200; Cooke, Australian Fungi *pl. 20, f. 153;* E. & P. Nat. Pfl. **1**[1]: 195, *f. 155, F–G;* Engler, Bot. Jahr. **14**: *pl. 6, f. 9;* Mycologia **5**: *pl. 88* (upper figure), *pl. 90, f. 3–5.*

3. **Cookeina tetraspora** Seaver; Stevenson, Jour. Depart. Agr. Porto Rico **2**: 160. 1918. (hyponym); Mycologia **17**: 45. 1925.

Apothecia gregarious, closely crowded, or occasionally becoming confluent with age, at first short-stipitate and turbinate, gradually expanding, becoming slightly concave or nearly plane, externally whitish and slightly tomentose, becoming strongly wrinkled when dry, reaching a diameter of 4 mm. and a height of 2 mm., yellowish or orange the margin fimbriate; fimbriae consisting of conical, fasciculate masses of delicate hairs, each reaching a length of 150 μ and a diameter of 100 μ at the base, hyaline; asci clavate, gradually tapering below into a slender stem, reaching a length of 200 μ and a diameter of 12–14 μ, 4-spored; spores ellipsoid, unequal-sided, usually containing two large oil-drops, appearing to have a long slender apiculus at the basal end which usually disappears when removed from the ascus, 10–11 × 24–27 μ, often showing faint striations; paraphyses filiform, slightly enlarged above.

On decaying leaves of some palm.

TYPE LOCALITY: El Yunque, Porto Rico.

DISTRIBUTION: Known only from the type locality.

4. **Cookeina Colensoi** (Berk.) Seaver, Mycologia **5**: 191. 1913. (PLATE 17, FIG. 3.)

Peziza Colensoi Berk. in Hooker's Fl. New Zealand **2**: 200. 1855.
Peziza aluticolor Berk. Proc. Linn. Soc. **13**: 176. 1873.
Phillipsia Venezuelae Berk. & Curt.; Cooke, Mycographia 120. 1876.
Geopyxis aluticolor Sacc. Syll. Fung. **8**: 64. 1889.
Sarcoscypha Colensoi Sacc. Syll. Fung. **8**: 157. 1889.
Peziza Venezuelae Massee, Jour. Linn. Soc. **31**: 473. 1896.

Apothecia stipitate or substipitate, shallow cup-shaped, reaching a diameter of 1–2 cm. and a depth of about 5 mm., dried specimens pale-yellow (probably much brighter when fresh), marked with concentric rings about the margin, externally covered with loose cells which sometimes approach rudimentary hairs but with no well-developed hairs, strongly wrinkled when dry especially near the base of the apothecium; stem very short and sometimes almost wanting, often not more than 1 mm. in length, never long as in other species; asci cylindric above, reaching a length of 400–475 μ and a diameter of 20 μ, gradually tapering below; spores 1-seriate or with the ends slightly overlapping, fusoid, with the ends quite strongly narrowed, containing one or two large oil-drops and granular within, the striations consisting of several broad, longitudinal bands extending the length of the spore, 12–15 $\times$ 30–40 μ; paraphyses filiform, scarcely enlarged above.

On wood and bark.

TYPE LOCALITY: New Zealand.

DISTRIBUTION: West Indies; also in New Zealand, Australia and Africa.

ILLUSTRATIONS: Hooker's Fl. New Zealand **2**: *pl. 105, f. 5;* Cooke, Mycographia *pl. 50, f. 198; pl. 52, f. 206;* Jour. Linn. Soc. **31**: *pl. 16, f. 4–5;* Hedwigia **39**: *pl. 11, f. 21;* Mycologia **5**: *pl. 89* (upper figure) and *pl. 90, f. 8.*

31. **PLECTANIA** Fuckel, Symb. Myc. 323. 1869.

Microstoma Milde, Bot. Zeit. **10**: 208. 1852. Not *Microstoma* Bruch. 1846.
Anthopeziza Wettst. Verh. Zool.-Bot. Wien **35**: 383. 1886.
Sarcoscypha Sacc. Syll. Fung. **8**: 153. 1889.

Apothecia at first subglobose, expanding but with the margin often incurved and even or laciniate, usually stipitate but with the stem very variable in length or sometimes almost wanting, externally more or less hairy; hymenium bright-colored; inclined



1. PLECTANIA COCCINEA
2. PLECTANIA HIEMALIS

to scarlet; hairs when present hyaline or very slightly yellowish; asci 8-spored; spores ellipsoid, hyaline or subhyaline, smooth; paraphyses filiform and slender.

Type species, *Peziza coccinea* Scop.

Apothecia large, 2–3 cm. in diameter.	1. *P. coccinea.*
Apothecia rarely exceeding a diameter of 1 cm.	
Hairs long, bristly and very conspicuous.	2. *P. floccosa.*
Hairs inconspicuous, consisting of delicate tomentum.	
Tomentum scant; spores 10–12 × 20–22 μ.	3. *P. occidentalis.*
Tomentum abundant; spores 13–16 × 35–50 μ.	4. *P. hiemalis.*

1. **Plectania coccinea** (Scop.) Fuckel, Symb. Myc. 324. 1869. (PLATE 19, FIG. 1.)

Elvela coccinea Scop. Fl. Car. **2**: 479. 1772.
Peziza coccinea Jacq. Fl. Austr. **2**: *pl. 163.* 1774.
Peziza epidendra Bull. Herb. Fr. *pl. 467, f. 3.* 1789.
Peziza poculiformis Hoffm. Veg. Crypt. **2**: 27. 1790.
?Peziza pulcherrima Raf. Med. Rep. II. **5**: 362. 1808.
Macroscyphus coccineus S. F. Gray, Nat. Arrang. Brit. Pl. **1**: *pl. 672.* 1821.
Lachnea coccinea Gill. Champ. Fr. Discom. 66. 1880.
Sarcoscypha coccinea Sacc. Syll. Fung. **8**: 154. 1889.
Peziza Dudleyi Peck, Ann. Rep. N. Y. State Mus. **47**: 23. 1894.
Geopyxis coccinea Massee, Brit. Fungus-Fl. **4**: 377. 1895.

Apothecia rather deep cup-shaped or infundibuliform, stipitate or rarely subsessile, the margin of the cup usually strongly incurved, externally nearly white and more or less floccose with hyaline, flexuous mycelial hairs; hymenium deeply concave, scarlet, fading when dry; stem stout, 4–5 mm. thick and of variable length, often 2–3 cm. long and again almost wanting, the length of the stem varying according to the depth at which the sticks are buried; asci very long, cylindric, gradually tapering into a stem-like base, reaching a length of 400–500 μ and a diameter of 12–14 μ; spores mostly 1-seriate, long-ellipsoid, often containing two large oil-drops and numerous small ones, 10–12 × 26–40 μ; paraphyses slightly enlarged above, containing numerous red granules.

On buried or partially buried sticks, early spring.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to Washington and West Virginia; also in Europe.

ILLUSTRATIONS: Boud. Ic. Myc. *pl. 322, 323;* Bull. Herb. Fr. *pl. 467, f. 3;* Cooke, Mycographia *pl. 25, f. 95;* E. & P. Nat. Pfl. 1^1: 195, *f. 155, A–B;* Gill. Champ. Fr. Discom. *pl. 58, f. 1;* Hoffm. Veg. Crypt. **2**: *pl. 7, f. 5;* Bull. Soc. Myc. Fr. *pl. 1, f. 1.*

Melvin R. Gilmore of the Museum of the American Indian, New York City writes: The Scarlet Cup-Fungus, *Sarcoscypha coccinea* (Jacq.) Sacc., is used medicinally by the Oneida, and probably by the other tribes of the Iroquois Six Nations. In such use this fungus is dried, pulverized, and applied as a styptic. Most particularly this is the medicament used for application to the navel of a newborn child when the umbilical cord is cut, and is not healing well. In old times this pulverized fungus was kept in place on the part under a bandage of soft-tanned deer-skin.

2. **Plectania floccosa** (Schw.) Seaver, comb. nov. (PLATE 20, FIG. 1.)

Peziza floccosa Schw. Trans. Am. Phil. Soc. II. **4**: 172. 1832.
Geopyxis floccosa Morgan, Jour. Myc. **8**: 188. 1902.
Sarcoscypha floccosa Sacc. Syll. Fung. **8**: 156. 1889.

Apothecia gregarious or occasionally cespitose, stipitate, infundibuliform, with the margin usually strongly curved especially in young plants, reaching a diameter of 5–8 mm. and a depth of 1 cm., externally clothed with very long, rigid, hyaline hairs which give to the apothecium a very shaggy appearance; stem slender, gradually expanding into the apothecium above, very short or reaching a length of several cm., the length varying with the depth at which the sticks on which the plants grow are buried in the ground; hairs about 15–18 μ in diameter at the base, gradually tapering into a bristle-like apex, reaching a length of more than 1 mm., septate, thick-walled; asci cylindric or subcylindric, rather abruptly narrowed into a long stem-like base, reaching a length of 300–325 μ (not including the constricted portion below) and a diameter of 20 μ at the apex; spores 1-seriate or with the ends partially overlapping, ellipsoid, with the ends rather strongly narrowed, smooth, hyaline or slightly yellowish, 15–17 $\times$ 20–35 μ; paraphyses slender, slightly enlarged above, reaching a diameter of 3 μ at their apices.

On buried sticks in woods.

TYPE LOCALITY: Nazareth, Pennsylvania.

DISTRIBUTION: Massachusetts to Iowa, Missouri and Virginia.

ILLUSTRATIONS: Grevillea **1**: *pl. 1, f. 2;* Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 22, f. 1;* Atkinson, Mush. *f. 222.*

EXSICCATI: N. Am. Fungi *435;* Rab.-Winter, Fungi Eu. *3171.*



1. PLECTANIA FLOCCOSA
2. PLECTANIA OCCIDENTALIS

The hairy cups of this species are rather common in woods in early spring.

3. **Plectania occidentalis** (Schw.) Seaver, comb. nov. (PLATE 20, FIG. 2; 45, FIG. 23.)

Peziza occidentalis Schw. Trans. Am. Phil. Soc. II. **4**: 171. 1832.
Peziza hesperidea Cooke & Peck, Grevillea **1**: 5. 1872.
Geopyxis occidentalis Morgan, Jour. Myc. **8**: 188. 1902.
Sarcoscypha occidentalis Sacc. Syll. Fung. **8**: 154. 1889.
Geopyxis hesperidea Sacc. Syll. Fung. **8**: 63. 1889.

Apothecia gregarious or cespitose, shallow cup-shaped to discoid, stipitate, externally whitish, reaching a diameter of 1 cm.; hymenium usually concave, bright-red, almost scarlet; stem varying in length according to the depth at which the sticks on which the plants grow are buried in the ground but often reaching a length of 2–3 cm. and a diameter of 2 mm.; asci cylindric or subcylindric, reaching a diameter of 15–18 μ at the apex, tapering below into a long stem-like base; spores 1-seriate, parallel with the ascus or oblique and with the ends slightly overlapping, ellipsoid, usually with two oil-drops one in either end which are often surrounded with smaller oil-drops and granules, hyaline or slightly yellowish, 10–12 $\times$ 20–22 μ; paraphyses slender, slightly thickened above, reaching a diameter of 3–4 μ at their apices.

On buried or partially buried sticks in wood.

TYPE LOCALITY: Muskingum, Ohio.

DISTRIBUTION: Connecticut to Iowa, Kansas and Louisiana.

ILLUSTRATIONS: Grevillea **1**: *pl. 1, f. 1;* Cooke, Mycographia *pl. 50, f. 195;* Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 22, f. 2.*

EXSICCATI: N. Am. Fungi *436.*

Rather common but not abundant in woods in midsummer.

4. **Plectania hiemalis** (Nees & Bernst.) Seaver, comb. nov. (PLATE 19, FIG. 2.)

Microstoma hiemale Nees & Bernst.; Milde, Bot. Zeit. **10**: 208. 1852.
Peziza cruciata Fries, Nov. Symb. Myc. 229. 1855.
Peziza prostrata Fries, Nov. Symb. Myc. 230. 1855.
Peziza mirabilis Borszcz. Beit. Pflanzenkunde **10**: 61. 1857.
Sclerotinia baccata Fuckel, Symb. Myc. 331. 1869.
Anthopeziza Winteri Wettst. Verh. Zool.-Bot. Wien **35**: 383. 1886.
Sarcoscypha alpina Ellis & Ev. Bull. Torrey Club **24**: 281. 1897.
Sarcoscypha cruciata Sacc. Syll. Fung. **8**: 154. 1889.

Sarcoscypha prostrata Sacc. Syll. Fung. **8**: 155. 1889.
Lachnea mirabilis Phill. Grevillea **18**: 83. 1890.

Apothecia solitary or cespitose, stipitate, at first infundibuliform, becoming expanded and at maturity shallow cup-shaped, the margin lobed and often reflexed, reaching a diameter of 1 cm., externally clothed with soft, white, flexuous hairs; hymenium bright-red, almost scarlet; stem slender, reaching a length of 3–4 cm. and a diameter of 1–2 mm., gradually expanding into the cup; hairs hyaline, blunt, and of about the same diameter throughout their entire length, reaching a diameter of 6–7 μ; asci cylindric, reaching a length of 200–250 μ and a diameter of 20–22 μ; spores 1-seriate, ellipsoid, subinequilateral, narrowed toward either end, usually containing one large oil-drop and one or two smaller ones, 13–16 $\times$ 25–50 μ, smooth; paraphyses slender, slightly enlarged above.

On the ground in woods.

TYPE LOCALITY: Near Breslau, Germany.

DISTRIBUTION: Minnesota to Colorado; also in Europe.

ILLUSTRATIONS: Verh. Zool.-Bot. Wien **35**: *pl. 16;* Cooke, Mycographia, *pl. 25, f. 89;* Beit. Pflanzenkunde **10**: *pl. 4, 5.*

The only North American specimen of this species seen is a single plant, the type of *Sarcoscypha alpina*. The accompanying photograph is from a plant in the Torrey collection from an unknown locality.

DOUBTFUL SPECIES

Sarcoscypha roseotincta Clements, Bot. Surv. Nebr. **4**: 11. 1896. A delicate rose-colored species with spores 5 $\times$ 10 μ described from material collected on horse dung. Nothing is known of the species in addition to that given in the original description.

32. **BULGARIA** Fries, Syst. Myc. **2**: 166. 1822.

Burcardia Schmidel, Ic. Pl. 261. 1797. Not *Burcardia* Schreb. 1789.
Sarcosoma Casp; Rehm in Rab. Krypt.-Fl. 1^3: 497. 1896.

Apothecia stipitate or sessile, externally hairy or tomentose, the tissue gelatinous, when dry becoming coriaceous or corky; hymenium partly or entirely covering the upper surface, lighter than the exterior which is black or blackish, the entire apothecium becoming strongly wrinkled when dry; asci cylindric or subcylindric, 8-spored; spores ellipsoid or subellipsoid; hyaline or subhyaline; paraphyses slender.

Type species, *Burcardia globosa* Schmidel.

The genus *Bulgaria* was established by Fries and based on an operculate form as indicated above. However, at the same time he included in the genus *Peziza inquinans* Pers. which is an inoperculate species and according to present notions not congeneric with the type of the genus. Unfortunately Saccardo in his "Sylloge Fungorum" has treated *Bulgaria inquinans* first so that it has come to be looked upon by collectors as the typical *Bulgaria*, especially since it is much more commonly collected than the authentic type. We are therefore using the genus *Bulgaria* in the restricted, and morphologically correct sense for the operculate forms. Since the inoperculate species are not treated in this work, *Bulgaria inquinans* (Pers.) Fries is left out of consideration, but will be properly dealt with under the Inoperculates when these are monographed.

Apothecium strongly swollen below into a large globose base. 1. *B. globosa.*

Apothecium not swollen into a globose base below the hymenium.

Hymenium light-colored when fresh, with a reddish tinge. 2. *B. rufa.*

Hymenium dark-colored, blackish. 3. *B. melastoma.*

1. **Bulgaria globosa** Fries, Syst. Myc. **2**: 166. 1822. (TEXT FIG. 2.)

Burcardia globosa Schmidel, Ic. Pl. 261. 1797.
Peziza Burcardia Pers. Syn. Fung. 632. 1801.
Burckhardia globosa Kuntze, Rev. Gen. Pl. **2**: 845. 1891.
Sarcosoma globosa Casp.; Rehm in Rab. Krypt.-Fl. 1^3: 498. 1896.

Apothecia gregarious, at first nearly globose, ranging in size from 1–5 cm. or rarely larger, at length ovoid with the hymenium appearing as a depression in the upper surface, becoming expanded but usually much narrower than the gelatinous base which remains strongly swollen and bulged out below the hymenium, externally, brownish-black, the substance consisting of a tough outer rind and a thick, gelatinous interior which shrinks much in drying, giving the exterior a peculiar wrinkled appearance, minutely brownish-tomentose; hymenium slightly concave, darker than the outside of the apothecium; asci cylindric or subcylindric above; spores 1-seriate, ellipsoid, about 10–12 $\times$ 23–26 μ; paraphyses dark-brown adhering together in masses, slender below, enlarged above.

On the ground among mosses in fir woods.

TYPE LOCALITY: Europe.

DISTRIBUTION: Ontario, Canada; also in Europe.

ILLUSTRATION: Schmidel, Ic. Pl. 69, *f. 1–13.*

On May 23, 1919 an excellent specimen of this species was received from Prof. J. H. Faull of the University of Toronto, Toronto, Canada. Although previously recorded from Ottawa, this is the only fresh specimen of the species seen by the writer. American specimens agree well with Schmidel's excellent illustrations except that the hymenium in our specimen is much more expanded which is probably due to a more advanced stage of development. This is a part of the material reported in Mycologia (**11**: 293. 1919) as *Bulgaria platydiscus* Casp. which is said to differ from *B. globosa* in the size of the spores. The writer has had no opportunity to study living material of either species.

2. **Bulgaria rufa** Schw. Trans. Phil. Soc. II. **4**: 178. 1832.

Apothecia gregarious or cespitose, sessile or more often short-stipitate, attaining a diameter of 2–3 cm. at first closed, finally opening and becoming shallow cup-shaped, with the margin incurved, externally blackish-brown and covered over with clusters of hairs, the substance tough, internally with a gelatinous layer several mm. thick, giving to the fresh plants a rubbery consistency, on drying becoming coriaceous and strongly wrinkled as a result of the gelatinous tissue within; hymenium slightly concave, pale-reddish or reddish-brown; stem reaching a length of 1 cm. and a diameter of 4–5 mm., attached below by means of a dense mass of black mycelium which reaches a diameter of 7–8 μ; hairs blackish-brown, similar to the mycelium at the base of the stem but shorter; asci cylindric above, gradually tapering below into a long stem-like base, reaching a length of 275–300 μ and a diameter of 12–14 μ; spores 1-seriate, ellipsoid, with the ends strongly narrowed, hyaline, granular within, 10 $\times$ 20 μ; paraphyses filiform, scarcely enlarged above.

On the ground in woods attached to sticks which are buried under leaf-mould and soil.

TYPE LOCALITY: Bethlehem, Pennsylvania.

DISTRIBUTION: New York to Minnesota, Missouri and North Carolina.

ILLUSTRATION: Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 37, f. 1.*

EXSICCATI: N. Am. Fungi *449;* Fungi Columb. *248;* Rav. Fungi Car. **4**: 23; Rab.-Winter, Fungi Eu. *3467* (as *Peziza pellita* Cooke & Peck).

Frequent in woods in midsummer.

3. **Bulgaria melastoma** (Sow.) Seaver, comb. nov.

Peziza melastoma Sow. Engl. Fungi *pl. 149* (with descr.). 1799.
Peziza rhizopus Alb. & Schw. Consp. Fung. 317. 1805.
Calycina melastoma S. F. Gray, Nat. Arrang. Brit. Pl. 670. 1821.
Peziza atrorufa Grev. Scot. Crypt. Fl. 315 (with descr.). 1828.
Peziza crenulata Fuckel, Bot. Zeit. **19**: 250. 1861.
Plectania melastoma Fuckel, Symb. Myc. 324. 1869.
Peziza hirtipes Cooke, Bull. Buff. Acad. Sci. **2**: 290. 1875.
Lachnea melastoma Gill. Champ. Fr. Discom. 66. 1880.
Peziza rhizomorpha Ellis & Ev. Jour. Myc. **4**: 98. 1888.
Plectania hirtipes Sacc. Syll. Fung. **8**: 163. 1889.
Plectania rhizomorpha Sacc. Syll. Fung. **8**: 164. 1889.
Urnula melastoma Boud. Hist. Class. Discom. Eu. 55. 1907.
Sarcoscypha melastoma Hone, Minn. Bot. Studies **4**: 97. 1909.
Scutellinia melastoma Kuntze, Rev. Gen. Pl. **3**^{3}: 520. 1898.
Scutellinia hirtipes Kuntze, Rev. Gen. Pl. **3**^{3}: 520. 1898.
Scutellinia rhizomorpha Kuntze, Rev. Gen. Pl. **3**^{3}: 520. 1898.

Apothecia gregarious or cespitose, short-stipitate or sessile, at first globose or subglobose, gradually opening with a circular aperture, but with the margin remaining incurved, entirely black or brownish-black, externally tomentose, the substance tough outside with an internal gelatinous layer, becoming much wrinkled in drying, reaching a diameter of 1–2 cm. and of about the same depth; hymenium deeply concave, a little paler than the outside of the apothecium, smooth and glistening, splitting into vein-like markings when dry; stem when present, short and stout, attached to the substratum by a dense tuft of black mycelium which penetrates into the humus and binds it together; hairs brown, flexuous, long, about 7 μ in diameter, often interspersed with brick-red granules; asci cylindric, reaching a length of 500–600 μ and a diameter of 15–18 μ, tapering below into a long, slender, stem-like base; spores ellipsoid, with the ends strongly narrowed, smooth, hyaline, 9–10 $\times$ 20–25 μ; paraphyses filiform, scarcely enlarged above, yellowish-brown.

Attached to sticks and twigs in woods.

TYPE LOCALITY: Great Britain.

DISTRIBUTION: Maine to Manitoba, Washington, Mexico and the West Indies; also in Europe, New Zealand and Australia.

ILLUSTRATIONS: Alb. & Schw. Consp. Fung. *pl. 1, f. 4;* Boud. Ic. Myc. *pl. 342;* Cooke, Mycographia *pl. 27, f. 103, 104;* Gill. Champ. Fr. Discom. *pl. 58, f. 3;* Grev. Scot. Crypt. Fl. *pl. 315;* Sow. Engl. Fungi *pl. 149.*

The generic position of this species is very uncertain as indicated by the synonymy. It shows affinities with both *Bulgaria* and *Urnula* but does not seem to fit either genus particularly well. What appears to be this species is very abundant in Bermuda on the bark of Bermuda cedar, *Juniperus bermudiana*, but never on anything else. Since the host is an endemic species the fungus may also be found to differ and be itself endemic. The writer has not yet had the opportunity to clear up this point but reference will be made to it in later publications.

DOUBTFUL SPECIES

Peziza aurantiopsis Ellis, Bull. Torrey Club **9**: 18. 1882; *Lachnea aurantiopsis* Sacc. Syll. Fung. **8**: 180. 1889; *Scutellinia aurantiopsis* Kuntze, Rev. Gen. Pl. **2**: 869. 1891. Apothecia gregarious, sessile, attached to the substratum by the entire under surface except the extreme margin which is free and slightly elevated and more or less incurved, externally brownish-black, the under side covered with numerous projecting folds, clothed with a dense growth of black mycelium, the substance within white, several mm. thick and in dried plants decidedly corky; hymenium pale-yellow, darker in dried plants; mycelium pale-brown, thin-walled, branched, septate, and externally often minutely rough; asci cylindric above, gradually tapering below into a stem-like base, reaching a length of 300 μ and a diameter of 17–18 μ; spores 1-seriate or with the ends slightly overlapping, broad-ellipsoid, granular within, hyaline or slightly yellowish, 16–18 $\times$ 27–33 μ; paraphyses slender, slightly enlarged above and hyaline or nearly so, reaching a diameter of 3–4 μ.

On bare soil and on decaying wood and leaves in low sandy oak and pine woods.

In general appearance this species resembles a *Bulgaria*. Dried specimens examined show the substance to be thick and decidedly corky and in this closely resembling a *Phillipsia*, although in other respects it does not seem to fit the genus well. Its generic position is therefore uncertain. In addition to the type, excellent dried specimens have been sent by Dr. L. O. Overholts from Pennsylvania. No living plants have been seen by the writer.

33. **URNULA** Fries, Summa Veg. Scand. 364. 1849.

Cenangium § *Microcrater* Endl. Gen. Pl. 32. 1836.
Chorioactis Kupfer, Bull. Torrey Club **29**: 142. 1902.

Apothecia opening with a stellate aperture or by a number of transverse slits through the apex, leaving the margin crenate

URNULA CRATERIUM

or split into a number of rays, externally tomentose; asci cylindric or subcylindric, 8-spored; spores ellipsoid to fusoid, hyaline or subhyaline; paraphyses very slender, often branched.

Type species, *Peziza Craterium* Schw.

Apothecium with a crenate margin, not splitting *Geaster*-like. 1. *U. Craterium.*
Apothecium splitting *Geaster*-like into rays. 2. *U. Geaster.*

1. **Urnula Craterium** (Schw.) Fries, Nova Acta Soc. Sci. Upsal. III. **1**: 122. 1851. (PLATE 21.)

Peziza Craterium Schw. Schr. Nat. Ges. Leipzig **1**: 117. 1822.
Cenangium Craterium Fries, Elench. Fung. **2**: 21. 1827.
Dermea Craterium Schw. Trans. Am. Phil. Soc. II. **4**: 237. 1832.
Peziza adusta Schulz. Verh. Zool.-Bot. Wien **16**: 62. 1866.
Craterium microcrater Haszl. Verh. Zool. Bot. Wien **37**: 167. 1887.
Scutellinia adusta Kuntze, Rev. Gen. Pl. **3**^{3}: 520. 1893.
Geopyxis Craterium Rehm in Rab. Krypt.-Fl. **1**^{3}: 974. 1896.
Plectania adusta Sacc. Syll. Fung. **8**: 164. 1889.

Apothecia at first closed above, finally opening by a star-shaped aperture which leaves the margin notched and infolded, externally black or brownish-black and clothed with a dense covering of tomentum, reaching a diameter of 3–4 cm. and a depth of 4–6 cm., the substance tough and leathery; hymenium brownish-black, a little paler than the outside of the apothecium; stem reaching a length of 3–4 cm. and a diameter of 5–8 mm., even or very slightly lacunose near the base, black or brownish-black and attached to the substratum by a dense mass of black mycelium, gradually expanding above into the apothecium; hairs variable in length, thin-walled sparingly septate, flexuous, about 10 μ in diameter and of nearly uniform thickness throughout their entire length; asci cylindric above, tapering below into a long stem-like base, reaching a length of 600 μ and a diameter of 15–17 μ; spores 1-seriate, broad-ellipsoid, smooth, hyaline, 12–14 $\times$ 25–35 μ; paraphyses filiform, slightly enlarged above, pale-brown.

On the ground in deciduous woods, attached to buried or partially buried sticks.

TYPE LOCALITY: North Carolina.

DISTRIBUTION: New York to North Dakota, Alabama and Mississippi; also in Europe.

ILLUSTRATIONS: Boud. Ic. Myc. *pl. 341;* Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 39, f. 2;* Bull. Soc. Myc. Fr. **14**:

pl. 11, f. 3; Freeman, Minn. Pl. Diseases *f. 61–62;* Minn. Bot. Studies **4**: *pl. 16;* Nees & Henry, Syst. Pilze *pl. 20, f. 1–4* (as *Microcrater*); Schr. Nat. Ges. Leipzig **1**: *pl. 1, f. 7–11.*

EXSICCATI: N. Am. Fungi *982;* Fungi Columb. *1890;* Rab.-Winter, Fungi Eu. *2968;* Rav. Fungi Am. Exsicc. *633;* Rav. Fungi Car. **1**: *39;* Rehm, Ascom. *2002.*

Common and conspicuous in woods in early spring.

2. **Urnula Geaster** Peck, Ann. Rep. N. Y. State Mus. **46**: 39. 1893. (PLATE 22.)

Chorioactis Geaster Bull. Torrey Club **29**: 142. 1902.

Apothecia at first elongated and closed or nearly closed at the top, reaching a diameter of 5 cm. and a depth of 10 cm., opening by several transverse fissures and finally splitting *Geaster*-like into 3–6 rays, the clefts extending nearly to the base of the apothecium, externally clothed with a dense covering of brownish tomentum, reaching a diameter of 10–12 cm. when expanded, substance tough and leathery, soft and spongy when dry; hymenium at first white or whitish, becoming yellowish with age; stem 1–2 cm. in diameter, and reaching a length of 1–3 cm., similar in color to the outside of the apothecium; hairs straight or bent and twisted into various forms, reaching a diameter of 8 μ and of nearly uniform thickness throughout their entire length, rather thin-walled, sparingly septate, deeply colored and minutely roughened on their outer surface, blunt at their apices, variable in length; asci cylindric or subcylindric, reaching a length of 600–700 μ and a diameter of 20–23 μ; spores 1-seriate, fusoid, often unequal-sided, hyaline or subhyaline, 12–16 $\times$ 50–70 μ; paraphyses strongly thickened above.

Attached to sticks of *Ulmus crassifolia* on the ground.

TYPE LOCALITY: Austin, Texas.

DISTRIBUTION: Texas.

ILLUSTRATION: Bot. Gaz. **49**: 184, *f. 1; 185, f. 2; 186, f. 3* and *pl. 12.*

EXCLUDED SPECIES

Urnula terrestris (Niessl) Sacc. Syll. Fung. **8**: 550. 1889; *Podophacidium terrestre* Niessl in Rab. Fungi Eu. *1152.* 1868; *Phacidium terrestre* Phill. Grevillea **18**: 86. 1890. While this species has been placed in the genus *Urnula* by Saccardo, examination of cotype material gives no indication that the asci are operculate as is true of other species of the genus *Urnula.* The genus *Podophacidium* was founded by Niessl on this species and the genus has been placed in the Phacidiaceae, a family of inoperculate ascomycetes where it doubtless belongs. North American specimens collected by Dr. B. O. Dodge in Wisconsin have been examined.

URNULA GEASTER

34. **PAXINA** Kuntze, Rev. Gen. Pl. **2**: 864. 1891.

Acetabula Fuckel, Symb. Myc. 330. 1869. Not *Acetabulum* Boehmer-Ludwig, 1760.

Macropodia Fuckel, Symb. Myc. 331. 1869. Not *Macropodia* R. Brown, 1812.

Fuckelina Kuntze, Rev. Gen. Pl. **2**: 852. 1891. Not *Fuckelina* Sacc., 1875.

Phleboscyphus Clements, Bull. Torrey Club **30**: 93. 1903.

Cyathipodia Boud. Hist. Class. Discom. Eu. 39. 1907.

Apothecia stipitate or substipitate, cup-shaped, usually regular or occasionally slightly contorted, externally hairy or tomentose, the substance tough and approaching leathery; stem slender or, more often stout, long or very short and immersed in the ground when the apothecia appear sessile, even or irregularly lacunose; asci cylindric to clavate, 8-spored; spores ellipsoid to fusiform, smooth or rough, hyaline or at most slightly yellowish; paraphyses rather stout.

Type species, *Peziza Acetabulum* L.

Spores remaining permanently smooth.
- Stem thick and strongly lacunose or corrugated.
  - Corrugations extending up the sides of the apothecium nearly to the margin. — 1. *P. Acetabulum.*
  - Corrugations extending only to the base of the apothecium.
    - Stem long, reaching a length of 4–5 cm. — 2. *P. macropus.*
    - Stem short, not exceeding 2–3 cm. in length.
      - Stem flattened. — 3. *P. platypodia.*
      - Stem rounded or angular, not flattened.
        - Stem and outside of the apothecium nearly white; stem short, rarely reaching a diameter of 1 cm. — 4. *P. leucomelas.*
        - Stem and outside of the apothecium cinereous to brown; stem reaching a length of 2–3 cm. — 5. *P. sulcata.*
        - Stem and outside of the apothecium olivaceous; stem reaching a length of 2–3 cm. — 6. *P. Barlae.*
- Stem slender, even or only slightly corrugated.
  - Apothecia and stems cinereous to brown; hairs hyaline.
    - Stems even, at least usually so.
      - Reaching a length of 3–4 cm. — 7. *P. hispida.*
      - Reaching a length of scarcely 1 cm. — 8. *P. subclavipes.*
    - Stems slightly corrugated. — 9. *P. Dupainii.*
  - Apothecia and stems black or blackish.
    - Spores 10–12 × 18–20 μ. — 10. *P. Corium.*
    - Spores 15 × 30–35 μ. — 11. *P. nigrella.*

Spores becoming rough at maturity.
  Spores fusiform; apothecia brown.
    Apothecia 2–3 cm. broad and high; stem strongly corrugated. 12. *P. semitosta.*
    Apothecia scarcely exceeding 2 cm. broad and high; stem not strongly corrugated. 13. *P. fusicarpa.*
  Spores ellipsoid; apothecia olivaceous. 14. *P. olivacea.*

1. **Paxina Acetabulum** (L.) Kuntze, Rev. Gen. Pl. **2**: 864. 1891. (PLATE 23.)

Peziza Acetabulum L. Sp. Pl. 1181. 1753.
Macroscyphus acetabuliforme S. F. Gray, Nat. Arrang. Brit. Pl. 672. 1821.
Acetabula vulgaris Fuckel, Symb. Myc. 330. 1869.
Aleuria Acetabulum Gill. Champ. Fr. Discom. 36. 1879.
Elvela Acetabulum Quél. Enchir. Fung. 275. 1886.
Phleboscyphus Acetabulum Clements, Crypt. Form. Colo. *298*. 1907.
Acetabula Acetabulum Underw. & Earle, Bull. Alab. Exp. Sta. **80**: 200. 1897.

Apothecia rather deep cup-shaped, reaching a depth of 2–3 cm. or more rarely expanded and shallow, externally cinereous to brownish, minutely tomentose and conspicuously ribbed, reaching a diameter of 3–5 cm., stipitate; hymenium darker than the exterior of the apothecium, brown to brownish-black; stem short and thick, reaching a length of 1–1.5 cm. and a diameter of 1 cm., deeply lacunose, the ribs of the stem being continuous with those of the apothecium, the stem not sharply defined but gradually expanding into the apothecium; tomentum consisting of bunches or fascicles of poorly developed, hyaline hairs, the component cells of which are strongly swollen but contracted at the point of union; asci cylindric above; spores obliquely 1-seriate, ellipsoid, hyaline, each containing one oil-drop, smooth, 12–14 × 18–22 μ; paraphyses straight, enlarged above, reaching a diameter of 5–6 μ at their apices.

On the ground in woods and open places.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to Minnesota, Colorado, Texas and Alabama; also in Europe.

ILLUSTRATIONS: Bull. Herb. Fr. *pl. 485, f. 4;* Vaill. Bot. Paris *pl. 13, f. 1;* Gill. Champ. Fr. Discom. *pl. 32, f. 1;* Bres. Fungi Trid. *pl. 213, f. 1;* Cooke, Mycographia *pl. 47, f. 183;* Massee, Brit. Fungus-Fl. **4**: 290, *f. 15, 16;* Phill. Brit. Discom. *pl. 3, f. 11;* Rab. Krypt.-Fl. 1^3: 923, *f. 1–3;* Sow. Engl. Fungi *pl. 59;* Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 19, f. 3;* Bull. Soc. Myc. Fr. **3**: *pl. 5, f. 1.*



PAXINA ACETABULUM

EXSICCATI: Rav. Fungi Am. Exsicc. *764;* Fungi Columb. *1801*; Clements Crypt. Form. Colo. *298*.

2. **Paxina macropus** (Clements) Seaver, comb. nov.

Phleboscyphus macropus Clements, Bull. Torrey Club **30**: 93. 1903.
Acetabula macropus Sacc. Syll. Fung. **18**: 13. 1906.
?Cyathipodia longipes Boud. Hist. Class. Discom. Eu. 39. 1907.

Apothecia reaching a diameter of 3–4 cm. and a depth of 1 cm., externally cinereous-whitish, clothed, especially about the margin with short, septate hairs; hymenium darker than the exterior of the apothecium, brownish; stem solid, long, rather tough, lacunose, usually with three or four longitudinal ribs, reaching a length of 4–5 cm. and a diameter of 7 mm.; asci cylindric, reaching a length of 300 μ and a diameter of 20 μ, tapering below into a stem-like base; spores irregularly 1-seriate, broad-ellipsoid, hyaline, each containing one large oil-drop which nearly fills the spore, about 12–14 $\times$ 20–22 μ; paraphyses enlarged above where they reach a diameter of 9 μ.

On damp soil.

TYPE LOCALITY: Minnehaha, Colorado.

DISTRIBUTION: Wisconsin to Colorado; also in (Europe?).

3. **Paxina platypodia** (Boud.) Seaver, comb. nov.

Cyathipodia platypodia Boud. Hist. Class. Discom. Eu. 39. 1907.
Macropodia platypodia Dodge, Trans. Wisc. Acad. Sci. **17**: 1041. 1914.

Apothecia scattered, rather shallow and more or less irregular, reaching a diameter of 2–3 cm. and usually a depth of 1 cm., externally cinereous and clothed with subconic fascicles of rather poorly developed hairs; hymenium darker than the exterior of the apothecium, almost black in dried specimens; stem much flattened and more or less costate or grooved on the two opposite sides, reaching a length of 2–3 cm. and a diameter of 4–5 mm., clothed with poorly developed hairs which give a pruinose appearance; hairs hyaline or subhyaline, consisting of loosely united cells which are strongly swollen in the center but contracted at the point of union; asci cylindric above, gradually tapering below, reaching a length 100–135 μ and a diameter of 15–17 μ; spores 1-seriate, usually diagonally disposed with the ends overlapping, fusoid, 10–11 $\times$ 21–27 μ, usually containing one large oil-drop and two smaller ones; paraphyses 3–4 μ in diameter below, rather strongly enlarged above where they reach a diameter of 7–9 μ.

On the ground.

TYPE LOCALITY: France.

DISTRIBUTION: Wisconsin; also in Europe.

4. **Paxina leucomelas** (Pers.) Kuntze, Rev. Gen. Pl. **2**: 864. 1891.

Peziza leucomelas Pers. Myc. Eu. **1**: 219. 1822.
Aleuria leucomelas Gill. Champ. Fr. Discom. 37. 1879.
Acetabula leucomelas Boud. Hist. Class. Discom. Eu. 40. 1907.

Apothecia gregarious or scattered, rather deep cup-shaped, reaching a diameter of 2–3 cm. and a depth of 2 cm. or occasionally expanded and more shallow, externally whitish or cinereous and rather conspicuously tomentose; hymenium darker than the outside of the apothecium, brownish-black; stem short and thick, not usually exceeding 1 cm. in length and 5–8 mm. thick, deeply lacunose, the ribs extending to the base of the apothecium; tomentum consisting of fascicles of hyaline hairs which are made up of rather loosely united and strongly swollen cells; asci cylindric above tapering below; spores ellipsoid, smooth, each containing one large oil-drop which nearly fills the spore, 12–13 × 21–23 μ; paraphyses straight, enlarged above, reaching a diameter of 5–6 μ.

On the ground.

TYPE LOCALITY: Europe.

DISTRIBUTION: Wisconsin to California; also in Europe.

ILLUSTRATIONS: Pers. Myc. Eu. *pl. 30, f. 1;* Gill. Champ. Fr. Discom. *pl. 32, f. 2;* Cooke, Mycographia *pl. 47, f. 186;* Boud. Ic. Myc. *pl. 249;* Bull. Soc. Myc. Fr. **3**: *pl. 5, f. 2.*

5. **Paxina sulcata** (Pers.) Kuntze, Rev. Gen. Pl. **2**: 864. 1891. (PLATE 24, FIG. 2.)

Peziza sulcata Pers. Syn. Fung. 643. 1801.
Acetabula sulcata Fuckel, Symb. Myc. 330. 1869.
Acetabula Calyx Sacc. Myc. Ven. 168. 1873.

Apothecia gregarious or scattered, shallow cup-shaped or occasionally expanded or nearly plane, reaching a diameter of 2–3 cm. and a depth of 1–1.5 cm., externally minutely rough, the roughenings consisting of poorly developed hairs, the color varying from cinereous to brown; the hymenium darker than the outside of the apothecium, brownish-black; stem usually short, seldom exceeding a length of 1 cm. or rarely 2 cm., deeply lacunose, paler than the outside of the apothecium, whitish to



1. PAXINA HISPIDA
2. PAXINA SULCATA

yellowish; hairs hyaline to subhyaline, each consisting of a few loosely united cells; asci cylindric or subcylindric, reaching a length of 300 μ and a diameter of 15–18 μ; spores 1-seriate, broad-ellipsoid, containing one large oil-drop, 12–14 $\times$ 15–20 μ; paraphyses rather stout, reaching a diameter of 7 μ, yellowish.

On the ground in woods.

TYPE LOCALITY: Europe.

DISTRIBUTION: New Jersey to Iowa; also in Europe.

ILLUSTRATIONS: Pers. Syn. Fung. *pl. 5, f. 1;* Sacc. Myc. Ven. *pl. 8, f. 14–18;* Boud. Ic. Myc. *pl. 246;* Cooke, Mycographia *pl. 47, f. 185;* E. & P. Nat. Pfl. 1^1: 186, *f. 150, E–F;* Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 19, f. 2.*

6. **Paxina Barlae** (Boud.) Seaver, comb. nov.

Acetabula Barlae Boud. Hist. Class. Discom. Eu. 40. 1907.

Apothecia gregarious deep cup-shaped, externally brownish-tomentose or with a faint olive tint, reaching a diameter of 2–4 cm. and about the same depth; stem short, scarcely exceeding a length of 5 mm., paler than the outside of the apothecium, deeply lacunose; hairs consisting of loosely united cells, hyaline or faintly yellowish; asci cylindric above, gradually tapering below, reaching a length of 300–325 μ and a diameter of 20 μ; spores ellipsoid, hyaline, containing one large oil-drop, 12–13 $\times$ 20–23 μ; paraphyses gradually enlarged above, reaching a diameter of 7 μ, pale olive-brown.

On the ground.

TYPE LOCALITY: Europe.

DISTRIBUTION: Alabama to California; also in Europe.

ILLUSTRATION: Boud. Ic. Myc. *pl. 245.*

7. **Paxina hispida** (Schaeff.) Seaver, comb. nov. (PLATE 24, FIG. 1.)

Elvela hispida Schaeff. Fung. Bavar. **4**: 108. 1774.
Octospora villosa Hedw. Descr. **2**: 64. 1788.
Peziza macropus Pers. Obs. Myc. **1**: 26. 1796.
Peziza stipitata Huds. Fl. Angl. 636. 1798.
Macroscyphus macropus S. F. Gray, Nat. Arrang. Brit. Pl. 672. 1821.
?Elvela ephippium Lév. Ann. Sci. Nat. II. **16**: 240. 1841.
Macropodia macropus Fuckel, Symb. Myc. 331. 1869.
Elvela macropus Karst. Myc. Fenn. **1**: 37. 1871.
Aleuria macropus Gill. Champ. Fr. Discom. 35. 1875.
Lachnea macropus Phill. Brit. Discom. 207. 1887.

Fuckelina villosa Kuntze, Rev. Gen. Pl. **2**: 852. 1891.
?Macropodia urceolata Clements, Bull. Torrey Club **30**: 91. 1903.

Apothecia scattered or gregarious, reaching a diameter of 2–3 cm. and a depth of 1 cm., externally grayish or densely whitish-tomentose; hymenium darker than the exterior of the apothecium, light-colored when young, becoming brownish with age; stem slender, reaching a length of 3–4 cm. and a diameter of 3–4 mm. at the base, gradually tapering upward toward the base of the cup where it is reduced to about 2 mm., sharply distinguished from the apothecium, even or irregularly and sparingly lacunose, tomentose; tomentum consisting of conical bunches of hairs which cling together in fascicles, the individual hairs made up of a chain of loosely united, hyaline cells which are strongly swollen in the center and contracted at the point of union; asci cylindric or slightly clavate above, reaching a length of 250–275 μ and a diameter of 17–20 μ, gradually tapering below into a stem-like base; spores 1-seriate, obliquely disposed in the ascus, narrow-ellipsoid to fusoid, smooth, containing one to three large oil-drops and numerous smaller ones, 12–14 $\times$ 25–30 μ, paraphyses gradually enlarged above where they reach a diameter of 8 μ.

On the ground in woods.

TYPE LOCALITY: Europe.

DISTRIBUTION: Massachusetts to Minnesota and Colorado (probably throughout North America); also in Europe.

ILLUSTRATIONS: Schaeff. Fung. Bavar. *pl. 167;* Bolton, Hist. Fung. **3**: *pl. 96;* Boud. Ic. Myc. *pl. 239;* E. & P. Nat. Pfl. **1**[1]: 186, *f. 150, G–L;* Gill. Champ. Fr. Discom. *pl. 31;* Hedw. Descr. **2**: *pl. 19, f. 1–4;* Pers. Obs. Myc. *pl. 1, f. 2;* Underw. Molds Mild. & Mush. *pl. 4, f. 9–10;* Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 19, f. 1.*

EXSICCATI: Clements, Crypt. Form. Colo. *297.*

Rather common in midsummer.

8. **Paxina subclavipes** (Phill. & Ellis) Seaver, comb. nov.

Peziza subclavipes Phill. & Ellis, N. Am. Fungi *985.* 1887.
?Peziza brachypus Ellis & Ev. Jour. Myc. **4**: 55. 1888.
?Geopyxis brachypus Sacc. Syll. Fung. **8**: 68. 1889.
Elvela macropus brevis Peck, Bull. Torrey Club **29**: 74. 1902.
Macropodia subclavipes Rehm, Ann. Myc. **2**: 354. 1904.

Apothecia gregarious, reaching a diameter of 2 cm. and a depth of about 1 cm., externally cinereous to yellowish or more

rarely olivaceous-brown, rough, the roughenings consisting of bunches of hairs; hymenium darker than the outside of the apothecium, brownish; stem slender, even or nearly so, reaching an extreme length of 1 cm. but often not over a few mm., about 2 mm. thick, covered like the outside of the apothecium with tufts of hairs; hairs clining together in conical clumps, each individual hair made up of a chain of loosely united cells the cells swollen in the center but contracted at the point of union, hyaline or faintly yellowish; asci cylindric, reaching a length of 250 μ and a diameter of 15 μ; spores 1-seriate or slightly crowded near the apex of the ascus, ellipsoid, simple, containing one large oil-drop and occasionally a few smaller ones, smooth, hyaline, 9–10 $\times$ 18–23 μ; paraphyses filiform, about 3 μ in diameter below, enlarged above where they reach a diameter of 7 μ, yellowish-brown.

On the ground in woods.

TYPE LOCALITY: Newfield, New Jersey.

DISTRIBUTION: New York to Iowa, Texas and South Carolina.

EXSICCATI: Rav. Fungi Am. Exsicc. *760* (as *Peziza Corium*); N. Am. Fungi *984* (as *Peziza Corium*).

9. **Paxina Dupainii** (Boud.) Seaver, comb. nov.

Acetabula Dupainii Boud. Bull. Soc. Myc. Fr. **14**: 17. 1898.
Elvela Faulknerae Copeland, Ann. Myc. **2**: 509. 1904.
Cyathipodia Dupainii Boud. Hist. Class. Discom. Eu. 39. 1907.

Apothecia gregarious, cup-shaped, regular or occasionally compressed, reaching a diameter of 2–3 cm. and a depth of 1 cm., externally rough, the roughenings consisting of conical tufts of hairs, yellowish to yellowish-brown; hymenium dark-brown to blackish; stem slightly swollen near the base and narrowed upward toward the base of the apothecium, yellowish, lighter than the outside of the apothecium; asci cylindric above, gradually tapering below, reaching a length of 300–325 μ and a diameter of 15 μ; spores ellipsoid, containing one large oil-drop which is nearly as large as the small diameter of the spore, hyaline, 12–15 $\times$ 20–23 μ; paraphyses strongly enlarged above where they reach a diameter of 8–10 μ.

On the ground.

TYPE LOCALITY: Europe.

DISTRIBUTION: Wisconsin; also in Europe.

ILLUSTRATIONS: Bull. Soc. Myc. Fr. **14**: *pl. 3, f. 1;* Boud. Ic. Myc. *pl. 242;* Ann. Myc. **2**: *pl. 12, f. 6–7.*

10. **Paxina Corium** (Weberb.) Seaver, comb. nov.

Peziza Corium Weberb. Pilze Norddeutsch. *pl. 3, f. 7.* 1873.
Aleuria Corium Gill. Champ. Fr. Discom. 39. 1879.
Lachnea Corium Phill. Brit. Discom. 204. 1887.
Macropodia Corium Sacc. Syll. Fung. **8**: 159. 1889.
Fuckelina Corium Kuntze, Rev. Gen. Pl. **2**: 852. 1891.
Sarcoscypha Corium Schröt. Krypt.-Fl. Schles. **3²**: 59. 1893.
Elvela nigra Peck, Bull. Torrey Club **26**: 70. 1899.
Cyathipodia Corium Boud. Hist. Class. Discom. Eu. 39. 1907.
Leptopodia Corium Boud. Ic. Myc. **6**: 1, 6. 1909.

Apothecia rather shallow cup-shaped or infundibuliform, with the margin more or less wavy, reaching a diameter of 3–5 cm. and a depth of 1–2 cm., externally blackish, with a brownish tomentum, the substance tough, becoming very hard when dry (white when broken); hymenium black or blackish; stem 2–3 cm. long and nearly 1 cm. in diameter at the base, tapering upward to a diameter of 3–4 mm. near the base of the apothecium then gradually expanding into the apothecium, irregularly and rather sparingly lacunose, tomentose; tomentum consisting of hairs which are composed of a chain of rather loosely united cells, often branched, pale-brown; asci cylindric above, reaching a diameter of 20 μ; spores 1-seriate, ellipsoid, smooth, usually containing one large oil-drop which nearly fills the spore, 10–12 $\times$ 18–20 μ; paraphyses strongly enlarged above where they reach a diameter of 7–9 μ, pale-brown.

On the ground.

TYPE LOCALITY: Europe.

DISTRIBUTION: Nebraska to Colorado and Idaho; also in Europe.

ILLUSTRATIONS: Weberb. Pilze Norddeutsch. *pl. 3, f. 7;* Cooke, Mycographia *pl. 48, f. 187;* Boud. Ic. Myc. *pl. 239* bis.

EXSICCATI: Fungi Columb. *2329;* Clements, Crypt. Form. Colo. *536.*

11. **Paxina nigrella** Seaver, sp. nov.

Apothecia scattered, shallow cup-shaped, with the margin more or less wavy, externally brownish-black, clothed with inconspicuous hairs, reaching a diameter of 2–3 cm. and a depth of about 1 cm. or less, becoming strongly wrinkled when dry (externally closely resembling *Pseudoplectania vogesiaca*); hymenium concave, black; stem reaching a length of 3–4 cm. and a diameter of 2–3 mm. of nearly uniform thickness throughout

its entire length, roundish or angular, clothed like the outside of the apothecium with brown hairs and attached at the base with a dense mass of coarse, black mycelium; hairs pale-brown, septate, thin-walled, flexuous, reaching a diameter of 10 μ; asci cylindric above, gradually tapering below, reaching a length of 500 μ and a diameter of 20 μ; spores 1-seriate, parallel with the ascus, ellipsoid, usually containing two small oil-drops and numerous granules, hyaline or faintly yellowish, 15 $\times$ 30–35 μ; paraphyses numerous slightly enlarged at their apices, dark-brown, reaching a diameter of 4–5 μ.

On rotten wood in coniferous woods.

TYPE LOCALITY: Tolland, Colorado.

DISTRIBUTION: Tolland and Caribou, Colorado.

This very characteristic species which externally very closely resembles *Pseudoplectania vogesiaca* is based on material collected June 30, 1914, by Dr. L. O. Overholts in Boulder Canyon, Colorado. A second collection was made by the late Professor E. Bethel at Caribou, Colorado, July 19, 1914. It differs from *Paxina Corium*, its closest ally, in the much larger size of its spores.

12. **Paxina semitosta** (Berk. & Curt.) Seaver, comb. nov. (PLATE 25, FIG. 2.)

Peziza semitosta Berk. & Curt. Grevillea **3**: 153. 1875.
Peziza Hainesii Ellis, Bull. Torrey Club **8**: 65. 1881.
Lachnea Hainesii Sacc. Syll. Fung. **8**: 186. 1889.
Fuckelina semitosta Kuntze, Rev. Gen. Pl. **2**: 852. 1891.
Scutellinia Hainesii Kuntze, Rev. Gen. Pl. **2**: 869. 1891.
Sepultaria gigantea Clements, Bot. Surv. Nebr. **5**: 8. 1901.
Sepultaria semitosta Morgan, Jour. Myc. **8**: 188. 1902.
Lachnea gigantea Sacc. & D. Sacc. in Sacc. Syll. Fung. **18**: 33. 1906.
Macropodia semitosta Durand, Jour. Myc. **12**: 31. 1906.

Apothecia gregarious or cespitose, rather deep cup-shaped, reaching a diameter of 2–5 cm. and a depth of 2–3 cm., externally brown and densely clothed with a soft tomentum; hymenium creamy-white when fresh, becoming brown when old or in drying; stem usually about 1 cm. long and 1–2 cm. in diameter, often immersed or partially immersed, deeply lacunose, brown and covered like the outside of the apothecium with tomentum; tomentum consisting of brown, rather thin-walled, sparingly septate, flexuous hairs which reach a diameter of 18–20 μ, of nearly uniform thickness throughout their entire length, ends

blunt, the length very variable; asci cylindric to clavate, reaching a length of 300–325 μ and a diameter of 15 μ; spores obliquely 1-seriate, with the ends overlapping, fusoid to fusiform, becoming sculptured when mature, containing two large oil-drops, 10–12 × 25–33 μ; spore-sculpturing consisting of coarse warts; paraphyses clavate, reaching a diameter of 7–8 μ at their apices.

On rich soil or more rarely on rotten wood.

TYPE LOCALITY: Pennsylvania.

DISTRIBUTION: New York to Wisconsin and Iowa.

ILLUSTRATIONS: Cooke, Mycographia *pl. 28, f. 109;* Grevillea **3**: *pl. 44, f. 225;* Jour. Linn. Soc. **31**: *pl. 16, f. 19.*

13. **Paxina fusicarpa** (Ger.) Seaver, comb. nov. (PLATE 25, FIG. 1.)

Peziza fusicarpa Ger. Bull. Torrey Club **4**: 64. 1873.
Peziza pubida Berk. & Curt. Grevillea **3**: 153. 1875.
Macropodia pubida Sacc. Syll. Fung. **8**: 159. 1889.
Lachnea fusicarpa Sacc. Syll. Fung. **8**: 172. 1889.
Fuckelina pubida Kuntzé, Rev. Gen. Pl. **2**: 852. 1891.
Scutellinia fusicarpa Kuntze, Rev. Gen. Pl. **2**: 869. 1891.
Sepultaria aspera Clements, Bot. Surv. Nebr. **4**: 12. 1895.
Lachnea aspera Sacc. & Sydow in Sacc. Syll. Fung. **14**: 757. 1899.
Peziza Morgani Massee; Morgan, Jour. Myc. **8**: 190. 1902.
Macropodia fusicarpa Durand, Jour. Myc. **12**: 29. 1906.

Apothecia gregarious or cespitose, rather deep cup-shaped, reaching a diameter of 2–3 cm. and a depth of 1 cm., externally brown and densely clothed with a soft tomentum; hymenium creamy-white when fresh, becoming brown when old or in drying; stem short, rarely exceeding 5 mm. in length and about 3–5 mm. in diameter, nearly even or in some cases deeply lacunose, clothed with tomentum similar to that on the outside of the apothecium, often immersed so that the apothecia appear to be sessile; tomentum consisting of thin-walled, flexuous, sparingly septate, brown hairs 12–15 μ in diameter and of nearly uniform thickness throughout their entire length, or occasionally enlarged at the base, slightly roughened on the outside, ends blunt; asci cylindric or clavate above, reaching a diameter of 20–22 μ and a length of 300–325 μ; spores 1-seriate, diagonally disposed and with the ends strongly overlapping, or becoming partially 2-seriate above, hyaline or slightly yellowish, fusiform, containing two large oil-drops, often slightly curved or unequal-sided, 10–12 × 30–45 μ, becoming sculptured; spore-sculpturing



1. PAXINA FUSICARPA
2. PAXINA SEMITOSTA

consisting of coarse warts; paraphyses cylindric or slightly clavate, reaching a diameter of 6–8 μ at their apices.

On soil in woods or more rarely on rotten wood.

TYPE LOCALITY: Poughkeepsie, New York.

DISTRIBUTION: Connecticut to Iowa and Virginia.

ILLUSTRATIONS: Cooke, Mycographia *pl. 28, f. 110* and *pl. 29, f. 113;* Grevillea **3**: *pl. 44, f. 226;* Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 21, f. 1.*

EXSICCATI: N. Am. Fungi *1269;* Fungi Columb. *1307;* Rab.-Winter, Fungi Eu. *325* (in part).

This and the preceding are frequently collected in midsummer.

14. **Paxina olivacea** (Clements) Seaver, comb. nov.

Phleboscyphus olivaceus Clements, Bull. Torrey Club **30**: 93. 1903.
Acetabula olivacea Sacc. & D. Sacc. in Sacc. Syll. Fung. **18**: 13. 1906.

Apothecia cespitose, cup-shaped, reaching a diameter of 4–5 cm. and a depth of 2–2.5 cm., margin becoming lacerate, externally brownish-olivaceous, furfuraceous-tuberculate; hymenium greenish-olivaceous; stem 1–2-sulcate or lacunose, furfuraceous, reaching a length of 2–2.5 cm. and a diameter of 1.5–2 cm.; asci cylindric, short-stipitate, reaching a length of 350 μ and a diameter of 14 μ; spores 1-seriate, ellipsoid, becoming sculptured, hyaline, 8 × 19 μ; spore-sculpturing consisting of minute warts; paraphyses slender, septate, hyaline, reaching a diameter of 3–4 μ at their apices.

On moist soil.

TYPE LOCALITY: Jack Brook, Minnehaha, Colorado.

DISTRIBUTION: Known only from the type locality.

DOUBTFUL SPECIES

Macropodia Schweinitzii Sacc. Syll. Fung. **8**: 160. 1889; *Peziza tomentosa* Schw. Trans. Am. Phil. Soc. II. **4**: 171. 1832; *Fuckelina Schweinitzii* Kuntze, Rev. Gen. Pl. **2**: 852. 1791. The identity of this species is uncertain.

Plectania rimosa Peck, Bull. Torrey Club **30**: 100. 1903. This may be *Paxina Corium.* No specimen has been seen.

35. **GEOPYXIS** (Pers.) Sacc. Syll. Fung. **8**: 63. 1889.

Peziza § *Geopyxis* Pers. Myc. Eu. **1**: 224. 1822.
Pustularia Fuckel, Symb. Myc. 328. 1869. Not *Pustularia* Roussel, 1806.

Apothecia medium sized, usually deep cup-shaped, externally pustulate, stipitate; stem usually slender and short; asci 8-

spored; spores ellipsoid, with or without oil-drops; paraphyses slender.

Type species, *Peziza carbonaria* Alb. & Schw.

Plants confined to burned places or rocky ledges.	1. *G. cupularis.*
Plants not confined to burned places.	
On damp soil; spores containing two conspicuous oil-drops.	
Apothecia bright-yellow, very coarsely verrucose.	2. *G. bronca.*
Apothecia dull-yellow, minutely verrucose.	3. *G. Catinus.*
Among mosses; spores without oil-drops.	4. *G. vulcanalis.*

1. **Geopyxis cupularis** (L.) Sacc. Syll. Fung. **8**: 72. 1889. (PLATE 26, FIG. 1.)

Peziza cupularis L. Sp. Pl. 1181. 1753.
Peziza carbonaria Alb. & Schw. Consp. Fung. 314. 1805.
Pustularia cupularis Fuckel, Symb. Myc. 328. 1869.
Peziza suburceolata Phill. Grevillea **7**: 21. 1878.
Aleuria cupularis Gill. Champ. Fr. Discom. 39. 1879.
Peziza crenata Bull. Herb. Fr. *pl. 396.* 1788.
Geopyxis carbonaria Sacc. Syll. Fung. **8**: 71. 1889.
Geopyxis suburceolata Sacc. Syll. Fung. **8**: 72. 1889.

Apothecia gregarious reaching a diameter of 3 mm. to 1 cm. (rarely larger) and of about the same depth, the mouth of the cup slightly constricted, at least in the younger specimens, and the margin crenate, the whole plant resembling a small acorn-cup, externally pustulate or nearly smooth, dull-yellowish; stem short, usually 2–3 mm. long and about 1 mm. thick, rather variable both in length and diameter, expanding rather abruptly into the apothecium; hymenium concave, a little paler than the outside of the apothecium; asci cylindric, reaching a length of 200 μ and a diameter of 10 μ; spores 1-seriate, obliquely arranged in the ascus when young and usually parallel when mature, ellipsoid, with the ends slightly narrowed, rather thick-walled, hyaline, smooth, about 9–10 $\times$ 18 μ, without conspicuous oil-drops; paraphyses enlarged above, reaching a diameter of 7 μ at their apices.

On the ground in burnt places.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to Washington, West Virginia and Colorado; also in Europe.

ILLUSTRATIONS: Boud. Ic. Myc. *pl. 338*, in part; Bull. Herb. Fr. *pl. 396, f. 3;* E. & P. Nat. Pfl. 1^1: 186, *f. 150, A.* & *B.;* Gill. Champ. Fr. Discom. *pl. 36;* Alb. & Schw. Consp. Fung. *pl. 4, f. 2.*

1. GEOPYXIS CUPULARIS
2. PEZIZA DOMICILIANA

EXSICCATI: N. Am. Fungi *2909*.

A specimen of *Peziza suburceolata* in the herbarium of The New York Botanical Garden collected by Dr. Harkness in California seems to be identical with the above.

2. **Geopyxis bronca** (Peck) Seaver, comb. nov.

Peziza bronca Peck, Ann. Rep. N. Y. State Mus. **29**: 54. 1878.
Aleuria humicola Boud. Hist. Class. Discom. Eu. 45. 1907.
Peziza humicola Sacc. & Trav. in Sacc. Syll. Fung. **22**: 614. 1913.

Apothecia gregarious or crowded, sessile or substipitate, at first subhemispheric, becoming more or less expanded with age, externally pale-yellow to bright lemon-yellow, the margin finely crenate and the entire exterior covered with coarse warts which give the apothecia a beautiful verrucose appearance, reaching a diameter of 2–3 cm. and of about half that depth; hymenium similar in color to the outside of the apothecium; asci cylindric or subcylindric, reaching a length of 400 μ and a diameter of 24–28 μ; spores 1-seriate, ellipsoid, hyaline, smooth, 12–14 $\times$ 20–24 μ, containing two conspicuous oil-drops; paraphyses rather strongly enlarged above.

On the ground.

TYPE LOCALITY: Knowersville, New York.

DISTRIBUTION: New York; also in Europe.

ILLUSTRATIONS: Ann. Rep. N. Y. State Mus. **29**: *pl. 2, f. 10–12;* Boud. Ic. Myc. *pl. 262.*

3. **Geopyxis Catinus** (Holmsk.) Sacc. Syll. Fung. **8**: 71. 1899.

Peziza Catinus Holmsk. Otia **2**: 22. 1799.
Pustularia Catinus Fuckel, Symb. Myc. 328. 1869.
Aleuria Catinus Gill Champ. Fr. Discom. 39. 1879.

Apothecia rather deep cup-shaped, externally dull-yellow, strongly but finely pustulate, the margin crenate and usually infolded, reaching a diameter of 2–3 cm. and nearly as deep; hymenium similar in color to the outside of the apothecium or a little lighter; stem reaching a length of 5 mm. to 1 cm. and usually 3–4 mm. thick, lacunose or nearly even; asci cylindric above, gradually tapering below, reaching a diameter of 15 μ; spores 1-seriate, broad-ellipsoid, rarely with the ends slightly narrowed, containing two large oil-drops, more or less granular, 10–12 $\times$ 20 μ; paraphyses about 2 μ in diameter, slightly enlarged above.

On the ground.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York and California; also in Europe.

ILLUSTRATIONS: Holmsk. Otia **2**: *pl. 8;* Boud. Ic. Myc. *pl. 336.*

4. **Geopyxis vulcanalis** (Peck) Sacc. Syll. Fung. **8**: 65. 1889.

?Peziza majalis Fries, Nova Acta Soc. Sci. Upsal. III. **1**: 120. 1855.
Peziza vulcanalis Peck; Coulter, Rep. U. S. Geol. Surv. Terr. **6**: 792. 1873.
Peziza pulchra Ger. Bull. Torrey Club **4**: 64. 1873.
Geopyxis pulchra Sacc. Syll. Fung. **8**: 65. 1889.
Phleboscyphus radicatus Clements, Bull. Torrey Club **30**: 94. 1903.
Sarcoscypha dawsonensis Peck, Bull. Torrey Club **33**: 220. 1906.
Acetabula radicata Sacc. Syll. Fung. **18**: 13. 1906.

Apothecia gregarious, short-stipitate, subinfundibuliform with the margin usually incurved and crenate, externally covered with an evanescent, pruinose or subtomentose coat, especially when young, often becoming entirely smooth with age, whitish or yellowish-white, darker when naked, reaching a diameter of 1–2 cm. and about half as deep; stem 1–2 mm. thick and of variable length up to 5 mm. but often so short that the apothecia appear sessile; hymenium pale-orange; asci cylindric above, tapering below into a long stem-like base, reaching a length of 275 μ and a diameter of 15–17 μ at the broadest point; spores ellipsoid, with the ends strongly narrowed, 8–10 $\times$ 14–18 μ; paraphyses slightly thickened above, reaching a diameter of 4 μ at their apices.

Nestling among mosses and leaf-mould in coniferous woods.

TYPE LOCALITY: Snake River, Wyoming.

DISTRIBUTION: New York to Wyoming and Colorado.

ILLUSTRATION: Cooke, Mycographia *pl. 74, f. 285.*

EXSICCATI: Clements, Crypt. Form. Colo. *126* (as *Phleboscyphus radicatus* Clements).

DOUBTFUL AND EXCLUDED SPECIES

Geopyxis nebulosa (Cooke) Sacc. Syll. Fung. **8**: 70. 1889; *Peziza nebulosa* Cooke, Mycographia 163. 1877. This is an inoperculate species and should be referred to the genus *Ciboria.*

Geopyxis nebulosoides Peck, Bull. Torrey Club **32**: 80. 1905. Probably not distinct from *Geopyxis nebulosa.*

Geopyxis cinerascens (Rehm) Sacc. Syll. Fung. **18**: 14. 1906; *Tarzetta cinerascens* Rehm, Ann. Myc. **2**: 352. 1904. This is also undoubtedly a *Ciboria.*

Geopyxis verruculosa (Berk. & Curt.) Sacc. Syll. Fung. **8**: 68. 1889;

Peziza verruculosa Berk. & Curt. Proc. Am. Acad. **4**: 127. 1860. This may be a *Pseudoplectania*.

Geopyxis Hicksii Peck, Ann. Rep. N. Y. State Mus. **46**: 38. 1893. This is probably a *Sclerotinia*. Type could not be found in the Peck collection at Albany.

36. **RHIZINA** Fries, Obs. Myc. **1**: 161. 1815.

Apothecia expanded, attached to the substratum by numerous root-like processes, the substance fleshy or subfleshy, rather tough when dry; asci cylindric to clavate, 8-spored; spores fusoid; paraphyses stout.

Type species, *Rhizina undulata* Fries.

1. **Rhizina inflata** (Schaeff.) Karst. Act. Fauna Fl. Fenn. **2**: 112. 1885. (PLATE 27.)

Elvela inflata Schaeff. Fung. Bavar. **4**: 102. 1774.
Elvela acaulis Pers. Obs. Myc. **2**: 29. 1799.
Phallus acaulis Batsch, Elench. Fung. 130. 1783.
Peziza rhizophora Willd. Fl. Berol. 402. 1787.
Octospora rhizophora Hedw. Descr. **2**: 15. 1788.
Rhizina undulata Fries, Obs. Myc. **1**: 161. 1815.
Rhizina laevigata Fries, Obs. Myc. **1**: 162. 1815.

Apothecia sessile, discoid or nearly so, attached to the substratum by root-like processes which cover the lower surface to near the margin, the substance firm, more or less spongy, becoming tough or rather corky when dry, reaching a diameter of 5–6 cm.; hymenium plane or slightly convex, even or undulated, brownish-black, the margin lighter; asci cylindric to clavate, reaching a length of about 300 μ and a diameter of 12–18 μ; spores 1-seriate or partially 2-seriate above, with the ends overlapping, fusiform, usually with an apiculus 3–4 μ long on either end, containing two oil-drops and occasionally with a pseudoseptate appearance, hyaline, about 10 $\times$ 35–43 μ; paraphyses enlarged and clinging together at their apices, brown, intermixed with brown, thick-walled, non-septate, hair-like structures.

Rare on the ground where it has recently been burned over.

TYPE LOCALITY: Europe.

DISTRIBUTION: Connecticut to Alabama and Montana; also in Europe.

ILLUSTRATIONS: Boud. Ic. Myc. *pl. 251;* E. & P. Nat. Pfl. 1^1: 172, *f. 143;* Gill. Champ. Fr. Discom. *pl. 30, f. 1;* Hedw.

Descr. *pl. 5, A;* Phill. Brit. Discom. *pl. 2, f. 10;* Rab. Krypt.-Fl. 1^3: *1136, f. 1–4;* Schaeff. Fung. Bavar. *pl. 153.*

EXSICCATI: N. Am. Fungi *2739.*

DOUBTFUL SPECIES

Rhizina spongiosa Berk. & Curt.; Berk. Jour. Linn. Soc. **10**: 364. 1869. Described as one and one-half inches in diameter with spores 20 × 60 μ. Growing vertically on bushes.

37. **DISCINA** Fries, Summa Veg. Scand. 348. 1849.

Apothecia medium to large, sessile or short-stipitate, fleshy or waxy, light- or dark-colored; asci cylindric or subcylindric, very long, usually showing a tendency to become spirally twisted; spores ellipsoid, appendiculate and often sculptured, hyaline or faintly colored; spore appendages apiculate or truncate; paraphyses rather stout.

Type species, *Discina perlata* Fries.

Apothecia dark-colored; spore appendages apiculate.
- Apothecia large, 6–7 cm. or more in diameter. 1. *D. ancilis.*
- Apothecia medium sized, not exceeding 5 cm. in diameter. 2. *D. apiculata.*

Apothecia bright-colored; spore appendages truncate.
- Hymenium strongly convolute. 3. *D. convoluta.*
- Hymenium even or only slightly undulated. 4. *D. leucoxantha.*

1. **Discina ancilis** (Pers.) Sacc. Syll. Fung. **8**: 103. 1899. (PLATE 28, FIG. 2.)

Peziza ancilis Pers. Myc. Eu. **1**: 219. 1822.
Peziza perlata Fries, Syst. Myc. **2**: 43. 1822.
Discina perlata Fries, Summa Veg. Scand. 348. 1849.
Rhizina helvetica Fuckel, Symb. Myc. Nacht. **2**: 66. 1873.
Peziza Warnei Peck, Ann. Rep. N. Y. State Mus. **30**: 59. 1878.
Aleuria ancilis Gill. Champ. Fr. Discom. 36. 1879.
Acetabula ancilis Lamb. Fl. Myc. Belg. **2**: 573. 1880.
Discina Warnei Sacc. Syll. Fung. **8**: 102. 1889.
Discina ancilis Sacc. Syll. Fung. **8**: 103. 1889.
Discina helvetica Sacc. Syll. Fung. **8**: 103. 1889.

Apothecia gregarious or scattered, more rarely congested, short-stipitate, at first subglobose, soon becoming discoid, finally repand, at first regular in form, becoming irregular and often angular as the margin rolls back, externally whitish or pallid, reaching a diameter of 7–8 cm., or in rare cases as large as 20–25 cm.; hymenium uneven, often beautifully veined or convolute, plane or convex, usually umbilicate, dark-brown, finally

RHIZINA INFLATA

almost black; stem very short and stout, often 1–3 cm. in diameter and rarely exceeding 1 cm. in length, or entirely wanting, more or less lacunose, whitish or overcast with a pinkish tint; asci cylindric or subcylindric, reaching a length of 300–350 μ and a diameter of 12–18 μ, 8-spored; spores obliquely 1-seriate, very large, ellipsoid, hyaline, 12–14 × 30–35 μ, or occasionally as long as 40 μ including apiculi; at first smooth, becoming sculptured; spore-sculpturing consisting of minute warts; spore appendages consisting of a minute apiculus 4–5 μ long and 3–4 μ broad at the base, one at either end of the spore; paraphyses strongly enlarged above, closely adhering together, dark yellowish-brown reaching a diameter of 8 μ.

On the ground in coniferous woods, more rarely on rotten wood.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to Manitoba, Washington, Oregon, and Colorado; also in Europe.

ILLUSTRATIONS: Ann. Rep. N. Y. State Mus. **30**: *pl. 1, f. 19–21;* Boud. Ic. Myc. *pl. 252;* Fuckel, Symb. Myc. Nacht. **2**: *f. 24;* Pat. Tab. Fung. *f. 596;* Cooke, Mycographia *pl. 103, f. 371;* Rab. Krypt.-Fl. 1^3: 922, *f. 1–4.*

EXSICCATI: Shear, New York Fungi *324;* N. Am. Fungi *2622.*

Rather frequently collected but scarcely common.

2. **Discina apiculata** (Cooke) Seaver, Mycologia **13**: 70. 1921.

Peziza apiculata Cooke, Mycographia 175. 1877.
Phaeopezia apiculata Sacc. Bot. Centr. **18**: 218. 1884.
Aleuria apiculata Boud. Hist. Class. Discom. Eu. 47. 1907.
?Peziza elaeodes Clements, Bot. Surv. Nebr. **5**: 6. 1901.

Apothecia scattered, sessile, at first cup-shaped, soon becoming discoid, fleshy, circular in outline not usually exceeding 1.5 cm. in diameter; hymenium dark brownish-black, slightly concave or nearly plane; asci cylindric or subcylindric, reaching a length of 200 μ and a diameter of 18 μ, 8-spored; spores obliquely 1-seriate, with the ends overlapping, narrow-ellipsoid to fusoid, often with a small apiculus at either end, becoming pale-brownish and often delicately sculptured, about 10 × 24 μ, containing two large oil-drops; spore-sculpturing consisting of minute warts; paraphyses slender, adhering together at their apices, dark-brown.

On damp soil.

TYPE LOCALITY: Italy.

DISTRIBUTION: (Nebraska?) and (Bermuda?); also in Europe.
ILLUSTRATION: Cooke, Mycographia *pl. 79, f. 305.*

A small specimen collected in Bermuda by B. O. Dodge seems to belong here so far as spore characters are concerned. However, the material was so limited that it was impossible to make a careful study of the gross characters. Also *Peziza elaeodes* of Clements seems to fit Cooke's description but here again it was impossible to be certain for lack of material. So the species must remain doubtful so far as its record from North America is concerned.

3. **Discina convoluta** Seaver, Mycologia **13**: 70. 1921. (PLATE 28, FIG. 1.)

Apothecia gregarious or cespitose, very short-stipitate, becoming shallow cup-shaped or subdiscoid, externally whitish at the base, becoming yellowish upwards toward the margin, reaching a diameter of 6 cm.; hymenium yellowish-brown, very deeply convolute, the convolutions consisting of more or less radiating ribs or veins, resembling those of *Peziza venosa* but more distinct; stem very short and stout, 1 cm. or more thick and usually not more than 1 cm. long, whitish, deeply corrugated; asci cylindric or subcylindric, reaching a length of 400–500 μ and a diameter of 20 μ, 8-spored but a part of the spores often remaining undeveloped; spores 1-seriate, with the ends usually overlapping, ellipsoid, becoming sculptured and appendiculate 12–14 $\times$ 35–40 μ; spore-sculpturing consisting of warts or short interrupted ridges which often approach very fine reticulations; spore appendages consisting of a cup-like structure at each end; paraphyses stout, usually straight, reaching a diameter of 8 μ at their apices, densely filled with yellow granules.

On the ground in woods.

Type collected in the woods near Yonkers, New York, May 22, 1916, by F. J. Seaver.

DISTRIBUTION: Known only from the type locality.

4. **Discina leucoxantha** Bres. Rev. Myc. **4**: 212. 1882.

Peziza leucoxantha Bres. Fungi Trid. **1**: 42. 1883.

Apothecia gregarious or substipitate, at first subglobose, expanding and becoming hemispheric or nearly plane, externally whitish, reaching a diameter of 4–7 cm., the margin regular or lobed; hymenium concave or nearly plane, even or undulated,



1. DISCINA CONVOLUTA
2. DISCINA ANCILIS

bright-yellow or becoming yellowish-brown with age; stem about 1 cm. in diameter and scarcely exceeding 5 mm. in length, irregularly corrugated at the base, the flesh thick and brittle; asci cylindric or subcylindric, attenuated at the base, reaching a length of 400 μ and a diameter of 20 μ, becoming twisted, 8-spored; spores ellipsoid, smooth, containing one large oil-drop and several smaller ones, becoming minutely sculptured, 10–15 $\times$ 27–35 μ; spore-sculpturing consisting of minute warts or occasionally minute ridges or indistinct reticulations; spore appendages consisting of truncate protuberances, one at either end of the spore; paraphyses slender, septate, branched, slightly enlarged above, filled with orange granules.

Rare on the ground in coniferous woods.

TYPE LOCALITY: France.

DISTRIBUTION: New York to Maryland; also in Europe.

ILLUSTRATIONS: Bres. Fungi Trid. *pl. 44;* Bull. Soc. Myc. Fr. **3**: *pl. 12;* Boud. Ic. Myc. *pl. 253;* Rab. Krypt.-Fl. 1^3: 922, *f. 5.*

38. **PEZIZA** (Dill.) L. Sp. Pl. 1180. 1753.

Plicaria Fuckel, Symb. Myc. 325. 1869 (in part).
Pustularia Fuckel, Symb. Myc. 328. 1869 (in part).
Otidea Fuckel, Symb. Myc. 329. 1869 (in part).
Peziza § *Galactinia* Cooke, Mycographia 253. 1879.
Galactinia Sacc. Syll. Fung. **8**: 106. 1889.
Aleuria Authors, not Fuckel Symb. Myc. 325. 1869.
Heteroplegma Clements, Bull. Torrey Club **30**: 92. 1903.
Disciotis Boud. Hist. Class. Discom. Eu. 42. 1907.
Pseudotis Boud. Hist. Class. Discom. Eu. 52. 1907.
Pachyella Boud. Hist. Class. Discom. Eu. 50. 1907.

Apothecia medium sized or large, usually not less than 1 cm. in diameter and occasionally reaching a diameter of 15–20 cm., usually cup-shaped or occasionally discoid or repand, the color variable; asci cylindric to clavate, usually 8-spored; spores, ellipsoid, hyaline or faintly colored, smooth or sculptured; spore-sculpturing consisting of minute papillae to rather coarse warts or short interrupted ridges, never truly reticulate; paraphyses variable.

Type species, *Peziza cochleata* L.

Spores at maturity sculptured; spore-sculpturing varying from minute papillae to coarse warts.
Hymenium becoming brown, brownish-black or olivaceous.

Apothecia also dark-colored, the color being similar to that of the hymenium.
Apothecia large, reaching a diameter of 8–10 cm. 1. *P. badia.*
Apothecia small, not exceeding 1 or 2 cm. in diameter.
Spores 10–11 × 20–22 μ. 2. *P. brunneoatra.*
Spores 12–14 × 25–27 μ. 3. *P. spissa.*
Apothecia externally light-colored, white, whitish, golden-yellow, or occasionally dingy with age.
Flesh when broken turning golden-yellow. 4. *P. succosa.*
Flesh when broken not turning golden-yellow.
Hymenium ochraceous-brown.
Apothecia externally pustulate; on burnt ground. 5. *P. pustulata.*
Not strongly pustulate; on wood. 6. *P. Emileia.*
Hymenium nearly black.
Spores 8–9 × 17–21 μ. 7. *P. melaleuca.*
Spores 10 × 14 μ. 8. *P. melaleucoides.*
Hymenium violaceous, becoming very dark with age; occurring on burnt ground. 9. *P. violacea.*
Hymenium white or whitish, more rarely rosaceous or faintly violet.
Hymenium rosaceous. 10. *P. griseorosea.*
Hymenium white, becoming smoky or faintly violet; on burnt ground. 11. *P. proteana.*
Spores remaining permanently smooth.
Apothecia externally reddish-brown, similar in color to the hymenium.
Apothecia marked with concentric rings, sessile. 12. *P. concentrica.*
Apothecia not marked with concentric rings, substipitate. 13. *P. abietina.*
Apothecia externally white or pallid, rarely dingy with age.
Apothecia cup-shaped or repand, not adhering to the substratum except at the central point of attachment.
Hymenium becoming venose-reticulate; spores very large, 12–17 × 22–30 μ. 14. *P. venosa.*
Hymenium not becoming venose-reticulate; spores rarely over 20 μ long.
Apothecia becoming repand at maturity.
Occurring about caves and cellars; flesh often turning yellow when broken. 15. *P. domiciliana.*
Occurring on rotten wood; flesh not turning yellow when broken. 16. *P. repanda.*
Apothecia normally remaining cup-shaped.

Occurring on manure or heavily fertilized soil.
Apothecia large; spores 10–11 × 20–23 μ. 17. *P. vesiculosa.*
Apothecia small; spores 8 × 16 μ. 18. *P. fimeti.*
Occurring on humus in woods. 19. *P. sylvestris.*
Apothecia becoming discoid, closely adhering to the substratum. 20. *P. clypeata.*

1. **Peziza badia** Pers. Obs. Myc. **2**: 78. 1799.

?Peziza cochleata L. Sp. Pl. 1181. 1753.
?Elvela cochleata Bolton, Fungi Halifax **3**: 99 (in part). 1789.
Scodellina badia S. F. Gray, Nat. Arrang. Brit. Pl. **1**: 669. 1821.
Plicaria badia Fuckel, Symb. Myc. 327. 1869.
Aleuria badia Gill. Champ. Fr. Discom. 43. 1879.
Galactinia badia Boud. Hist. Class. Discom. Eu. 48. 1907.

Apothecia scattered, gregarious, or more commonly cespitose, sessile, at first globose, expanding and becoming deep cup-shaped, regular in form, or infolded and cochleate or auricular, occasionally one-sided, rarely *Otidea*-like, externally varying from tan-colored when young to dark-brown with age, whitish near the base, pustulate, the pustules reddish or reddish-purple, becoming dark with age, reaching a diameter of 10 cm.; hymenium usually concave, dark-brown; asci tapering below and often forked at the base, cylindric above; spores 1-seriate, usually obliquely arranged and often irregularly crowded, ellipsoid, becoming sculptured, hyaline or faintly colored, 8–10 × 17–23 μ; spore-sculpturing consisting of rather coarse warts or short interrupted ridges; paraphyses rather strongly enlarged above, yellowish.

On the ground in deciduous woods and open places.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to Oregon, California and Alabama; also in Europe. Probably widely distributed.

ILLUSTRATIONS: Bolton, Fungi Halifax, *pl. 99;* Boud. Ic. Myc. *pl. 283;* Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 14, f. 2; pl. 15, f. 1;* Cooke, Mycographia *pl. 57, f. 226;* Gill. Champ. Fr. Discom. *pl. 42;* Mycologia **7**: *pl. 155* (lower figure); (?) Sow. Engl. Fungi *pl. 5.*

EXSICCATI: N. Am. Fungi *981.*

The species is rather common and often abundant.

2. **Peziza brunneoatra** Desm. Ann. Sci. Nat. II. 6: 244. 1836.

?Peziza macrospora Wallr. Fl. Crypt. Germ. **2**: 500. 1833.
?Humaria macrospora Fuckel, Symb. Myc. 323. 1869.
Aleuria brunneoatra Gill. Champ. Fr. Discom. 53. 1879.
Plicaria brunneoatra Rehm in Rab. Krypt.-Fl. 1^3: 1010. 1896.
?Phaeopezia vinacea Clements, Bot. Surv. Nebr. **4**: 16. 1896.
?Aleurina vinacea Sacc. & Sydow in Sacc. Syll. Fung. **16**: 740. 1902.
?Peziza nana Massee & Morgan, Jour. Myc. **8**: 190. 1902.
Galactinia brunneoatra Boud. Hist. Class. Discom. Eu. 49. 1907.

Apothecia scattered or gregarious, sessile, expanding, and at maturity becoming scutellate to discoid, not usually exceeding 1–2 cm. in diameter; hymenium at first concave, becoming plane, finally a little convex, often umbilicate, brownish-black with a slight tinge of green; asci cylindric above, tapering rather abruptly below into a stem-like base, reaching a length of 300–350 μ and a diameter of 15–17 μ, 8-spored; spores 1-seriate or occasionally slightly crowded, parallel with the ascus or oblique, at first smooth, becoming sculptured, very faintly yellowish or olive, containing two conspicuous oil-drops or more rarely one, 10–11 $\times$ 20–22 μ; spore-sculpturing taking the form of rather conspicuous warts about 2 μ in diameter; paraphyses rather strongly enlarged above, reaching a diameter of 7–8 μ at their apices, yellowish-brown.

On the ground in damp places.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to Iowa; also in Europe.

ILLUSTRATIONS: Boud. Ic. Myc. *pl. 298;* Cooke, Mycographia *pl. 20, f. 78;* Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 14, f. 3.*

3. **Peziza spissa** Berk. Grevillea **3**: 152. 1875.

Humaria spissa Sacc. Syll. Fung. **8**: 141. 1889.

Apothecia gregarious, at first shallow cup-shaped, becoming discoid, attached to the substratum by a short stem-like base which is obscured at maturity, reaching a diameter of 1–2 cm.; hymenium concave, becoming convex, dark-brown; asci cylindric or subcylindric above, tapering below into a stem-like base, becoming spirally twisted when young, reaching a length of 260–275 μ and a diameter of 23–24 μ; spores 1-seriate, obliquely or irregularly disposed, ellipsoid, usually containing two large oil-drops and often several smaller ones, granular within and slightly yellowish, smooth, then becoming very minutely sculp-

tured, 12–14 × 25–27 μ; spore-sculpturing consisting of minute warts; paraphyses stout, strongly enlarged above where they reach a diameter of 10–12 μ, filled with a mass of dark-brown coloring matter which often separates in such a way as to give the tip of the paraphyses a septate or muriform appearance, hyaline below, often spirally twisted like the ascus.

On damp soil.

TYPE LOCALITY: Alabama.

DISTRIBUTION: Alabama and Washington.

ILLUSTRATION: Cooke, Mycographia *pl. 20, f. 79.*

Except in habitat and mode of attachment, resembling a *Psilopezia.*

4. **Peziza succosa** Berk. Ann. Mag. Nat. Hist. I. **6**: 358. 1841.

Aleuria succosa Gill. Champ. Fr. Discom. 45. 1879.
Otidea succosa Thüm. Mycoth. Univ. *1411.* 1879.
Galactinia succosa Sacc. Syll. Fung. **8**: 106. 1889.
Plicaria succosa Rehm in Rab. Krypt.-Fl. **1**[3]: 1016. 1896.

Apothecia gregarious or scattered, sessile, at first hemispheric, expanding and becoming shallow cup-shaped or more rarely subdiscoid, regular in outline or irregularly contorted, the margin even or occasionally more or less lobed, externally whitish or yellowish with a slight shade of olive, reaching a diameter of 3–4 cm.; hymenium concave or occasionally nearly plane, even, brown, becoming darker with age, often with an olivaceous tint apparently from the spores which have dusted out over its surface; asci cylindric or subcylindric, reaching a length of 200–225 μ and a diameter of 12–15 μ, 8-spored; spores obliquely 1-seriate, ellipsoid, usually containing two distinct oil-drops, hyaline or very faintly yellowish (olivaceous in mass), finally becoming sculptured, 8–12 × 16–20 μ; spore-sculpturing consisting of rather conspicuous warts 1–2 μ in diameter; paraphyses slender, septate, enlarged above, filled with yellowish coloring matter.

On damp soil in woods.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to Iowa and Wisconsin; also in Europe.

ILLUSTRATIONS: Ann. Mag. Nat. Hist. I. **6**: *pl. 10, f. 5–7;* Gill. Champ. Fr. Discom. *pl. 46;* Boud. Ic. Myc. *pl. 284;* Cooke, Mycographia *pl. 63, f. 243;* Pat. Tab. Fung. *f. 166;* Phill. Brit.

Discom. *pl. 4, f. 16;* Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 16, f. 2.*

Quite frequently found in midsummer.

5. **Peziza pustulata** (Hedw.) Pers. Syn. Fung. 646. 1801. (PLATE 29, FIG. 1.)

Octospora pustulata Hedw. Descr. **2**: 19. 1787.
Scodellina pustulata S. F. Gray, Nat. Arrang. Brit. Pl. **1**: 668. 1821.
Plicaria pustulata Fuckel, Symb. Myc. 327. 1869.
Peziza assimilata Karst. Not. Fauna Fl. Fenn. **10**: 113. 1869.
?Peziza echinosperma Karst. Not. Fauna Fl. Fenn. **10**: 115. 1869.
Peziza Petersii Berk. Grevillea **3**: 150. 1875.
Aleuria pustulata Gill. Champ. Fr. Discom. 45. 1879.
Peziza umbrina Boud.; Cooke, Mycographia 226 (in part). 1879.
Aleuria umbrina Gill. Champ. Fr. Discom. 42. 1879.
Galactinia pustulata Boud. Hist. Class. Discom. Eu. 48. 1907.

Apothecia gregarious, scattered, or cespitose, at first closed and globose, gradually expanding, reaching a diameter of 3–5 cm., regular, or much contorted, externally whitish and densely pustulate, the pustules giving rise to bran-like particles as the plant matures, becoming dingy with age, margin usually crenate; hymenium pale- to dark-brown; asci cylindric above, reaching a length of 275 μ and a diameter of 12–14 μ; spores 1-seriate, ellipsoid, becoming minutely sculptured, hyaline to faintly yellowish, 8–10 $\times$ 15–17 μ; spore-sculpturing consisting of very minute warts; paraphyses strongly enlarged above and reaching a diameter of 7–8 μ at their apices.

On charcoal and burned areas.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to Wisconsin and Wyoming; also in Europe.

ILLUSTRATIONS: Boud. Ic. Myc. *pl. 279;* Cooke, Mycographia *pl. 64, f. 246, pl. 106, f. 378;* Grevillea **2**: *pl. 24, f. 2;* Hedw. Descr. **2**: *pl. 6, f. 1–4;* Gill. Champ. Fr. Discom. *pl. 47, f. 2;* Mycologia **7**: *pl. 156* (upper figure).

The species is quite common in midsummer.

6. **Peziza Emileia** Cooke, Mycographia 226. 1879.

Aleuria Emileia Boud. Hist. Class. Discom. Eu. 46. 1907.

Apothecia gregarious or scattered, sessile, large, at first cup-shaped, becoming expanded and often irregularly repand, the margin even or more or less lobed, regular in form or contorted, externally white or whitish-farinose, reaching a diameter of



1. PEZIZA PUSTULATA
2. PEZIZA VESICULOSA

8–10 cm.; hymenium concave, plane or occasionally convex, even or undulated; asci cylindric or subcylindric, reaching a length of 200 μ and a diameter of 12–15 μ; spores 1-seriate, ellipsoid, hyaline, becoming sculptured, 8 × 16–18 μ; spore-sculpturing consisting of very minute warts or papillae; paraphyses slender, septate, strongly enlarged above.

On damp soil in woods.

TYPE LOCALITY: France.

DISTRIBUTION: Wisconsin; also in Europe.

ILLUSTRATIONS: Cooke, Mycographia *pl. 106, f. 379;* Boud. Ic. Myc. *pl. 280.*

7. **Peziza melaleuca** (Bres.) Seaver, comb. nov.

Discina melaleuca Bres. Fungi Trid. **2**: 74. 1898.
?Heteroplegma caeruleum Clements, Bull. Torrey Club **30**: 92. 1903.
?Peziza caerulea Sacc. & D. Sacc. in Sacc. Syll. Fung. **18**: 20. 1906.

Apothecia gregarious, soon becoming expanded, finally often repand or with the extreme margin slightly elevated, the center usually depressed or umbilicate, externally white or whitish, reaching a diameter of 2–4 cm., narrowed below into a stem-like base; hymenium concave, becoming plane or slightly convex, brownish-black, when dry entirely black; asci cylindric or subcylindric, reaching a length of 360–400 μ and a diameter of 16–18 μ; spores 1-seriate, ellipsoid, containing 1–3 (usually 2) oil-drops, 8–9 × 17–21 μ, becoming sculptured; spore-sculpturing consisting of papillae or minute warts; paraphyses rather stout, enlarged above where they reach a diameter of 6–8 μ.

On the ground in coniferous woods.

TYPE LOCALITY: Europe.

DISTRIBUTION: Colorado; also in Europe.

ILLUSTRATION: Bres. Fungi Trid. *pl. 186.*

EXSICCATI: Clements, Crypt. Form. Colo. *124?* (as *Plicaria Fuckelii caerulescens*).

8. **Peziza melaleucoides** Seaver, sp. nov.

Apothecia sessile or short-stipitate, reaching a diameter of 10 cm., cup-shaped, becoming somewhat repand, usually whitish near the base, brownish above or occasionally with a lavender tint; stem when present lacunose, very short; hymenium concave, often umbilicate, becoming dark-brown or tinted with olive; asci cylindric or subcylindric, reaching a length of 250 μ and a diameter of 12–15 μ; spores 1-seriate, ellipsoid, each containing

two oil-drops, 10 × 14 μ, hyaline or subhyaline, becoming sculptured; spore-sculpturing consisting of minute papillae; paraphyses about 4 μ in diameter below, enlarged above where they reach a diameter of 8 μ, filled with brown coloring matter.

On the ground or more often on rotten wood in coniferous woods.

Type collected by Dr. L. O. Overholts at Tolland, Colorado, June, 1914.

DISTRIBUTION: Colorado.

This species differs from the preceding, which it resembles externally, in the broad-ellipsoid spores.

9. **Peziza violacea** Pers. Syn. Fung. 639. 1801. (PLATE 30, FIG. 2.)

Peziza Boltoni Quél. Bull. Soc. Bot. Fr. **25**: 290. 1878.
Aleuria violacea Gill. Champ. Fr. Discom. 47. 1879.
Peziza ampelina Quél. Grevillea **8**: 116. 1880.
Humaria violacea Sacc. Syll. Fung. **8**: 149. 1889.
Aleuria Boltoni Gill. Champ. Fr. Discom. 206. 1886.
Galactinia Boltoni Boud. Hist. Class. Discom. Eu. 48. 1907.

Apothecia gregarious, sessile or substipitate when young, at first closed and subglobose, gradually expanding and becoming shallow cup-shaped, discoid, or occasionally repand, at first regular in form, becoming irregular, the margin often splitting, externally at first white, finally becoming pale-violaceous, reaching a diameter of 3–4 cm., although often much smaller; hymenium concave, plane or convex, pale-violet when young, becoming deep-violet with age, finally almost black, even or slightly undulated, usually umbilicate; asci cylindric or subcylindric, gradually attenuated below, reaching a length of 200–250 μ and a diameter of 12–15 μ, 8-spored; spores 1-seriate, or irregularly disposed, ellipsoid, usually containing two small oil-drops, hyaline, becoming sculptured, 8 × 12–13 μ; spore-sculpturing consisting of very minute warts; paraphyses slender, septate, enlarged above, usually curved and filled with violet granules, reaching a diameter of 8 μ at their apices.

On burnt ground and charcoal beds.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to Wisconsin and Colorado; also in Europe.

ILLUSTRATIONS: Grevillea **8**: *pl. 131, f. 4;* Boud. Ic. Myc. *pl. 276;* Gill. Champ. Fr. Discom. (suites).



1. PEZIZA PROTEANA
2. PEZIZA VIOLACEA

Quite common on burned places and rather attractive because of its violet color, although not very conspicuous.

10. **Peziza griseorosea** Ger. Bull. Torrey Club **6**: 31. 1875.

Apothecia shallow cup-shaped, externally whitish-pruinose, sessile, the flesh rather thin, the color grayish-ochraceous; reaching a diameter of 2–4 cm.; hymenium concave, pale-rosaceous to subochraceous; asci cylindric or subcylindric; spores 1-seriate, ellipsoid, becoming sculptured, 7–10 × 15–18 μ, hyaline or slightly yellowish; spore-sculptured taking the form of minute warts; paraphyses slightly enlarged at their apices.

On the ground.

TYPE LOCALITY: Poughkeepsie, New York.

DISTRIBUTION: New York and Connecticut to Wisconsin and Virginia.

ILLUSTRATION: Cooke, Mycographia *pl. 65, f. 249.*

An authentic specimen of this species from the herbarium of Gerard is deposited in our collection at The New York Botanical Garden. A number of specimens from various localities have been referred to this species but always with more or less doubt, since it is difficult to determine the exact color from the dried specimens and the species was originally distinguished chiefly on that character. When possible, color records should be preserved in such cases as this in connection with field records.

11. **Peziza proteana** (Boud.) Seaver, Mycologia **9**: 1. 1917. (PLATE 30, FIG. 1.)

Aleuria proteana Boud. Bull. Soc. Myc. Fr. **15**: 50. 1899.
Galactinia proteana Sacc. Syll. Fung. **16**: 709. 1902.

Apothecia sessile, cup-shaped, finally becoming expanded and often more or less repand, entirely white when young, becoming overcast with a faint reddish or lilac tint, reaching a diameter of 3–6 cm.; hymenium concave, becoming plane or convex and usually umbilicate, the color varying from white when young to rosy, pale-lilac or slightly brownish, becoming darker with age; asci cylindric or subcylindric, reaching a length of 225–250 μ and a diameter of 10 μ; spores 1-seriate or becoming slightly crowded, small, ellipsoid, usually containing two small oil-drops, at first smooth, becoming sculptured, 5–7 × 12–13 μ; spore-sculpturing assuming the form of minute warts or papillae; paraphyses slender, septate, enlarged above where they reach a diameter of 7–8 μ.

On old burnt places which have been overrun with mosses.

TYPE LOCALITY: France.

DISTRIBUTION: New York and Texas; also in Europe.

ILLUSTRATIONS: Bull. Soc. Myc. Fr. **15**: *pl. 3, f. 1;* Boud. Ic. Myc. *pl. 293;* Mycologia **9**: *pl. 1* (upper figure).

In habitat and spore characters this species resembles *Peziza violacea* but differs much in color.

12. **Peziza concentrica** Seaver, sp. nov.

Apothecia subglobose, becoming more or less widely expanded, externally pale-brownish, reaching a diameter of 2–4 cm., both externally and internally marked with concentric rings consisting of depressions and elevations which somewhat resemble ripple-marks; hymenium concave or nearly plane, showing the same concentric markings found on the exterior of the apothecium, similar in color to the outside of the apothecium or a little darker, often slightly olivaceous; asci cylindric above, tapering below rather abruptly into a stem-like base with an enlargement at the extreme base, reaching a length of 300 μ and a diameter of 16 μ; spores 1-seriate, ellipsoid, hyaline or subhyaline, remaining smooth, 10 $\times$ 18–20 μ; paraphyses reaching a diameter of 2 μ below and a diameter of 4 μ at their apices, yellowish-olive.

On the ground.

TYPE LOCALITY: Humboldt County, California.

DISTRIBUTION: California.

This species is based on material submitted to the writer by the late C. G. Lloyd. The specimens were collected by Mrs. H. E. Wilder, January 22, 1914, in the locality indicated above.

13. **Peziza abietina** Pers. Syn. Fung. 637. 1801.

Otidea abietina Fuckel, Symb. Myc. 320. 1869.
Aleuria abietina Gill. Champ. Fr. Discom. 41. 1879.
Discina abietina Rehm in Rab. Krypt.-Fl. **1**[3]: 977. 1896.
Heteroplegma crenatum Clements, Bull. Torrey Club **30**: 92. 1903.
Peziza crenata Sacc. & D. Sacc. in Sacc. Syll. Fung. **18**: 19. 1906.
Pseudotis abietina Boud. Hist. Class. Discom. Eu. 52. 1907.
Plicaria rubrofusca Rehm, Ann. Myc. **3**: 517. 1905.
Peziza rubrofusca Sacc. & Trott. in Sacc. Syll. Fung. **22**: 619. 1913.

Apothecia gregarious or cespitose, stipitate or substipitate, cup-shaped, regular in form or becoming irregular with age, the margin entire or crenate, externally pale- to dark-brownish,

PEZIZA VENOSA

reaching a diameter of 3–5 cm. and a depth of 2–3 cm.; hymenium concave, brown, of nearly the same shade as the outside of the apothecium or a little darker; stem very short, stout, not exceeding 1 cm. in length and about 5 mm. thick, or often nearly wanting, the substance rather tough, not shrinking much in drying; asci cylindric or subcylindric, reaching a length of 200 μ and a diameter of 12–15 μ, 8-spored; spores ellipsoid, smooth, usually containing two large oil-drops and numerous smaller ones, 9–12 $\times$ 18–20 μ; paraphyses filiform, septate, enlarged above, with numerous small knots or branches near their apices, filled with coarse granules, reaching a diameter of 7 μ at their tips.

On the ground in coniferous woods.

TYPE LOCALITY: Europe.

DISTRIBUTION: Colorado; also in Europe.

ILLUSTRATIONS: Boud. Ic. Myc. *pl. 333;* Bres. Fungi Trid. *pl. 183, f. 2;* Cooke, Mycographia *pl. 57, f. 223;* Gonnerm. & Rab. Myc. Eu. **3**: *pl. 1, f. 7;* Fl. Dan. *pl. 1853, f. 1;* Payer, Bot. Crypt. 89, *f. 401.*

14. **Peziza venosa** Pers. Syn. Fung. 638. 1801. (excl. syn.) (PLATE 31.)

Peziza reticulata Grev. Scot. Crypt. Fl. 156. 1825.
Plicaria reticulata Fuckel, Symb. Myc. 328. 1869.
Aleuria venosa Gill. Champ. Fr. Discom. 37. 1879.
Acetabula venosa Lamb. Fl. Myc. Belg. **2**: 574. 1880.
Discina reticulata Sacc. Syll. Fung. **8**: 100. 1889.
Discina venosa Sacc. Syll. Fung. **8**: 104. 1889.
Peziza convoluta Peck, Bull. Torrey Club **30**: 101. 1903.
Disciotis venosa Boud. Hist. Class. Discom. Eu. 42. 1907.

Apothecia solitary or gregarious, substipitate with the base lacunose, at first shallow cup-shaped, soon becoming depressed and finally resting on the substratum with the extreme margin slightly elevated, when young entire but often irregularly radially splitting at maturity, externally whitish or dingy, slightly pruinose or scurfy, reaching a diameter of 20 cm.; hymenium reddish-brown, dried specimens often dusted over with the spores which are rust-colored in mass, at first even, becoming convolute, the convolutions either radial or giving a reticulate appearance; asci cylindric or subcylindric clavate, reaching a diameter of 25 μ; spores 1-seriate or slightly crowded together, ellipsoid, smooth, pale-yellowish, 12–17 $\times$ 22–30 μ; paraphyses clinging

together in masses, enlarged above, reaching a diameter of 8 μ, yellowish-brown.

On the ground in deciduous woods.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to Ohio; also in Europe.

ILLUSTRATIONS: Boud. Ic. Myc. *pl. 254;* Rab. Krypt.-Fl. 1^3: 922, *f. 1–3.*

EXSICCATI: N. Am. Fungi *2621.*

Very distinct with its reticulated hymenium but not very commonly collected.

15. **Peziza domiciliana** Cooke, Gardner's Chronicle **41**: 793. 1877. (PLATE 26, FIG. 2.)

?Octospora varia Hedw. Descr. **2**: 22. 1788.
Peziza Adae Sadler; Cooke, Trans. Bot. Soc. Edinburgh **13**: 45. 1877.
Peziza odorata Peck, Bull. Torrey Club **23**: 420. 1896.
Peziza varia f. *typica* Bres. Fungi Trid. **2**: 75. 1898.

Apothecia gregarious or occasionally cespitose, usually short-stipitate when young, at first concave, soon becoming repand, leaving a depression in the center, at first rounded, becoming irregular and often decidedly angular in outline, externally white, the margin entire or occasionally splitting, reaching a diameter of 10 cm., the substance when broken turning golden-yellow (it is uncertain whether this character is constant but it is very conspicuous in some cases); hymenium at first concave, becoming plane or convex and distinctly umbilicate, at first white, becoming dingy-buff or brownish; stem not exceeding 1 cm. in length, thick, irregular, white, becoming obsolete with age; asci cylindric or subcylindric, reaching a length of 225–250 μ and a diameter of 15 μ; spores ellipsoid, hyaline when young, often containing two small oil-drops, 8–10 $\times$ 13–15 μ; paraphyses slender, septate, slightly enlarged above.

In cellars, mushroom caves, and occasionally in greenhouses, usually growing on plaster, sand, gravel, or coal-dust.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to Iowa and Missouri; also in Europe.

ILLUSTRATIONS: Trans. Bot. Soc. Edinburgh **13**: *pl. 3, f. a–i;* Cooke, Mycographia, *pl. 97, f. 349;* Bres. Fungi Trid. *pl. 188;* Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 15, f. 2* (as *Peziza repanda*).

Rather frequently encountered.

PEZIZA REPANDA

16. **Peziza repanda** Pers. Ic. Pict. Fung. 49. 1806.

?Peziza micropus Pers. Ic. & Descr. Fung. 30. 1800.
Peziza pallida Cooke & Peck; Cooke, Buff. Acad. Sci. **2**: 288. 1875.
Peziza repanda amplispora Cooke & Peck; Cooke, Bull. Buff. Acad. Sci. **2**: 288. 1875.
Peziza truncicomes Ger.; Cooke, Mycographia 147. 1877.
Peziza amplispora Cooke, Mycographia 167. 1877.
Aleuria repanda Gill. Champ. Fr. Discom. 43. 1879.
Peziza Stevensoniana Ellis; Rehm, Ascom. Lojk. 3. 1882. (as synonym.)
Geopyxis pallidula Sacc. Syll. Fung. **8**: 70. 1889.
Geopyxis amplispora Sacc. Syll. Fung. **8**: 71. 1889.
Discina repanda Sacc. Syll. Fung. **8**: 100. 1889.
Plicaria repanda Rehm in Rab. Krypt.-Fl. 1^3: 1007. 1896.
Pustularia Stevensoniana Rehm in Rab. Krypt.-Fl. 1^3: 1019. 1896.
Peziza varia f. *lignicola* Bres. Fungi Trid. **2**: 76. 1898.

Apothecia sessile or very short-stipitate, at first cup-shaped, the margin even or crenate, externally white or whitish, expanding and becoming repand, the margin remaining entire or splitting, regular in outling or irregularly revolute, reaching a diameter of 8–10 cm.; hymenium concave, becoming plane or convex, pale-brown, becoming darker with age, even or convolute; stem short, stout, usually only a few mm. long or entirely wanting; asci cylindric or subcylindric, reaching a length of 225 μ and a diameter of 12–15 μ; spores ellipsoid, hyaline, smooth, 8–10 $\times$ 14–16 μ; paraphyses slender, slightly enlarged above, yellowish or brownish.

On rotten logs or occasionally on soil or chip-piles.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to Iowa and Maryland; also in Europe.

ILLUSTRATIONS: Pers. Ic. Pict. *pl. 20, f. 2;* ? Pers. Ic. Descr. Fung. *pl. 8, f. 3;* Cooke, Mycographia *pl. 62, f. 240;* Bres. Fungi Trid. *pl. 189;* Minn. Bot. Studies **4**: *pl. 15;* Freeman, Minn. Pl. Diseases *f. 64.*

One of the most conspicuous species on rotten logs in woods, its variability in form at different ages probably accounting for its numerous synonyms. When young it appears short-stipitate but when mature the stem is inconspicuous or obsolete.

17. **Peziza vesiculosa** Bull. Herb. Fr. *pl. 457, f. 1.* 1789. (PLATE 29, FIG. 2.)

Scodellina vesiculosa S. F. Gray, Nat. Arrang. Brit. Pl. 669. 1821.
?Elvela cochleata Bolton, Hist. Fung. **3**: 99 (in part). 1789.
Pustularia vesiculosa Fuckel, Symb. Myc. 329. 1869.
Aleuria vesiculosa Gill. Champ. Fr. Discom. 45. 1879.

Apothecia gregarious or more often densely cespitose, at first closed and globose, gradually expanding and becoming deep cup-shaped, regular in form or irregularly contorted, sessile or with a very stout stem-like base, externally whitish or yellowish, with minute wart-like pustules, reaching a diameter of 7–8 cm.; hymenium pale-brown, paler than the outside of the apothecium; asci cylindric or subcylindric; spores obliquely 1-seriate, ellipsoid, smooth, hyaline, 10–11 × 20–23 μ; paraphyses enlarged above granular within, subhyaline.

On manure piles and heavily fertilized soil.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to Washington, California, Alabama and probably throughout North America; also in Europe.

ILLUSTRATIONS: Bull. Herb. Fr. *pl. 457, f. 1;* Boud. Ic. Myc. *pl. 257;* Cooke, Mycographia *pl. 63, f. 242;* Gill. Champ. Fr. Discom. *pl. 44;* Rab. Krypt.-Fl. **1**[3]: 992, *f. 1–4;* Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 16, f. 1;* Sow. Engl. Fungi *pl. 4;* Massee, Brit. Fungus-Fl. **4**: 290, *f. 22;* Mycologia **7**: *pl. 155* (upper figure).

Very common on manure piles and on account of its large size conspicuous and attractive. It occurs less commonly on soil in greenhouses and occasionally in richly manured gardens. The cups are often regular in form or may become beautifully contorted or crimped.

18. **Peziza fimeti** (Fuckel) Seaver, comb. nov.

Humaria fimeti Fuckel, Symb. Myc. Nacht. **1**: 50. 1871.
?Peziza fimetaria Schum. Pl. Saell. **2**: 426. 1803.
?Peziza chlorophysa Sacc. & D. Sacc. in Sacc. Syll. Fung. **18**: 18. 1906.

Apothecia scattered, sessile, or substipitate, at first subglobose, becoming expanded and reaching a diameter of 2 cm., externally granulose, pallid-brown; hymenium concave to nearly plane, similar in color to the outside of the apothecium; asci cylindric or subcylindric, reaching a length of 280 μ and a diameter of 18 μ; spores ellipsoid, 8 × 16 μ; paraphyses abundant, slender, reaching a diameter of 8 μ.

On cow dung.

TYPE LOCALITY: Europe.

DISTRIBUTION: Kentucky to Iowa and Colorado; also in Europe.

ILLUSTRATION: Bres. Fungi Trid. *pl. 192, f. 2.*



PEZIZA SYLVESTRIS

Similar in habitat to the preceding but apparently distinguished by the small size of both apothecia and spores. The plants occur more on isolated cow dung than on manure piles as does the preceding.

19. **Peziza sylvestris** (Boud.) Sacc. & Trott. in Sacc. Syll. Fung. **22**: 612. 1913. (PLATE 33.)

Peziza varia f. *terrestris* Bres. Fungi Trid. **2**: 76. 1898.
Aleuria sylvestris Boud. Hist. Class. Discom. Eu. 44. 1907.

Apothecia gregarious, sessile, deep cup-shaped to subdiscoid, externally whitish, nearly smooth or minutely pustulate, the margin even or slightly wavy and crenate, reaching a diameter of 3–8 cm.; hymenium umber-brown; asci cylindric above, gradually tapering below, reaching a length of 200–235 μ and a diameter of 13–15 μ; spores 1-seriate, with the ends slightly overlapping, hyaline, smooth, 9–10 $\times$ 15–20 μ; paraphyses strongly enlarged above, reaching a diameter of 7–8 μ at their apices.

On rubbish piles and soil in woods.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to California; also in Europe.

ILLUSTRATIONS: Boud. Ic. Myc. *pl. 261;* Bres. Fungi Trid. *pl. 190;* Mycologia **7**: *pl. 156* (lower figure).

This may be *Peziza cerea* Sow. the identity of which seems to be uncertain. The species is common in woods and often approaches *Peziza repanda* in size and general appearance. The latter seems to differ however in its more repand habits as well as in habitat, occurring on rotten logs.

20. **Peziza clypeata** Schw. Schr. Nat. Ges. Leipzig **1**: 117. 1822. (PLATE 34.)

Bulgaria bicolor Peck, Ann. Rep. N. Y. State Mus. **32**: 49. 1879.
Peziza orbicularis Peck, Bull. N. Y. State Mus. **2**: 20. 1887.
Discina clypeata Sacc. Syll. Fung. **8**: 101. 1889.
?Peziza Barleana Bres. Fungi Trid. **2**: 74. 1898.
?Peziza paraphysata Clements, Bot. Surv. Nebr. **4**: 9. 1896.
Psilopezia juruensis P. Henn. Hedwigia **43**: 273. 1904.
?Pachyella Barleana Boud. Hist. Class. Discom. Eu. 50. 1907.
Psilopezia orbicularis Dodge, Trans. Wisc. Acad. Sci. **17**: 1052. 1914.

Apothecia gregarious but not usually crowded, at first globose, gradually expanding, soon becoming shallow cup-shaped, finally discoid, regular in outline or occasionally more or less distorted when old, becoming closely adnate to the substratum on the under surface, except the extreme margin, which is free and

often slightly upturned, varying in size from a few mm. when young to 3 or 4 cm. when mature; hymenium at first dark reddish-brown, smooth and glistening, later assuming an olive tint and losing its luster, when old greenish-black (almost entirely black when dried), splitting in dried specimens, giving rise to whitish vein-like markings where the whitish interior is exposed through the crevices, the substance soft, inclined to waxy; asci cylindric, reaching a diameter of 17 μ, often spirally twisted when dry; spores 1-seriate, ellipsoid, smooth, hyaline, containing one or two large oil-drops, 12–14 $\times$ 25–27 μ or rarely 30–35 μ; paraphyses adhering together, septate, strongly enlarged above, reaching a diameter of 7–8 μ, densely filled with yellowish-brown coloring matter.

On much decayed deciduous logs of various kinds, especially where they are saturated with water.

TYPE LOCALITY: North Carolina.

DISTRIBUTION: New York to Wisconsin and North Carolina; also in South America.

ILLUSTRATIONS: Bull. N. Y. State Mus. **2**: *pl. 2, f. 4–6;* Hedwigia **43**: *273, f. 1–3;* (?) Boud. Ic. Myc. *pl. 310;* (?) Bres. Fungi Trid. *pl. 187.*

EXSICCATI: N. Am. Fungi *568* (as *Psilopezia nummularia* Berk.).

Very distinct with its adnate, dark-colored, discoid apothecia. This species has been collected in New York City year after year in the same place on the decaying trunk of a large tree, occurring in the late summer or autumn.

DOUBTFUL SPECIES

Peziza caeruleomaculata (Rehm) Sacc. & D. Sacc. in Sacc. Syll. Fung. **18**: 19. 1906; *Plicaria caeruleomaculata* Rehm, Ann. Myc. **2**: 351. 1904. This may be an abnormal specimen of *Peziza pustulata* (Hedw.) Pers.

Humaria lacteocinerea Rehm, Ann. Myc. **3**: 517. 1905. A whitish species reaching a diameter of 2 cm. and with coarsely verrucose spores 7–8 $\times$ 12–13 μ. On a pine drain in greenhouse at Madison, Wisconsin. The identity of the specimens is uncertain.

Peziza brunneovinosa Clements, Bot. Surv. Nebr. **4**: 8. 1896. A brownish or wine-colored species, with discoid apothecia reaching a diameter of 1.5 cm. and with smooth spores 10 $\times$ 16–20 μ. On soil, Keya Paha County, Nebraska. No specimen has been seen.

Peziza chrysopela Cooke, Mycographia 156. 1877. A small species described as one-half inch in diameter, subdiscoid, with pale-orange hymenium and smooth spores 6 $\times$ 12 μ. On the ground in South Carolina. No specimen has been seen.

PEZIZA CLYPEATA

Peziza roseolilacina Clements, Bot. Surv. Nebr. **5**: 7. 1901. Described as yellowish-ochraceous with a rose-lilac hymenium, reaching a diameter of 1.5 cm., cup-shaped, with smooth spores 10 × 17–18 μ. On wet soil among mosses, Keya Paha County, Nebraska. No specimens seen.

Peziza caligans Cooke, Mycographia 149. 1877. The description suggests a small specimen of *Peziza badia*. Its identity however is uncertain.

Peziza irrorata Berk. & Curt.; Berk. Grevillea **3**: 150. 1875. This species is described as applanate, fuligineous, 2–3 cm. in diameter, with spores 12 × 18–20 μ. On soil in Texas. No specimen seen.

Peziza repandoides (Rehm) Sacc. & Trott. in Sacc. Syll. Fung. **22**: 614. 1913; *Plicaria repandoides* Rehm, Ann. Myc. **3**: 518. 1905. This appears from the description to be a small specimen of *Peziza repanda*.

Peziza irregularis Berk. & Curt.; Berk. Jour. Linn. Soc. **10**: 365. 1868. This species is described as 6–8 cm. in diameter, with an orange hymenium and spores 12 × 25 μ. Both the size of the apothecia and the size and form of the spores are strongly suggestive of a large *Phillipsia* which occurs in the West Indies. The color of the *Phillipsia* is reddish-purple, however, rather than orange as described and illustrated by Cooke. No authentic specimen has been seen.

Peziza convoluta Peck, Bull. Torrey Club **30**: 101. 1903. Apothecia yellowish or tawny-brown with a wavy margin and a contorted hymenium, the spores 12–15 × 20–22 μ. On sandy soil in California. No specimens have been seen.

39. **SARCOSPHAERA** Auersw. Hedwigia **8**: 82. 1869.

Apothecia gregarious or scattered, at first entirely immersed in the substratum, the margin splitting *Geaster*-like at maturity; asci 8-spored; spores smooth or nearly so, ellipsoid or subellipsoid, hyaline; paraphyses filiform.

Type species, *Peziza macrocalyx* Reiss.

1. **Sarcosphaera coronaria** (Jacq.) Schröt. Krypt.-Fl. Schles. **3**[2]: 49. 1893.

Peziza coronaria Jacq. Misc. Austr. **1**: 140. 1778.
Peziza amplissima Fries, Summa Veg. Scand. 349. 1849.
Peziza Geaster Rab. Sitzber. Nat. Ges. Isis. **1867**: 22. 1867.
Sarcosphaera macrocalyx Auersw. Hedwigia **8**: 82. 1869.
?Peziza sicula Inzenga, Fung. Sic. **2**: 39. 1869.
Peziza Clissoni Ripart. Bull. Soc. Bot. Fr. **23**: 309. 1876.
Aleuria eximia Gill. Champ. Fr. Discom. 48. 1879.
Pustularia gigantea Rehm, Ann. Myc. **3**: 517. 1905.
Peziza gigantea Sacc. & Trott. in Sacc. Syll. Fung. **22**: 611. 1913.

Apothecia occurring singly or two to five closely crowded together, at first buried, later becoming partially exposed, subglobose or more or less irregularly contorted, splitting from the top downward into 7–10 pointed rays, externally dirty-pallid and

at first clothed with a transient mycelium, at length naked, reaching a diameter of 6 cm. and a depth of 8–10 cm. when expanded, extended below into a short stem-like base; hymenium at first pallid, becoming bluish or violaceous; asci cylindric or subcylindric, reaching a length of about 200 μ and a diameter of 10–12 μ; spores 1-seriate, smooth or very minutely roughened, usually containing two oil-drops, 8–9 $\times$ 15–18 μ; paraphyses enlarged and branched above, filled with bluish coloring matter when fresh, the color fading when dry.

In coniferous woods, on or under humus.

TYPE LOCALITY: Europe.

DISTRIBUTION: British Columbia to Michigan and New York; also in Europe and North Africa.

ILLUSTRATIONS: Boud. Ic. Myc. *pl. 302;* Cooke, Mycographia *pl. 61;* E. & P. Nat. Pfl. 1[1]: 181, *f. 147, F–G;* Fres. Beit. Myk. *pl. 9, f. 7–11;* Gill. Champ. Fr. Discom. *pl. 48;* Gonnerm. & Rab. Myc. Eu. Pez. *pl. 3, f. 5;* Inzenga, Fung. Sic. **2**: *pl. 8, f. 4;* Jacq. Misc. Austr. **1**: *pl. 10;* Jour. Bot. **7**: *pl. 98;* Bull. Torrey Club **45**: *pl. 3;* Lloyd, Myc. Notes 56, *f. 1255;* Sitzber. Nat. Ges. Isis. **1867**: *pl. 1.*

A most beautiful species but seldom collected. The only freshly collected material examined was that collected by Mr. Stewart H. Burnham in Washington county, New York. Recognized by its semisubterranean habits and the splitting of the apothecia.

Family 2. **ELVELACEAE**

Ascophores consisting of a distinct stem and pileus; pileus bell-shaped, saddle-shaped, subglobose or entirely covering the outside of the stem when it assumes a columnar form, the surface of the hymenium even, lacunose or costate, the margin of the pileus either free or partially to entirely attached to the stem; stem very slender or stout, even, lacunose, or distinctly fluted, naked or pubescent, usually hollow within; asci cylindric or subcylindric, 2–8-spored; spores ellipsoid, smooth or sculptured, hyaline or faintly colored; paraphyses variable in form, hyaline or slightly colored.

Upper surface of the pileus costate.
  With a sterile stem and fertile head. 1. *Morchella.*
  Without stem and fertile to the base. 2. *Durandiomyces.*
Upper surface of the pileus even, lacunose, or gyrose, not costate.

Pileus bell-shaped. 3. *Verpa.*
Pileus saddle-shaped or irregularly subglobose. 4. *Elvela.*
Pileus columnar. 5. *Underwoodia.*

1. **MORCHELLA** (Dill.) Pers. Tent. Disp. Fung. 36. 1797.

Morchella Dill. App. Cat. Pl. 74. 1719.
Phalloboletus Adans. Fam. Pl. 9. 1763. Hyponym.
Morilla Quél. Ench. Fung. 270. 1886.
Phalloboletus Kuntze, Rev. Gen. Pl. **2**: 864. 1891.

Pileus subglobose or elongated, obtuse or acute at the apex, the margin closely adnate or rarely free from the stem at the base, the surface irregularly costate, the rounded or elongated pits entirely lined within by the hymenium; the ribs irregularly anastomosing; hymenium yellow to brown; asci cylindric or subcylindric, usually 8-spored; spores ellipsoid, smooth, hyaline or faintly colored; paraphyses few, usually stout.

Type species, *Phallus esculentus* L.

Pileus adnate to the stem at the base.
Ribs of the pileus similar in color to the interior of the pits or lighter.
Pileus large, reaching a length of 4–8 cm. at maturity.
Pits large, usually shallow; ribs thin; stem strongly enlarged at the base. 1. *M. crassipes.*
Pits small, deep; ribs thick; stem only slightly enlarged at the base.
Pileus subglobose or only slightly elongated. 2. *M. esculenta.*
Pileus elongated or strongly attenuated upwards. 3. *M. conica.*
Pileus small, not exceeding 2–3 cm. in length. 4. *M. deliciosa.*
Ribs of the pileus much darker than the interior of the pits, becoming smoky-brown. 5. *M. angusticeps.*
Pileus free from the stem at the base. 6. *M. hybrida.*

1. **Morchella crassipes** (Vent.) Pers. Syn. Fung. 621. 1801. (PLATE 35.)

Phallus crassipes Ventenat, Mem. Inst. 509. 1798.
Phalloboletus crassipes Kuntze, Rev. Gen. Pl. **2**: 865. 1891.

Pileus usually elongated and often subconic, reaching a length of 6–8 cm. and a diameter of 5–6 cm.; pits roundish or irregularly elongated, reaching a diameter of 1 cm., yellowish within; ribs irregularly anastomosing, thin, often sharp-edged; stem stout, reaching a length of 10–11 cm. and a diameter of 5–6 cm. at the base, tapering upwards to a diameter of 4 cm., yellowish or whitish, more or less lacunose, or nearly even,

slightly pubescent; asci cylindric or subcylindric, reaching a length of 200 μ and a diameter of 20 μ; spores 1-seriate, ellipsoid, hyaline or subhyaline, 12–14 $\times$ 20–22 μ; paraphyses stout, clavate, hyaline or subhyaline.

On the ground in open places, often in margins of wooded areas.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to Montana and Iowa; also in Europe.

ILLUSTRATIONS: Atkinson, Stud. Am. Fungi ed. 2, *pl. 86;* Boud. Ic. Myc. *pl. 194;* Cooke, Mycographia *pl. 84, f. 319;* Gill. Champ. Fr. Discom. *pl. 14;* Krombh. Abbild. *pl. 16, f. 1–2;* Hedwigia **25**: 136; Mem. Inst. 509, *f. 2.*

2. **Morchella esculenta** (L.) Pers. Syn. Fung. 618. 1801. (PLATE 36, FIG. 2.)

Phallus esculentus L. Sp. Pl. 1178. 1753.
Morchella esculenta rotunda Pers. Syn. Fung. 619. 1801.
Morchella esculenta vulgaris Pers. Syn. Fung. 619. 1801.
Morchella continua Tratt. Fungi Austr. 67. 1830 (in part).
Morchella esculenta fulva Fries, Syst. Myc. **2**: 7. 1822.
Morchella esculenta longipes Peck, Ann. Rep. N. Y. State Mus. **28**: 87. 1876.
Phalloboletus esculentus Kuntze, Rev. Gen. Pl. **2**: 865. 1891.
Morchella rotunda Boud. Soc. Myc. Fr. **13**: 135. 1897.
Morchella vulgaris Boud. Bull. Soc. Myc. Fr. **13**: 139. 1897.
Morilla esculenta Quél. Ench. Fung. 271. 1886.

Pileus subglobose, ovoid, or elongated, occasionally attenuated upwards but obtuse at the apex, varying much in size but often reaching a diameter of 4–5 cm. and a length of 7–9 cm.; pits rounded, irregular, or occasionally longitudinally elongated, yellowish within, becoming brownish or blackish when dry, reaching a diameter of 5 mm. to 1 cm.; ribs irregularly anastomosing, not longitudinally disposed, the edges rounded, about 1 mm. thick and lighter colored than the interior of the pits, usually whitish or yellowish; stem stout but not usually exceeding two-thirds the diameter of the pileus, usually a little enlarged at the base and irregularly lacunose, nearly even above, whitish to yellowish, lighter than the pileus, delicately pubescent, reaching a length of 200–250 μ and a diameter of 18–20 μ; spores 1-seriate, ellipsoid, hyaline, yellowish in mass, 12–14 $\times$ 20–24 μ; paraphyses strongly enlarged above, where they reach a diameter of 15 μ, faintly colored.

MORCHELLA CRASSIPES

On the ground in woods and open places.

TYPE LOCALITY: Europe.

DISTRIBUTION: Massachusetts to Michigan and Colorado, south to South Carolina; probably throughout North America; also in Europe.

ILLUSTRATIONS: Sow. Engl. Fungi *pl. 51* (in part); Bolton, Hist. Fung. *pl. 91;* Boud. Ic. Myc. *pl. 195*, *196* and *202;* Bull. Herb. Fr. *pl. 216;* Cooke, Mycographia *pl. 81* (in part); Grev. Scot. Crypt. Fl. **2**: *pl. 68;* Gill. Champ. Fr. Discom. *pl. 13;* Massee, Brit. Fungus-Fl. **4**: 188, *f. 1–3;* Phill. Brit. Discom. *pl. 1*, *f. 1;* Schaeff. Fung. Bavar. *pl. 298*, *299;* Krombh. Abbild. *pl. 19*, *f. 6–7*.

EXSICCATI: N. Am. Fungi *979*.

3. **Morchella conica** Pers. Champ. Comest. 257. 1819.

?Phallus acuminatus Batsch. Elench. Fung. 133. 1783.
?Morchella continua Tratt. Fungi Austr. 67. 1830 (in part).
Morchella esculenta conica Fries, Syst. Myc. **2**: 7. 1822.
Morilla conica Quél. Ench. Fung. 271. 1886.
Phalloboletus conicus Kuntze, Rev. Gen. Pl. **2**: 865. 1891.

Pileus elongated, subconic, acute at the apex or subobtuse, reaching a length of 6–8 cm. and a diameter of 4 cm.; pits elongated and more or less inclined to rectangular in form, yellowish within, becoming brown or blackish when dry, reaching a length of 5 mm. to 1 cm. and about half as broad; ribs more or less longitudinally disposed, anastomosing or connected by cross ribs, the edges about 1 mm. thick, rounded and lighter in color than the interior of the pits; stem about two-thirds the diameter of the base of the pileus, irregularly lacunose below, nearly even above, whitish to yellowish, lighter than the pileus, delicately pubescent, reaching a length of 4–7 cm.; asci cylindric or subcylindric, reaching a length of 200–250 μ and a diameter of 18–20 μ; spores 1-seriate, ellipsoid, hyaline, yellowish in mass, 12–14 $\times$ 20–24 μ; paraphyses strongly enlarged above, where they reach a diameter of 12–15 μ, faintly colored.

On the ground in woods and open places in the margins of woods.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to Washington and Iowa; also in Europe.

ILLUSTRATIONS: Tratt. Fungi Austr. *pl. 6*, *f. 11;* Rab. Krypt.-Fl. 1^3: 1178, *f. 1–4;* Atkinson, Stud. Am. Fungi ed. 2, 218, *f. 217;*

Gill. Champ. Fr. Discom. *pl. 16, f. 1;* Krombh. Abbild. *pl. 16, f. 7–12.*

The species is regarded by many as only a form of *M. esculenta.*

4. **Morchella deliciosa** Fries, Syst. Myc. **2**: 8. 1822.

Morchella conica deliciosa Phill. Brit. Discom. 5. 1887.
Phalloboletus deliciosus Kuntze, Rev. Gen. Pl. **2**: 865. 1891.

Pileus conic or subconic, reaching a length of 2–3 cm. and a diameter of 1–1.5 cm.; pits usually elongated, cinereous to blackish within; ribs inclined to be longitudinally disposed, irregularly anastomosing, the edges rounded and about 1 mm. thick, much lighter than the interior of the pits, whitish; stem half to two-thirds as thick as the base of the pileus, often enlarged at the base and irregularly lacunose, lighter than the pileus, whitish or yellowish; asci cylindric or subcylindric, reaching a length of about 200 μ and a diameter of 12–15 μ; spores 1-seriate, ellipsoid, hyaline, yellowish in mass, $10 \times 20\ \mu$; paraphyses enlarged at their apices and slightly colored.

On the ground in grassy places, usually in the edges of wooded places.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to Pennsylvania, Iowa and Colorado; also in Europe.

ILLUSTRATIONS: Trans. Bot. Soc. Edinburgh **10**: 441, *f. 1;* Cooke, Mycographia *pl. 84, f. 320;* Gill. Champ. Fr. Discom. *pl. 15;* Massee, Brit. Fungus-Fl. **4**: 188, *f. 31;* Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 2, f. 2;* Krombh. Abbild. *pl. 16, f. 17–19.*

5. **Morchella angusticeps** Peck, Ann. Rep. N. Y. State Mus. **32**: 44. 1879.

?Morchella elata Fries, Syst. Myc. **2**: 8. 1822.
?Morilla elata Quél. Ench. Fung. 271. 1886.
?Phalloboletus elatus Kuntze, Rev. Gen. Pl. **2**: 865. 1891.
Phalloboletus angusticeps Kuntze, Rev. Gen. Pl. **2**: 865. 1891.

Pileus elongated, subobtuse or narrowly conic and acute at the apex, reaching a length of 2–5 cm. and about half as broad at the base; pits elongated, reaching a length of 5–10 mm. and about half as broad, yellowish within, becoming smoky-brown at the margins; ribs inclined to be longitudinally disposed, irregularly anastomosing, usually less than 1 mm. thick and often almost sharp-edged, black at the extreme edge, lighter



1. MORCHELLA HYBRIDA
2. MORCHELLA ESCULENTA

below; stem of almost uniform thickness throughout its entire length, the diameter almost equal to that of the pileus, irregularly lacunose near the base, nearly even above, delicately pubescent; asci cylindric or subcylindric, reaching a length of 275 μ and a diameter of 20 μ; spores 1-seriate, ellipsoid, hyaline or very faintly colored, yellowish in mass, 12–14 × 24–28 μ; paraphyses enlarged upwards.

On sandy soil in the borders of woods and open places.

TYPE LOCALITY: West Albany, New York.

DISTRIBUTION: New York to Pennsylvania and Michigan; also in Europe.

ILLUSTRATIONS: Bull. N. Y. State Mus. 1[2]: *pl. 1, f. 19–20;* Ann. Rep. N. Y. State Mus. **48**: *pl. 4, f. 5–9;* Boud. Ic. Myc. *pl. 206* and (? *213*).

6. **Morchella hybrida** (Sow.) Pers. Syn. Fung. 621. 1801. (PLATE 36, FIG. 1.)

?Phallus Rete Batsch. Elench. Fung. 131. 1783.
Elvela hybrida Sow. Engl. Fungi *pl. 238*. 1799.
?Morchella patula Pers. Syn. Fung. 619. 1801.
Morchella rimosipes DC. Fl. Fr. **2**: 214. 1815.
?Morchella Rete Pers. Myc. Eu. **1**: 205. 1822.
Mitrophora semilibera Lév. Ann. Sci. Nat. III. **5**: 250. 1846.
?Mitrophora patula Lév. Ann. Sci. Nat. III. **5**: 250. 1846.
Mitrophora rimosipes Lév. Ann. Sci. Nat. III. **5**: 250. 1846.
Morilla semilibera Quél. Ench. Fung. 271. 1886.
Morchella Mitra Lenz, Schwamme 103. 1831.
Phalloboletus Rete Kuntze, Rev. Gen. Pl. **2**: 865. 1891.
Morchella hybrida rimosipes P. Henn. Verh. Bot. Brandenb. **36**: 68. 1895.
Phalloboletus rimosipes Kuntze, Rev. Gen. Pl. **2**: 865. 1891.
Morchella punctipes Peck, Bull. Torrey Club **30**: 99. 1903.
Mitrophora hybrida Boud. Hist. Class. Discom. Eu. 33. 1907.

Pileus bell-shaped or subconic, reaching a length of 2 cm. and a diameter of 1.5 cm., the margin free about half way up; pits irregularly rounded or elongated, reaching a diameter of 5–10 mm., yellowish within; ribs inclined to be longitudinal or irregularly anastomosing, about 1 mm. thick, whitish, lighter than the interior of the pits; stem thick, at first very short, finally reaching a length of 8–10 cm. and a diameter of 2 cm. at the base, gradually tapering upward to about half that diameter, delicately pubescent, whitish or yellowish, often irregularly lacunose at the base, nearly even above; asci cylindric or subcylindric, reaching a length of 250 μ and a diameter of 20 μ;

spores 1-seriate, ellipsoid, 12–14 × 22–26 μ, hyaline or faintly colored; paraphyses enlarged upwards, hyaline or subhyaline.

On the ground, usually in the margins of woods.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to Iowa and Oregon; also in Europe.

ILLUSTRATIONS: Boud. Ic. Myc. *pl. 216;* Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 3, f. 2;* Cooke, Mycographia *pl. 85;* Minn. Bot. Studies **3**: *pl. 48, f. 7, 8;* Lenz, Schwamme *pl. 16, f. 67;* Mycologia **19**: 141, *f. 3.*

EXSICCATI: N. Am. Fungi *2628, 2907.*

2. **DURANDIOMYCES** Seaver, gen. nov.

Ascophores large, consisting of numerous contorted and frequently anastomosing ribs resembling a *Sparassis*, the numerous cavities lined everywhere with the hymenium, the color ranging from white to rosy or faintly violaceous; flesh very brittle; asci 8-spored; spores minutely sculptured; spore-sculpturing consisting of minute warts or papillae.

Type species, *Gyromitra Phillipsii* Massee.

Originally thought and still maintained by some to be an anomalous form of *Peziza proteana* which it resembles in spore characters and in color. Although familiar with *Peziza proteana*, I have never seen any tendency to assume this form and therefore regard it as a well-defined genus. It is here dedicated to the late Dr. E. J. Durand, to whom we are indebted for the excellent photograph. (PLATE 37.)

Durandiomyces Phillipsii (Massee) Seaver, comb. nov. (PLATE 37.)

Gyromitra Gigas Phill. Jour. Bot. **31**: 129. 1893. Not *G. Gigas* Krombh.
Gyromitra Phillipsii Massee, Brit. Fungus-Fl. **4**: 478. 1895.
Aleuria proteana sparassoides Boud. Bull. Soc. Myc. Fr. **15**: 51. 1899.
Galactinia proteana sparassoides Boud. Ic. Myc. 162. 1911.
Peziza proteana sparassoides Durand, Mycologia **11**: 1. 1919.

Ascophores large, reaching a height of 20–25 cm. and a diameter of 15–20 cm.; asci cylindric or subcylindric, scarcely attenuated at the base, reaching a length of 250–300 μ and a diameter of 10–11 μ; spores ellipsoid, usually containing two oil-drops, becoming minutely sculptured, 7 × 11–12 μ; spore-sculpturing consisting of minute warts or papillae; paraphyses enlarged upwards, reaching a diameter of 5–7 μ at their apices.



DURANDIOMYCES PHILLIPSII

On the ground in burnt places.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to Washington and Oregon; also in Europe.

ILLUSTRATIONS: Bull. Soc. Myc. Fr. **15**: *pl. 3, f. 2;* Boud. Ic. Myc. *pl. 294;* Jour. Bot. **31**: *pl. 334;* Mycologia **11**: *pl. 1.*

3. **VERPA** Swartz, Sv. Vet. Akad. Handl. **1815**: 129.

Pileus bell-shaped, with the stem attached to the center of the under surface; hymenium covering the entire outside of the pileus, varying in color from yellow to dark-brown, even or more or less lacunose; stem rather stout, hollow or loosely stuffed, lighter colored than the hymenium, usually even or very slightly lacunose, naked or delicately pubescent; asci cylindric or clavate, 2–8-spored; spores ellipsoid, smooth, medium sized to very large; paraphyses rather stout.

Type species, *Phallus conicus* Müller.

Asci 8-spored; spores medium sized.	1. *V. conica.*
Asci typically 2-spored; spores very large.	2. *V. bohemica.*

1. **Verpa conica** (Müll.) Swartz, Sv. Vet. Akad. Handl. **1815**: 136. 1815. (PLATE 38, FIG. 2.)

Phallus conicus Müller, Fl. Dan. *pl. 654, f. 2.* 1777.
Leotia conica Pers. Syn. Fung. 613. 1801.
Verpa digitaliformis Pers. Myc. Eu. **1**: 202. 1822.
Verpa Krombholzii Corda in Sturm, Deutsch. Fl. **2**: 5. 1829.
Verpa helvelloides Krombh. Abbild. **1**: 76. 1831.
Phalloboletus conicus Kuntze, Rev. Gen. Pl. **2**: 865. 1891.
Verpa chicoensis Copeland, Ann. Myc. **2**: 508. 1904.

Pileus bell-shaped or subconic, whitish beneath, the margin often slightly reflexed, exposing a whitish border, entirely free from the stem, reaching a depth of 2 cm. and a diameter of 1–2 cm.; hymenium becoming dark-brown, even or slightly lacunose; stem cylindric or subcylindric, slightly narrowed above, whitish or slightly floccose or scaly, the scales giving a transversely striate appearance, hollow or very loosely stuffed, reaching a length of 5–6 cm.; asci cylindric or subcylindric, gradually tapering below into a more or less contorted, stem-like base, reaching a length of 300 μ and a diameter of 20 μ, 8-spored; spores 1-seriate or irregularly crowded, ellipsoid, slightly yellowish at maturity, 12–16 $\times$ 22–26 μ; paraphyses stout, sparingly septate, reaching a diameter of 10 μ.

On the ground.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to Colorado; also in Europe.

ILLUSTRATIONS: Boud. Ic. Myc. *pl. 220;* Cooke, Mycographia *pl. 101, f. 364;* Fl. Dan. *pl. 654, f. 2;* Pers. Myc. Eu. **1**: *pl. 7, f. 1–3;* Sturm, Deutsch. Fl. **2**: *pl. 1;* Ann. Myc. **2**: *pl. 12, f. 1–5;* Rab. Krypt.-Fl. 1^3: 1175, *f. 2–3.*

2. **Verpa bohemica** (Krombh.) Schröt. Krypt. Fl. Schles. 3^2: 25. 1893. (PLATE 38, FIG. 1.)

Morchella bohemica Krombh. Abbild. **3**: 3. 1834.
Morchella gigaspora Cooke, Trans. Bot. Soc. Edinb. **10**: 442. 1870.
Morchella bispora Sorokin, Mycol. Unters. 21. 1872.
Morchella bohemica bispora Cooke, Mycographia 188. 1878.
Morilla bohemica Quél. Ench. Fung. 271. 1886.
Morilla gigaspora Quél. Ench. Fung. 271. 1886.
Phalloboletus bohemicus Kuntze, Rev. Gen. Pl. **2**: 865. 1891.
Phalloboletus gigasporus Kuntze, Rev. Gen. Pl. **2**: 865. 1891.
Phalloboletus bisporus Kuntze, Rev. Gen. Pl. **2**: 865. 1891.
Morchella bispora truncata Peck, Ann. Rep. N. Y. State Mus. **46**: 38. 1893.

Pileus bell-shaped or subconic, white underneath, often with the margin slightly reflexed, exposing a white border below, 2–3 cm. deep and reaching a diameter of 1–2 cm.; hymenium folded into longitudinal and rather freely anastomosing ribs, yellowish to brownish; stem almost cylindric, hollow or loosely stuffed, externally white or slightly floccose, not wrinkled or lacunose, reaching a length of 6–8 cm.; asci cylindric or subcylindric, tapering below into a much contorted stem-like base, the entire ascus reaching a length of 300–325 μ and a diameter of 20–22 μ, typically 2-spored; spores ellipsoid, straight or slightly curved, smooth externally, and minutely granular within, 15–18 $\times$ 60–80 μ, subhyaline or slightly yellowish; paraphyses clavate, stout, reaching a diameter of 7–8 μ at their apices.

On the ground.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to Minnesota; also in Europe.

ILLUSTRATIONS: Boud. Ic. Myc. *pl. 218;* Krombh. Abbild. *pl. 15, f. 1–13;* Minn. Bot. Studies **3**: *pl. 49, f. 7–9;* Trans. Bot. Soc. Edinb. **10**: 442, *f. 2;* Rab. Krypt.-Fl. 1^3: 1176, *f. 1–3;* Cooke, Mycographia *pl. 87, f. 324–326;* Bot. Zeit. **34**: *pl. 10;* E. & P. Nat. Pfl. 1^1: 168, *f. 141, E–F* and 169, *f. 142, G.*



1. VERPA BOHEMICA
2. VERPA CONICA

DOUBTFUL SPECIES

Verpa atroalba Fries, Syst. Myc. **2**: 23. 1822. A specimen referred to this name was collected at Jamesville, New York, May 17, 1890. The specimen is identical with *Verpa conica*. No authentic specimen of Fries' species has been seen and its identity is therefore uncertain.

Verpa perpusilla Rehm, Ann. Myc. **7**: 526. 1909. Specimens of this species distributed by Rehm in his Ascomycetes *1857* appear to be identical with *Elvela elastica* Bull.

4. **ELVELA** L. Sp. Pl. 1180. 1753. (Often written *Helvella*.)

Gyromitra Fries, Summa Veg. Scand. 346. 1849.
Leptopodia Boud. Bull. Soc. Myc. Fr. **1**: 99. 1885.
Physomitra Boud. Hist. Class. Discom. Eu. 35. 1907.

Pileus mitrate, saddle-shaped or subglobose, even or irregularly convolute, the convolutions often taking the form of gyrose ridges, the margin reflexed and free or more or less adnate to the stem; stem very slender or stout, even, lacunose, or strongly fluted, white, yellow, or smoky; asci cylindric or subcylindric, 8-spored; spores 1- or 2-seriate, ellipsoid, often apiculate, smooth or rough; paraphyses slender, more or less enlarged above.

Type species, *Elvela Mitra* L.

Stem distinctly fluted, *i.e.* with sharp-cornered ridges which extend almost the entire length of the stem.
  Stem stout and usually strongly swollen at the base.
    Pileus becoming dark-colored and stem smoky. — 1. *E. Mitra.*
    Pileus and stem remaining light-colored, almost white. — 2. *E. crispa.*
  Stem very slender and of nearly uniform thickness. — 3. *E. palustris.*
Stem not distinctly fluted but even or irregularly lacunose.
  Stem very slender, rarely exceeding 5 mm. in diameter.
    Pileus and stem entirely black. — 4. *E. atra.*
    Pileus and stem not entirely black.
      Pileus entirely free from the stem.
        Stem very long. — 5. *E. elastica.*
        Stem very short. — 6. *E. Klotzschiana.*
      Pileus adnate to the stem. — 7. *E. adhaerens.*
  Stem stout, usually 1 to several cm. in diameter.
    Spores ellipsoid to fusoid.
      Spores small, smooth, not exceeding 20–24 μ long.

Pileus large and umbrella-like; spores small, 8–9 × 17 μ.
Stem long, reaching a length of 10 cm. 8. *E. californica.*
Stem short, reaching a length of 3–5 cm. 9. *E. umbraculiformis.*
Pileus not widely spreading or umbrella-like.
Spores narrow-ellipsoid, containing two oil-drops.
Pileus medium large; stem reaching a length of 6–8 cm. 10. *E. infula.*
Pileus very small; stem not exceeding 1–2 cm. in length. 11. *E. brevissima.*
Spores broad-ellipsoid, containing one oil-drop. 12. *E. albipes.*
Spores large, rough, 12–14 × 30–35 μ.
Spores fusoid, strongly apiculate. 13. *E. caroliniana.*
Spores ellipsoid, not strongly apiculate. 14. *E. Underwoodii.*
Spores globose. 15. *E. sphaerospora.*

1. **Elvela Mitra** L. Sp. Pl. 1180. 1753. (PLATE 39, FIG. 1.)

Elvela nigricans Schaeff. Fung. Bavar. **4**: Ind. 102. 1774.
Elvela lacunosa Afz. Sv. Vet. Akad. Handl. 304. 1783.
Elvela sulcata Afz. Sv. Vet. Akad. Handl. 305. 1783.
?Elvela grisea Clements, Bot. Surv. Nebr. **4**: 8. 1896.
Leptopodia elastica Boud. Hist. Class. Discom. 37. 1907.

Pileus mitrate, irregularly saddle-shaped or occasionally subglobose, the margin reflexed and adnate with the stem at various points, occasionally extending down the stem for some distance, reaching a diameter of 2–5 cm.; hymenium even or irregularly convolute, dark brownish-black; stem reaching a length of 10 cm. and a diameter of 2 cm., usually enlarged below and gradually attenuated above, deeply fluted, yellowish, becoming smoky; asci cylindric, 8-spored, reaching a length of 200 μ and a diameter of 16 μ; spores ellipsoid, usually containing one large oil-drop which almost fills the spore, about 12 × 20 μ; paraphyses enlarged at their apices, reaching a diameter of 5 μ, pale yellowish-brown.

On damp soil.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to British Columbia and south to Mexico; also in Europe.

ILLUSTRATIONS: Micheli, Nov. Pl. Gen. *pl. 86, f. 7;* Schaeff.



1. ELVELA MITRA
2. ELVELA CRISPA

Fung. Bavar. *pl. 154;* Boud. Ic. Myc. *pl. 228;* Cooke, Mycographia *pl. 41, f. 160;* Gill. Champ. Fr. Discom. *pl. 2, f. 1* and *pl. 3;* Bull. Herb. Fr. *pl. 466, f. B;* Minn. Bot. Studies **3**: *pl. 48, f. 11–13.*

EXSICCATI: Clements, Crypt. Form. Colo. *130, 302;* N. Am. Fungi *3039;* Fungi Columb. *1931.*

2. **Elvela crispa** (Scop.) Fries, Syst. Myc. **2**: 14. 1822. (PLATE 39, FIG. 2.)

?Elvela pallida Schaeff. Fung. Bavar. **4**: 112. 1770.
Phallus crispus Scop. Fl. Carn. **2**: 475. 1772.
Elvela nivea Schrad. Jour. Bot. Schrad. **2²**: 66. 1799.
Elvela leucophaea Pers. Obs. Myc. **2**: 19. 1799.
Elvela fusca Gill. Champ. Fr. Discom. 9. 1879.

Pileus mitrate or more often saddle-shaped, reflexed and usually irregularly lobed, reaching a diameter of 4–5 cm.; hymenium white, becoming cream or yellowish with age and especially when dried, even or more or less convoluted; stem reaching a length of 6–7 cm. and a diameter of 2–3 cm., deeply fluted, entirely white when fresh, becoming cream or yellowish with age, still darker when dry; asci cylindric or subcylindric, reaching a length of 300 μ and a diameter of 15–18 μ; spores 1-seriate, ellipsoid, containing one large oil-drop, about 12 $\times$ 20 μ; paraphyses enlarged above, where they reach a diameter of 8 μ.

On soil usually in wooded places.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to Alberta and California; also in Europe.

ILLUSTRATIONS: Boud. Ic. Myc. *pl. 225;* Cooke, Mycographia *pl. 41, f. 159;* Gill. Champ. Fr. Discom. *pl. 2, f. 2;* Krombh. Abbild. *pl. 19, f. 27–29;* Massee, Brit. Fungus-Fl. **4**: 188, *f. 17–18;* E. & P. Nat. Pfl. **1¹**: 168, *f. 141, G–H;* Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 5, f. 1;* Bull. Herb. Fr. *pl. 466, f. A;* Minn. Bot. Studies **3**: *pl. 49, f. 10;* Schaeff. Fung. Bavar. *pl. 282.*

EXSICCATI: Fungi Columb. *3329.*

3. **Elvela palustris** Peck, Ann. Rep. N. Y. State Mus. **33**: 31. 1880.

?Elvela pallescens Schaeff. Fung. Bavar. **4**: Ind. 114. 1774.

Pileus widely expanded, the margin reflexed, at first slightly adnate with the stem but at maturity entirely free, reaching a

diameter of 2–3 cm., light-colored and rugose underneath; hymenium even or slightly undulated, the color varying from brown to blackish; stem slender, of nearly uniform diameter throughout its entire length or very slightly enlarged at the base, deeply fluted throughout its entire length, dark, the color similar to that of the hymenium, reaching a length of 4–5 cm. and a diameter of 5–6 mm. at the base; asci cylindric or subcylindric, reaching a length of 200 μ and a diameter of 16 μ; spores ellipsoid, containing one large oil-drop which often breaks up in dried specimens, about 9 $\times$ 18 μ; paraphyses enlarged at their apices, where they reach a diameter of 6–8 μ.

Among mosses and liverworts in swamp.

TYPE LOCALITY: Manilius Center, New York.

DISTRIBUTION: New York; also in Europe?

ILLUSTRATIONS: Ann. Rep. N. Y. State Mus. **33**: *pl. 2, f. 16–18;* ? Schaeff. Fung. Bavar. *pl. 322;* Cooke, Mycographia *pl. 94, f. 341.*

Little is known of this species and it is possible that it is only a slender form of *Elvela Mitra.*

4. **Elvela atra** Oed. Fl. Dan. **9**: 7. 1770.

Elvela nigricans Pers. Obs. Myc. **1**: 72. 1796.
?Leptopodia atra Boud. Hist. Class. Discom. 37. 1907.

Pileus mitrate or more often saddle-shaped, the margin free from the stem, at least when mature and irregularly lobed, the color underneath smoky-brown, becoming almost black when mature, especially when dry; hymenium nearly even or slightly undulated, entirely black; stem slender, slightly enlarged at the base, entirely smoky-black except the base, which is whitish, slightly pruinose or tomentose, reaching a length of 4–5 cm. and a diameter of 2–3 mm.; asci cylindric or subcylindric, reaching a length of 230 μ and a diameter of 16 μ; spores 1-seriate, ellipsoid, containing one large oil-drop, about 12 $\times$ 20 μ; paraphyses strongly enlarged at their apices, where they reach a diameter of 8–10 μ, smoky-brown.

On the ground, usually in woods.

TYPE LOCALITY: Europe.

DISTRIBUTION: Wisconsin to Montana; also in Europe.

ILLUSTRATIONS: Fl. Dan. *pl. 534;* Boud. Ic. Myc. *pl. 238;* Cooke, Mycographia *pl. 43, f. 167;* Gill. Champ. Fr. Discom. *pl. 7;* Krombh. Abbild. *pl. 21, f. 18–20;* Rab. Krypt.-Fl. 1^3: 1173, *f. 8.*

1–3. ELVELA ELASTICA
4–5. ELVELA ELASTICA GIGANTEA

This species although listed separately here scarcely differs from *E. elastica* except in color; whether the species is well founded is difficult to say without more extended observations in the field. This is made difficult through the fact that the plants are not often collected.

5. **Elvela elastica** Bull. Herb. Fr. *pl. 242.* 1785. (PLATE 40.)

?Elvela capucinoides Peck, Bull. N. Y. State Mus. **157**: 27. 1912.
Elvela gracilis Peck, Ann. Rep. N. Y. State Mus. **24**: 94. 1872.
?Elvela fistulosa Alb. & Schw. Consp. Fung. 299. 1805.
?Elvela fuliginosa Pers.; Cooke, Mycographia 92. 1876.
Elvela Stevensii Peck, Bull. Torrey Club **31**: 182. 1904.

Pileus irregularly saddle-shaped, the margin reflexed and usually free from the stem, reaching a diameter of 2–3 cm.; hymenium even or more or less convolute, the color varying from yellowish to dark-brown or nearly black; stem slender, 3–10 mm. thick, reaching a length of 5–10 cm., even or slightly lacunose, never fluted, yellowish, paler than the hymenium; asci cylindric or subcylindric, reaching a length of 200–250 μ and a diameter of 20 μ; spores 1-seriate, ellipsoid, containing one large oil-drop which often becomes indistinct in dried specimens, 10–12 $\times$ 18–20 μ; paraphyses strongly enlarged above, where they reach a diameter of 10 μ, faintly colored.

On the ground, usually in woods.

TYPE LOCALITY: Europe.

DISTRIBUTION: New York to California, Idaho and Texas; also in Europe.

ILLUSTRATIONS: Bull. Herb. Fr. *pl. 242;* Boud. Ic. Myc. *232;* Cooke, Mycographia *pl. 42, f. 163, 164;* Gill. Champ. Fr. Discom. *pl. 11, 12;* Rehm, Krypt.-Fl. 1^3: 1173, *f. 6–7;* Krombh. Abbild. *pl. 21, f. 21;* Pat. Tab. Fung. *f. 100;* E. & P. Nat. Pfl. 1^1: 168, *f. 141, J;* Bull. Lab. Nat. Hist. State Univ. Iowa **6**: *pl. 5, f. 2;* Minn. Bot. Studies **3**: *pl. 49, f. 14, 15.*

EXSICCATI: Underw. & Cook, Illust. Fungi *81;* Clements, Crypt. Form. Colo. *129.*

The large specimens shown on the accompanying plate were collected and photographed by C. G. Lloyd. It is unusually large and might be regarded as a different form.

6. **Elvela Klotzschiana** Corda; Sturm, Deutsch. Fl. **3**: 121. 1837.

?Leptopodia Klotzschiana Boud. Hist. Class. Discom. 37. 1907.

Pileus mitrate, saddle-shaped or irregularly lobed, reaching a diameter of 2 cm., the color underneath bright-yellow or tan;

hymenium even or more or less undulated, dark brownish-black; stem short, slender, of nearly uniform diameter throughout its entire length, or slightly enlarged at the base, reaching a length of 1.5 cm. and a diameter of 3–4 mm., similar in color to the under side of the pileus; asci cylindric or subcylindric, reaching a length of 300 μ and a diameter of 16 μ; spores 1-seriate, ellipsoid, containing one large oil-drop, 10 $\times$ 17–18 μ; paraphyses enlarged above, where they reach a diameter of 8 μ, slightly colored.

On soil, usually in woods.

TYPE LOCALITY: Europe.

DISTRIBUTION: Idaho; also in Europe.

ILLUSTRATION: Sturm, Deutsch. Fl. **3**: *pl. 57.*

7. **Elvela adhaerens** Peck, Bull. N. Y. State Mus. **54**: 956. 1902.

Pileus thin, irregularly reflexed and more or less saddle-shaped, reaching a diameter of 2–3 cm., the margin adhering to the stem, underneath whitish or yellowish; hymenium even or very slightly undulated, whitish to tan, becoming brownish with age; stem slender, reaching a diameter of 5 mm. and a length of 6–8 cm., similar in color to the outside of the pileus; asci cylindric or subcylindric, 8-spored; spores ellipsoid, usually containing one large oil-drop, about 12 $\times$ 18–20 μ; paraphyses strongly thickened at their apices.

On the ground in woods.

TYPE LOCALITY: Bolton, New York.

DISTRIBUTION: New York.

ILLUSTRATION: Bull. N. Y. State Mus. **54**: *pl. L, f. 11–14.*

8. **Elvela californica** Phill. Trans. Linn. Soc. II. **1**: 423. 1880.

Pileus strongly spreading, reflexed or occasionally irregularly saddle-shaped, reaching a diameter of 10 cm., strongly inflated, the substance very thin and papery when dry; hymenium even or irregularly convoluted, light- to dark-brown; stem stout, reaching a diameter of 3 cm. and a length of 10 cm., deeply and irregularly lacunose, with the ribs extending out on the under side of the pileus like the ribs of an umbrella, the entire stem white or occasionally with a pinkish tint, becoming yellowish with age; asci clavate or subcylindric, reaching a length of 200 μ and a diameter of 12 μ; spores 1-seriate or partially 2-seriate above, narrow-ellipsoid, containing two small oil-drops,



ELVELA UMBRACULIFORMIS

about 8–9 × 17 μ; paraphyses rather stout, enlarged above, reaching a diameter of 8 μ at their apices.

On soil, usually in woods.

TYPE LOCALITY: Sierra Nevada Mountains, California.

DISTRIBUTION: California and Nevada.

ILLUSTRATION: Trans. Linn. Soc. II. **1**: *pl. 48*.

EXSICCATI: N. Am. Fungi *2737*.

9. **Elvela umbraculiformis** Seaver, sp. nov. (PLATE 41.)

Pileus inflated and widely expanded and reflexed, the margin free or attached to the stem at various points, reaching a diameter of 10 cm.; hymenium light-brown or tan to dark-brown or brownish-black, even or irregularly undulated to convolute; stem stout, deeply and irregularly lacunose, reaching a length of 4–5 cm. and a diameter of 3–4 cm., the ribs of the stem extending out on the under side of the pileus like the ribs of an umbrella, the color white, whitish or pinkish; asci cylindric or subcylindric, reaching a length of 225 μ and a diameter of 10 μ; spores 1-seriate, each containing two small oil-drops, ellipsoid, the ends narrowed, about 7–8 × 16–17 μ; paraphyses clavate, enlarged at their apices, where they reach a diameter of 10 μ.

On damp soil.

TYPE LOCALITY: Morgan Hill, Yosemite Park, California, elevation 7,000 feet.

DISTRIBUTION: California and Idaho.

This species resembles the preceding but differs in its much shorter stem as well as in other minor points.

10. **Elvela infula** Schaeff. Fung. Bavar. **4**: Ind. 105. 1774. (PLATE 42.)

?Phallus Monacellus Scop. Fl. Carn. ed. 2, **2**: 476. 1772.
Elvela Mitra Schaeff. Fung. Bavar. **4**: Ind. 105. 1774. Not *E. Mitra* L.
Elvela brunnea L. Syst. Nat. 1450. 1796.
Elvela esculenta Pers. Comm. Fung. Bavar. 64. 1800.
Elvela infula Pers. Syn. Fung. 617. 1801.
Elvela rhodopus Krombh. Abbild. **3**: 23. 1834.
Gyromitra esculenta Fries, Summa Veg. Scand. 346. 1849.
Gyromitra infula Quél. Ench. Fung. 272. 1886.
Gyromitra esculenta crispa Peck, Ann. Rep. N. Y. State Mus. **51**: 299. 1898.
Physomitra infula Boud. Hist. Class. Discom. Eu. 35. 1907.
Physomitra esculenta Boud. Hist. Class. Discom. Eu. 35. 1907.

Pileus reaching a diameter of 6–8 cm., reflexed and more or less adnate to the stem, very irregular, mitrate, saddle-shaped or

occasionally subglobose, even or variously contorted or convoluted, the color varying from reddish-brown to dark-brown and occasionally almost black; stem reaching a length of 6–8 cm. and a diameter of 5–15 mm., even or more or less lacunose, never strongly fluted, the color varying from whitish to yellowish or occasionally with a pinkish tint; asci cylindric or subcylindric, reaching a length of 200 μ and a diameter of 12–14 μ, 8-spored; spores 1-seriate or partially 2-seriate, rather narrow-ellipsoid, containing two oil-drops, about 8–12 $\times$ 18–24 μ; paraphyses strongly enlarged at their apices, reaching a diameter of 10 μ.

On the ground or occasionally on rotten wood, usually summer and autumn.

TYPE LOCALITY: Europe.

DISTRIBUTION: Maine to British Columbia and California; also in Europe.

ILLUSTRATIONS: Schaeff. Fung. Bavar. **4**: *pl. 159, 161;* Rab. Krypt.-Fl. **1**[3]: 1174, *f. 1–3;* Boud. Ic. Myc. *pl. 223, 224;* Pers. Champ. Comest. *pl. 4;* Cooke, Hand. Brit. Fungi **2**: 657, *f. 322;* Cooke, Mycographia *pl. 89, f. 328* and *pl. 90, f. 330;* Gill. Champ. Fr. Discom. *pl. 19;* Massee, Brit. Fungus-Fl. **4**: 188, *f. 14;* Phill. Brit. Discom. *pl. 1, f. 2;* E. & P. Nat. Pfl. **1**[1]: 168, *f. 141, B;* Fries, Sv. Aetl. Svamp. *pl. 82* and *83;* Krombh. Abbild. *pl. 20, f. 6–12* and *pl. 21, f. 12–17;* Minn. Bot. Studies **3**: *pl. 49, f. 1–3.*

EXSICCATI: N. Am. Fungi *1267;* Clements, Crypt. Form. Colo. *141.*

The writer regards *Gyromitra esculenta* as only a gyrose form of *Elvela infula*, although there may be some difference of opinion on this point.

11. **Elvela brevissima** Peck, Bull. Torrey Club **30**: 100. 1903.

Pileus irregularly undulated or convolute, with the margin reflexed and usually free, whitish or pallid beneath, reaching a diameter of 1–2 cm.; hymenium dark-brown, becoming blackish when dry; stem very short and rather stout, even, solid, whitish or pallid, reaching a length of 1–1.5 cm. and a diameter of 3–4 mm.; asci cylindric or subcylindric, 8-spored; spores 1-seriate, ellipsoid, usually containing two small oil-drops, about 10–12 $\times$ 15–20 μ; paraphyses slender, enlarged above and colored.

On the ground.

TYPE LOCALITY: California.

DISTRIBUTION: Known only from the type locality.

ELVELA INFULA

12. **Elvela albipes** Fuckel, Symb. Myc. 334. 1869.

Elvela Hegani Copeland, Ann. Myc. **2**: 510. 1904.

Pileus lobate, deflexed, the lobes subreniform, reaching a diameter of 2 cm. or more, whitish or yellowish outside; hymenium brown, becoming blackish with age; stem white or whitish, subpubescent, even or scarcely lacunose, reaching a diameter of 1 cm. and a length of 3–4 cm.; asci cylindric or subcylindric, reaching a length of 300 μ and a diameter of 16–18 μ; spores 1-seriate, ellipsoid, each containing one large oil-drop, 12–14 $\times$ 20–22 μ; paraphyses enlarged above, where they reach a diameter of 7–8 μ, yellowish.

On damp soil.

TYPE LOCALITY: Europe.

DISTRIBUTION: California and Idaho; also in Europe.

ILLUSTRATIONS: Fuckel, Symb. Myc. *pl. 5, f. 2;* Boud. Ic. Myc. *pl. 231;* Ann. Myc. **3**: *pl. 12, f. 8, 9.*

13. **Elvela caroliniana** (Bosc) Nees, Syst. Pilze 176. 1817.

Morchella caroliniana Bosc Ges. Nat. Freunde Berlin Mag. **5**: 86. 1811.
?Elvela costata Schw. Schr. Nat. Ges. Leipzig **1**: 114. 1822.
Elvela Gigas Krombh. Abbild. **3**: 28. 1834.
Mitrophora caroliniana Lév. Ann. Sci. Nat. III. **5**: 250. 1846.
Gyromitra curtipes Fries, Sv. Aetl. Svamp. 34. 1861.
Gyromitra caroliniana Fries, Ofv. Sv. Vet.-Akad. Förh. **28**: 173. 1871.
Gyromitra Gigas Cooke, Mycographia 191. 1878.

Pileus irregularly lobed and plicate and in places more or less gyrose, the ridges often anastomosing and giving to the surface a reticulate appearance, whitish underneath, the margin adhering to the stem in various places, reaching a diameter of 5–12 cm.; hymenium brown to brownish-black; stem rather short and stout, more or less enlarged at the base, reaching a diameter of 2–5 cm. and a length of 8–10 cm., rather deeply lacunose, the color white or whitish; asci clavate or subcylindric; spores 1-seriate, narrow-ellipsoid and often apiculate, usually containing one large oil-drop and often two smaller ones, at first smooth, becoming sculptured; spore-sculpturing assuming the form of minute warts or occasionally reticulations; paraphyses thickened above and colored.

On the ground.

TYPE LOCALITY: Carolina.

DISTRIBUTION: West Virginia to Colorado; also in Europe.

ILLUSTRATIONS: Krombh. Abbild. *pl. 20, f. 1–5;* Boud. Ic. Myc. *pl. 221;* Hard, Mush. *f. 419* and (? *f. 418* as *G. esculenta*); Ges. Nat. Freunde Berlin Mag. **5**: *pl. 5, f. 6;* Cooke, Mycographia *pl. 90, f. 330.*

14. **Elvela Underwoodii** Seaver, nom. nov. (PLATE 43.)

Gyromitra brunnea Underw. Proc. Ind. Acad. Sci. **1893**: 33. 1894.

Pileus much contorted, irregularly lobed and plicate, in places faintly marked into areas by indistinct anastomosing ridges, adhering to the stem at various points, whitish underneath, reaching a diameter of 5–12 cm. in the widest point; hymenium chocolate-brown (lighter when covered by leaves); stem reaching a length of 8–13 cm. and a diameter of 2–5 cm., rarely slightly lacunose, hollow or loosely stuffed, white; asci cylindric or subcylindric; spores ellipsoid, 14 × 28–30 μ, usually containing two rather large oil-drops, becoming sculptured; spore-sculpturing consisting of small warts or faint reticulations; paraphyses slender, enlarged above.

In rich woods on leaf-mould.

TYPE LOCALITY: Putnam County, Indiana.

DISTRIBUTION: Indiana and Ohio.

ILLUSTRATION: Hard, Mush. *f. 419.*

The name assigned by Underwood is untenable in this genus since the *Elvela brunnea* was used by Linnaeus for an entirely different species. The name *Elvela Underwoodii* is therefore adopted.

15. **Elvela sphaerospora** Peck, Ann. Rep. N. Y. State Mus. **27**: 106. 1875.

Gyromitra sphaerospora Sacc. Syll. Fung. **8**: 16. 1889.

Pileus rather widely spreading, the margin quite evenly reflexed and usually free from the stem, whitish underneath and slightly tomentose, reaching a diameter of 6–8 cm.; hymenium nearly even or slightly undulated, the color varying from brown to brownish-black; stem stout, reaching a diameter of 2–3 cm. and a length of 4–8 cm., whitish or slightly pinkish, deeply lacunose or fluted, the ribs extending up underneath the pileus like the ribs of an umbrella; asci cylindric or subcylindric, reaching a length of 150 μ and a diameter of 10–12 μ; spores 1-seriate, globose, reaching a diameter of 10 μ; paraphyses slightly enlarged above, where they reach a diameter of 4 μ.

ELVELA UNDERWOODII

On old stumps in woods.

TYPE LOCALITY: Indian Lake, New York.

DISTRIBUTION: Vermont to Wisconsin and Manitoba.

ILLUSTRATION: Ann. Rep. N. Y. State Mus. **51**: *pl. B, f. 21–25.*

DOUBTFUL SPECIES

Elvela pileata Clements, Bull. Torrey Club **30**: 94. 1903. The description of this species suggests that it may be a *Verpa*. No specimen has been seen.

5. **UNDERWOODIA** Peck, Ann. Rep. N. Y. State Mus. **43**: 32. 1890.

Pileus fleshy, more or less elongated or columnar, entirely covering the stem; hymenium covering the entire outer surface of the pileus, even or undulated; stem externally lacunose or fluted, internally containing several longitudinal cavities; asci cylindric above, 8-spored; paraphyses slender below, clavate above.

Type species, *Underwoodia columnaris* Peck.

1. **Underwoodia columnaris** Peck, Ann. Rep. N. Y. State Mus. **43**: 32. 1890. (PLATE 44.)

Pileus clavate, columnar, or slightly tapering above, straight or curved and often horn-shaped, entirely overspreading the stem; hymenium at first light-colored, becoming brownish, the entire fruiting body like the stem of an *Elvela* entirely overspread and surrounded by the pileus, reaching a height of 10 cm. and a diameter of 2–3 cm.; asci reaching a length of 350 μ and a diameter of 20 μ, tapering below into a stem-like base with an abrupt enlargement at the extreme base; spores 1-seriate or occasionally crowded, ellipsoid, at first smooth, becoming sculptured, 12–14 $\times$ 25–27 μ; spore-sculpturing taking the form of rather coarse warts or small tubercles; paraphyses rather strongly thickened above, brownish.

On soil among leaves.

TYPE LOCALITY: Kirkville, New York.

DISTRIBUTION: New York to Michigan and Manitoba.

ILLUSTRATION: Ann. Rep. N. Y. State Mus. **43**: *pl. 4, f. 1–4.*

Since this manuscript was submitted to the printers, excellent specimens of *Underwoodia columnaris* and *Elvela sphaerospora* have been received from G. R. Bisby of Winnipeg, Manitoba.

These additional specimens have materially extended the range of the species northward.

EXCLUDED GENUS

Spragueola Massee, Jour. Bot. **34**: 149. 1896. This genus was founded by Massee on *Spragueola americana*, which according to Durand is *Mitrula irregularis* (Peck) Durand, a species which becomes irregularly distorted as indicated by its name.

BIBLIOGRAPHICAL NOTE

Since most of the citations in this work are familiar to mycologists, it hardly seems necessary at this time to append a complete bibliography. If a second volume is eventually published on the Inoperculates, it would be much more fitting that the bibliography, if one is thought necessary, should appear at that time.



UNDERWOODIA COLUMNARIS

EXPLANATION OF PLATES

The plates accompanying this work consist of colored drawings made from living plants, photographs, and drawings of spores and other microscopic characters. All spore drawings are made with the aid of the camera lucida and are drawn to a common scale. The reduction is also approximately equal so that the proportionate sizes are kept throughout. Other microscopic characters such as the hairs from the apothecia are also drawn with the aid of the camera lucida but occasionally it became necessary to use a lower magnification. If desired, the exact magnification can be determined by comparing reproductions with sizes indicated in descriptions. An especial effort has been made to keep drawings of spores approximately proportionate throughout the work.

Since the plates are distributed through the body of the work, instead of being bunched at the end, as a matter of convenience the page opposite which the plate appears is placed in parenthesis directly after the plate number.

In order to save space and bring the various characters together, a combination of photographs and drawings has been used. Where not otherwise indicated, the photographs of apothecia and ascophores are approximately natural size, although they may not indicate the maximum size which the apothecia may attain.

Frontispiece

Phillipsia Chardoniana. Vertical view of two plants showing the expanded apothecia, about natural size. Drawn and colored from life by Mr. Mario Brau of the Department of Agriculture of Porto Rico and the plate furnished by the courtesy of that Department.

Plate 1 (p. 48)

Upper figure (1). *Pseudoplectania vogesiaca.* Three apothecia, about natural size. Also drawing of a portion of an ascus with spores; one of the branching and hooked paraphyses; and a portion of one of the hairs from the outside of the apothecium.

Lower figure (2). *Pseudoplectania nigrella.* Several apothecia, about natural size, showing the habitat among *Sphagnum* moss. At the left, drawing of a portion of an ascus with spores and a paraphysis. Below, two of the coiled and twisted hairs from the outside of the apothecia.

Plate 2 (p. 52)

Figure 1. *Boudiera areolata.* Portion of an ascus with spores and a tip of one of the much thickened paraphyses. Drawn with the aid of the camera lucida from material collected in Maine by Dr. Roland Thaxter and compared with authentic material collected in Europe. The spores are marked with long spines or interrupted reticulations.

Figure 2. *Boudiera echinulata.* Drawn from material collected in Iowa and originally described as *Sphaerosoma echinulatum* by the author.

Spores scarcely differ from those of the preceding except that they are slightly larger.

FIGURE 3. *Sphaerosoma fuscescens.* A portion of an ascus with spores, showing the definite reticulations in contrast with the indefinite reticulations of the two preceding.

PLATE 3 (p. 64)

Drawings of a portion of the ascus and spores of various species of *Lamprospora:* 1, *L. Crec'hqueraultii;* 2, *L. Crouani;* 3, *L. areolata;* 4, *L. spinulosa;* 5, *L. tuberculatella;* 6, *L. tuberculata;* 7, *L. ascoboloides;* 8, *L. annulata;* 9, *L. Wrightii;* 10, *L. lobata.*

PLATE 4 (p. 78)

UPPER FIGURE (1). *Pithya Cupressi.* Photographs of a large number of apothecia on the twigs of Bermuda cedar, *Juniperus bermudiana,* collected in Bermuda by the writer. Also drawing of a portion of an ascus with spores, and the tips of two of the slender paraphyses. Lower right-hand corner, the end of an ascus showing the operculum.

CENTRAL FIGURE (2). *Pithya lacunosa.* Photograph of the type collected in Newfoundland by Waghorne. Also drawing of a portion of the ascus with spores, and a tip of a paraphysis.

LOWER FIGURE (3). *Pithya pithya.* Photographs of several apothecia on some fir. Also drawing of a portion of an ascus with spores, and a tip of a paraphysis.

PLATE 5 (p. 80)

FIGURES 1–5. *Ascodesmis microscopica.* 1, cluster of immature asci and paraphyses; 2, asci and paraphyses with spores, in various stages of maturity; 3, ascus with immature spores; 4, ascus with mature spores; 5, mature spores removed from the ascus, showing the regular character of the reticulations.

FIGURES 6–9. *Ascodesmis porcina.* 6, cluster of asci and paraphyses in various stages of development; 7, isolated ascus with immature spores; 8, ascus with mature spores and a tip of one of the paraphyses; 9, mature spores removed from ascus, showing character of the sculpturing.

PLATE 6 (p. 86)

Ascobolus magnificus. Photographs of a large number of apothecia in various stages of development, the photographs made by Dr. B. O. Dodge, the author of the species. Above, a drawing of a portion of an ascus with several spores, showing the characteristic marking.

PLATE 7 (p. 90)

Drawings of the asci and spores of various species of *Ascobolus:* 1, *A. viridis.* Drawing of a portion of an ascus with spores for comparison with *A. geophilus,* the American form with which it has been confused; 2, *A. geophilus.* Drawing of a portion of an ascus with a part of the spores undeveloped; also isolated spores in various stages of development; 3, *A. subglobosus.* Drawing of a portion of an ascus with spores and several isolated

spores in various stages of development; 4, *A. carbonarius.* Drawing of a portion of an ascus with isolated spores; 5, *A. immersus.* Drawing of an ascus with spores.

PLATE 8 (p. 98)

UPPER FIGURE (1). *Aleuria aurantia.* Photograph of a number of apothecia, about natural size. At the right, a drawing of an ascus with mature spores showing reticulate markings and a paraphysis with rather strongly enlarged tip. Below, two immature spores.

LOWER FIGURE (2). *Aleurina atrovinosa.* Photograph of a cluster of apothecia, about natural size. At the left, a drawing of an ascus with mature spores showing irregular reticulations and a paraphysis with enlarged tip. Above, two immature spores.

PLATE 9 (p. 102)

UPPER FIGURE (1). *Aleuria rutilans.* Photograph of several apothecia at different stages in their development and showing habitat among mosses. Above, drawing of a mature and an immature spore, the former showing the regular and fine reticulate markings.

LOWER FIGURE (2). *Aleuria rhenana.* Photograph of several apothecia showing habitat in coniferous woods and dense mycelial growth which binds the substratum together. Above at the right, a mature and an immature spore, the former showing the typical reticulate markings.

PLATE 10 (p. 110)

Pyronema omphalodes. Photograph of a mass of mycelium and apothecia, showing the natural habitat on charcoal (a). Upper right-hand corner, a germinating ascospore showing the very coarse mycelium (c). Lower right-hand corner, cluster of oögonia with trichogynes (d). Lower left-hand corner, antheridium and oögonium with the connecting trichogyne (e). Upper left-hand corner, an ascus with spores and a paraphysis (b).

PLATE 11 (p. 120)

Drawings of asci and spores of various species of *Ascophanus:* 1, *A. lacteus;* 2, *A. bermudiensis;* 3, *A. minutissimus* (later referred to *A. microsporus?*); 4, *A. sarcobius;* 5, *A. argentius;* 6, *A. glumarum* = *Humarina glumarum;* 7, *A. Aurora;* 8, *A. Holmskjoldii;* 9, *A. microsporus* with paraphysis showing enlarged tip; 10, *A. granuliformis;* 11, *A. cinereus;* 12, *A. granulatus;* 13, *A. isabellinus.*

PLATE 12 (p. 144)

Drawings of asci and spores of various species: 1, *Ryparobius sexdecimsporus;* 2–3, *Streptotheca woolhopensis,* the latter showing the method of dehiscence of the ascus; 4, *Streptotheca obscura;* 5, *Ryparobius polysporus;* 6, *Ryparobius monascus;* 7, *Thecotheus Pelletieri,* showing the large size of the asci and spores.

PLATE 13 (p. 150)

Sepultaria Longii. Upper figure, a vertical view of a cluster of apothecia which make their appearance as a series of holes in the surface of the ground. Lower figure, a section through such a cluster, giving some idea of the subterranean habits of the species. At the left, a drawing of an ascus with spores and a paraphysis. In the center, a portion of the mycelium which clothes the outside of the apothecia and binds the soil together.

PLATE 14 (p. 160)

UPPER FIGURE (1). Photograph of several apothecia of *Patella albida*, made by Dr. F. E. Clements under the name of *Lachnea chaetoloma.* Upper right-hand corner, sketch of a section through one of the apothecia, showing general form of cup, also drawings of hairs from outside of apothecia. Lower left-hand corner, drawing of an ascus with spores.

LOWER FIGURE (2). *Patella scutellata.* Several apothecia on the bark of a dead stump. Above, a sketch of a section through an apothecium. Right, a drawing of a hair from outside of apothecium. Left, a drawing of an ascus with spores and a paraphysis with its enlarged tip.

PLATE 15 (p. 176)

UPPER FIGURE (1). *Patella gregaria.* Photograph of group of apothecia, showing natural habitat under pine trees. Right, drawings of hairs from outside of apothecia. Left, drawing of a portion of an ascus with spores and a paraphysis.

LOWER FIGURE (2). *Patella melaloma.* Photograph of a group of apothecia, showing natural habitat on charcoal. Right, drawing of portion of an ascus with spores and a paraphysis. Left, cluster of club-shaped hairs from outside of apothecia.

PLATE 16 (p. 180)

Wynnea americana. Photograph of a sclerotium with a branched apothecium. Right, drawing of a portion of an ascus with spores, showing the longitudinal striations. Left, an isolated spore showing its unequal-sided character. Above, sketch of the end of an ascus, showing the eccentricity of the ascostome.

PLATE 17 (p. 184)

UPPER FIGURE (1). *Phillipsia gigantea.* Three apothecia photographed from dried specimens but about natural size. Also drawing of a portion of an ascus with spores and a paraphysis.

CENTRAL FIGURE (2). *Phillipsia domingensis.* Several apothecia on a dead stick. Spores apparently identical with those of the preceding.

LOWER FIGURE (3). *Cookeina Colensoi.* Photograph of a number of apothecia on moss-covered bark, made from dried specimens, which, however, shrink little in drying. Left, drawing of a portion of an ascus with spores, showing characteristic striations. Also the tip of a paraphysis.

PLATE 18 (p. 188)

UPPER FIGURE (1). *Cookeina sulcipes.* Photograph of several apothecia on dead wood, showing variation in the length of the stem, from dried specimens which shrink little in drying. Right, drawing of a portion of an ascus with spores, showing characteristic striations. Also empty ascus showing eccentric position of ascostome and operculum and the abrupt tail-like stem at the base of the ascus. Near the center, a drawing of one of the fasciculate hairs to be found about the margin of the cup.

LOWER FIGURE (2). *Cookeina Tricholoma.* Photograph of a cluster of apothecia on moss-covered bark, made from dried specimens which shrink little in drying. Near the left, a cluster of apothecia showing the variation in the length of the stem; also the conspicuous hairs which clothe the outside of the cup. Left, a drawing of a portion of an ascus with spores, showing characteristic striations. Right, drawing of tip of one of the long fasciculate hairs from the outside of the apothecium.

PLATE 19 (p. 190)

UPPER FIGURE (1). *Plectania coccinea.* Photograph of two apothecia growing from a partially buried stick. Photograph by C. G. Lloyd. Left, drawing of a portion of an ascus with spores. Upper right-hand corner, drawing of hairs from the outside of the apothecium.

LOWER FIGURE (2). *Plectania hiemalis.* Photograph of two apothecia, showing long stems and crenate margins. Photograph by C. G. Lloyd. Lower right-hand corner, drawing of a portion of an ascus with spores. Upper left-hand corner, tip of ascus showing the eccentricity of the ascostome and operculum. Near the center, two hairs from outside of the apothecium.

PLATE 20 (p. 192)

UPPER FIGURE (1). *Plectania floccosa.* Cluster of apothecia, showing long stems and densely hairy exteriors. Also a drawing of a portion of an ascus with spores and the tip of one of the very long hairs from the outside of the apothecium.

LOWER FIGURE (2). *Plectania occidentalis.* Photographs of several apothecia from dead sticks, showing the variation in the length of the stems and the comparatively smooth exterior of the apothecia. Near the center, drawing of a portion of an ascus with spores and a slender paraphysis. Lower right-hand corner, drawing of hairs from exterior of cups.

PLATE 21 (p. 198)

Urnula Craterium. Photograph of a cluster of apothecia from a partially buried stick, with their notched margins. Photograph by C. G. Lloyd. Right, drawing of a portion of an ascus with spores and tips of two paraphyses. Upper left-hand corner, drawing of a portion of hair from exterior of cup. Below, three isolated spores.

PLATE 22 (p. 200)

Urnula Geaster. Photographs of several apothecia in different stages of development, showing the *Geaster*-like method of dehiscence of the apothecia.

Near the center, drawing of a portion of an ascus with spores. Lower left-hand corner, one spore isolated. Photograph by Dr. W. H. Long.

PLATE 23 (p. 202)

Paxina Acetabulum. Upper figure, photograph of an isolated apothecium, showing the vein-like markings on the outside of the cup. Also drawing of an ascus with spores and the tip of a paraphysis. Lower figure, two apothecia showing habitat. Lower figure from photograph by Dr. F. E. Clements.

PLATE 24 (p. 204)

UPPER FIGURE (1). *Paxina hispida.* Photograph of several apothecia, showing long stems. Left, drawing of an ascus with spores and a paraphysis with its enlarged tip. Right, two spores isolated.

LOWER FIGURE (2). *Paxina sulcata.* Photograph of apothecium, showing thick, corrugated stem. Lower right-hand corner, drawing of an ascus with spores and the tip of a paraphysis. Left, two spores isolated.

PLATE 25 (p. 210)

UPPER FIGURE (1). *Paxina fusicarpa.* Photograph of a cluster of apothecia, showing habitat on the ground. Left, drawing of a portion of an ascus with spores and the tip of a paraphysis. Right, drawing of a hair from outside of apothecium; also one spore isolated.

LOWER FIGURE (2). *Paxina semitosta.* Photograph of a cluster of apothecia, showing comparatively large cups with short, thick, furrowed stems. Right, drawing of a portion of an ascus with spores; also the tip of a paraphysis. Left, drawings of portions of hairs from outside of cups.

PLATE 26 (p. 212)

UPPER FIGURE (1). *Geopyxis cupularis.* Photograph of several apothecia. Photograph by W. R. Fisher. Left, drawing of ascus with spores; also tip of paraphysis.

LOWER FIGURE (2). *Peziza domiciliana.* Photograph of several apothecia in different stages of development, showing the short, stem-like base in young specimens. Left, drawing of a portion of an ascus with spores; also tip of a paraphysis.

PLATE 27 (p. 216)

Rhizina inflata. Upper figure, photograph of a section through one of the apothecia, showing expanded hymenium and root-like attachment. Lower figure, vertical views of apothecia. Photographs by Dr. J. R. Weir. Left, drawing of a portion of an ascus with spores; also tips of paraphyses. Center, drawing of one of hairs from hymenium. Right, two spores, showing apiculate ends.

PLATE 28 (p. 218)

UPPER FIGURE (1). *Discina convoluta.* Photograph of type specimen, showing two apothecia with their convoluted hymenia. Right, drawing of a

portion of an ascus with spores; also the tip of a paraphysis. Left, drawing of one spore isolated, showing peculiar truncate ends.

LOWER FIGURE (2). *Discina ancilis.* Photographs of two apothecia with side and vertical view. Photographs by Dr. J. R. Weir. Center, drawing of a portion of an ascus with spores; also the tip of a paraphysis.

PLATE 29 (p. 224)

UPPER FIGURE (1). *Peziza pustulata.* Photograph of group of apothecia, showing habitat on charcoal. Left, drawing of an ascus with spores; also the tip of a paraphysis.

LOWER FIGURE (2). *Peziza vesiculosa.* Photograph of a cluster of apothecia. Right, drawing of a portion of an ascus with spores.

PLATE 30 (p. 226)

UPPER FIGURE (1). *Peziza proteana.* Photograph of several apothecia, showing habitat on old burnt place which has been subsequently overrun with mosses. Right, drawing of a portion of an ascus with spores; also tip of a paraphysis.

LOWER FIGURE (2). *Peziza violacea.* Photograph of several apothecia, showing habitat on charcoal. Left, drawing of a portion of an ascus with spores; also tips of paraphyses. Note similarity in size and markings of the spores of this and the preceding.

PLATE 31 (p. 228)

Peziza venosa. Upper figure, an expanded apothecium, showing the vein-like folds characteristic of the hymenium. Lower figure, a younger specimen in which the veins are not apparent. Left, drawing of a portion of an ascus with spores; also a paraphysis. Right, an isolated spore.

PLATE 32 (p. 230)

Peziza repanda. Photographs of a number of apothecia in different stages of development, on rotten wood which has been partially overrun with mosses. Left, drawing of a portion of an ascus with spores; also tip of a paraphysis. Center, drawing of tip of an ascus, showing operculum.

PLATE 33 (p. 232)

Peziza sylvestris. Photographs of a number of apothecia in different stages of development, and about natural size, although they are often much larger. Right, drawing of an ascus with spores; also tip of paraphysis. Center, drawing of two spores isolated.

PLATE 34 (p. 234)

Peziza clypeata. Photographs of a number of apothecia at different stages of development, from a rotten log which was partially overgrown with mosses. Right, drawing of a portion of an ascus with spores; also tip of one of the stout paraphyses. Left, drawing of one spore isolated. Note the unusually large size of the spores of this species.

PLATE 35 (p. 238)

Morchella crassipes. Photograph of two ascophores, showing the enlarged bases and rather thin, almost sharp ribs. Lower right-hand corner, a germinating spore. Near center, drawing of an ascus with spores; also a part of one of the stout paraphyses.

PLATE 36 (p. 240)

UPPER FIGURE (1). *Morchella hybrida.* Photograph of three ascophores of different ages. The stems elongate much as the plants mature. Near center, drawing of a portion of an ascus with spores and an isolated spore.

LOWER FIGURE (2). *Morchella esculenta.* Photograph of two ascophores, one bent over. Upper left-hand corner, drawing of a portion of an ascus with spores; also the tip of one of the stout paraphyses and an isolated spore.

PLATE 37 (p. 242)

Durandiomyces Phillipsii. Photograph of the ascophore which appears like a compound apothecium. Photograph obtained from the late Dr. E. J. Durand. Right, drawing of an ascus with spores; also a paraphysis. Note the similarity of the spores to those of *Peziza proteana* and *Peziza violacea* (PLATE 29).

PLATE 38 (p. 244)

UPPER FIGURE (1). *Verpa bohemica.* Photograph of three ascophores from dried specimens. Near center, drawing of a portion of an ascus with spores; also tip of paraphysis and sketch of section through pileus.

LOWER FIGURE (2). *Verpa conica.* Photograph of three ascophores from fresh material. Left, drawing of portion of an ascus with spores; also tip of a paraphysis.

PLATE 39 (p. 246)

UPPER FIGURE (1). *Elvela Mitra.* Photograph of three ascophores from dried material. Also drawing of a portion of an ascus with spores and tip of a paraphysis.

LOWER FIGURE (2). *Elvela crispa.* Photograph of several ascophores at different stages of development, from dried material. Also drawing of a portion of an ascus with spores and tip of a paraphysis.

PLATE 40 (p. 248)

FIGURES 1–3. *Elvela elastica.* Photographs of ascophores in different stages of development. Right, drawing of an ascus with spores; also a paraphysis and an isolated spore.

FIGURES 4–5. *Elvela elastica gigantea.* Unusually large specimens of this species which might be regarded as a different form. Photographed by C. G. Lloyd.

PLATE 41 (p. 250)

Elvela umbraculiformis. Photographs of two ascophores, showing the large pilei and the short, corrugated stems. Left, drawing of an ascus with

spores; also a paraphysis. Right, isolated spores. Lower right-hand corner, a bit of a hair from outside of stem.

PLATE 42 (p. 252)

Elvela infula. Photographs of ascophores at different stages of development and showing different degrees of gyrosity. Center, drawing of a portion of an ascus with spores; also tip of a paraphysis. Right, two isolated spores.

PLATE 43 (p. 254)

Elvela Underwoodia. Photograph of a large ascophore by C. G. Lloyd. Left, drawing of a portion of an ascus with spores; also tip of a paraphysis. Lower right-hand corner, drawing of a mature and an immature spore, showing the faintly reticulate character of the markings.

PLATE 44 (p. 256)

Underwoodia columnaris. Photograph of a double ascophore, consisting of a corrugated stem overspread with the hymenium; also a sketch of a section of the ascophore. Left, drawing of an ascus with spores; also tips of several paraphyses.

PLATE 45 (p. 266)

1, *Lamprospora haemastigma;* 2–3, *Lamprospora Crouani;* 4, *Lamprospora discoidea;* 5–6, *Lamprospora spinulosa;* 7, *Lamprospora Constellatio;* 8, *Lamprospora Crec'hqueraultii;* 9–10, *Lamprospora tuberculatella;* 11–12, *Patella albocincta;* 13–14, *Patella gregaria;* 15–16, *Patella ricciophila;* 17, *Melastiza Charteri;* 18, *Patella coprinaria;* 19, *Patella scutellata;* 20, *Peziza violacea;* 21, *Ascobolus stercorarius;* 22, *Aleuria rutilans;* 23, *Plectania occidentalis.*

PEZIZACEAE

INDEX TO RECOGNIZED GENERA

Aleuria, 96
Aleurina, 101
Ascobolus, 81
Ascodesmis, 79
Ascophanus, 111
Boudiera, 51
Bulgaria, 194
Cookeina, 187
Cubonia, 80
Discina, 216
Durandiomyces, 242
Elvela, 245
Geopyxis, 211
Humarina, 121
Lamprospora, 54
Lasiobolus, 155
Melastiza, 103
Morchella, 237
Patella, 156
Paxina, 201
Perrotia, 154
Peziza, 219
Phillipsia, 182
Pithya, 76
Plectania, 190
Pseudombrophila, 141
Pseudopithyella, 153
Pseudoplectania, 47
Psilopezia, 105
Pyronema, 109
Rhizina, 215
Ryparobius, 144
Saccobolus, 92
Sarcosphaera, 235
Scodellina, 184
Sepultaria, 148
Sphaerosoma, 51
Sphaerospora, 43
Streptotheca, 142
Thecotheus, 147
Underwoodia, 255
Urnula, 198
Verpa, 243
Wynnea, 180

INDEX TO GENERA AND SPECIES

(*Synonyms in italics*)

Acetabula, 201
Acetabulum, 202
ancilis, 216
Barlae, 205
Calyx, 204
Dupainii, 207
leucomelas, 204
macropus, 203
olivacea, 211
radicata, 214
sulcata, 204
venosa, 229
vulgaris, 202
Aleuria, 40, **96**, 101, 219
abietina, 228
Acetabulum, 202
ancilis, 216
apiculata, 217
araneosa, 130
aurantia, **96**–98
auricula, 185
auriflava, 62
Aurora, 115
badia, 221
bicucullata, 96, **99**
Boltoni, 226
brunneoatra, 222
Catinus, 213
cestrica, 96, **98**
Constellatio, 69
convexula, 128
corallina, 137
Corium, 208
Crouani, 56

cupularis, 212
Emileia, 224
eximia, 235
fulgens, 50
glumarum, 133
granulata, 116
hepatica, 139
humicola, 213
humosa, 129
ionella, 138
leiocarpa, 73
leporina, 185
leucoloma, 129
leucomelas, 204
macropus, 205
melaloma, 167
ollaris, 135
omphalodes, 109
onotica, 185
proteana, 227
proteana sparassoides, 242
pustulata, 224
repanda, 231
rhenana, 96, **99**
rutilans, 96, **100**
subhirsuta, 170
succosa, 223
sylvestris, 233
tetraspora, 134
trachycarpa, 71
umbrina, 224
venosa, 229
vesiculosa, 231
violacea, 226
wisconsinensis, 96, **98**
Aleurina, 40, **101**
aquehongensis, 101, **102**
atrovinosa, 101
Lloydiana, 98
retiderma, 101
vinacea, 222
Angelina rufescens, 92
Anthopeziza, 190
Winteri, 193
Anthracobia melaloma, 168
Ascobolus, 10, 16, 26, 40, 55, 60, 79, **81**, 91, 140, 142
aerugineus, 86
albinus, 81, **89**
americanus, 81, **85**
argenteus, 114
atrofuscus, 87, 92
Boudieri, 85
brunneus, 92
candidus, 81, **86**
caninus, 79
carbonarius, 8, 12, 25, 26, 81, **87**, 90, 92
carbonicola, 87
carneus, 115
Cesatii, 121
ciliatus, 155
cinereus, 118
coccineus, 137
conglomeratus, 92
Cookei, 145
Crec'hqueraultii, 62
crenulatus, 86
Crouani, 56, 142
crustaceus, 145
cubensis, 92
depauperatus, 95
diversisporus, 155
epimyces, 82, **91**
furfuraceus, 8, 82
geophilus, 81, **88**, 92
gigasporus, 83
glaber, 81, **83**, 85
grœnulatus, 116
granuliformis, 117
hyperboreus, 79
immersus, 14, 81, **83**
incanus, 112
Kerverni, 93
lacteus, 114
Leveillei, 81, **85**
Leveillei americanus, 85
macrosporus, 83
magnificus, 9, 12, 81, **87**
major, 92
microscopicus, 79
microsporus, 120
miniatus, 54, 56
niveus, 79, 147
obscurus, 96
ochraceus, 117
papillatus, 155
Pelletieri, 148

pezizoides, 82
pilosus, 155
polysporus, 145, 146
pulcherrimus, 172
purpurascens, 92
pusillus, 92
raripilus, 173
saccharinus, 115
sexdecimsporus, 144
stercorarius, 81, **82**, 92
stercorarius retisporus, 82
striisporus, 82, **90**
subglobosus, 81, **89**
testaceus, 125
tetrasporus, 134
Trifolii, 92
vinosus, 85, **92**
viridis, 88, 89, **92**
viridulus, 81, **86**, 91, 92
Winteri, 81, **84**
woolhopensis, 143
xylophilus, 82, **90**
Ascodesmis, 39, 54, **79**
microscopica, 79
nigricans, 79
porcina, 25, 79, **80**
reticulata, 79
Ascophanus, 16, 41, **111**
argenteus, 111, **114**
Aurora, 111, **115**
bermudensis, 111, **113**
carneus, 111, **115**, 116, 126, 127
carneus difformis, 115
carneus saccharinus, 115
carneus testaceus, 125
Cesatii, 112, **121**
cinerellus, 112, **120**
cinereus, 111, **118**
Coemansii, 120
crustaceus, 118
flavus, 117
gallinaceus, 111, **118**
glaucellus, 112, **121**
granulatus, 111, **116**, 141
granuliformis, 111, **117**
hepaticus, 139
Holmskjoldii, 111, **112**
humosoides, 115
incanus, 112
isabellinus, 111, **112**
lacteus, 111, **114**
lilacinus, 111, **120**
microsporus, 112, **120**
minutissimus, 120
ochraceus, 111, **117**
pallens, 124
pilosus, 155
ruber, 156
saccharinus, 115
sarcobius, 111, **113**
sexdecimsporus, 144
subgranuliformis, 117
testaceus, 125
tetraonalis, 111, **119**
vicinus, 111, **114**
Ascozonus, 142
Boudieri, 142
Crouani, 142
oligoascos, 146
Barlaea, 34, 54
amethystina, 67
areolata, 52
asperella, 62
carbonaria, 67
cinnabarina, 58
Constellatio minuta, 69
convexella, 68
Crec'hqueraultii, 62
discoidea, 70
exasperata, 75
fulgens, 50
fuliginea, 76
gemmea, 70
hinnulea, 46
lacunosa, 77
laeterubra, 69
lobata, 74
miniata, 57
modesta, 62
Polytrichi, 75
schizospora, 46
verrucosa, 65
Wrightii, 66
Barlaeina, 34, 54
carbonicola, 67
Constellatio, 69
Constellatio minuta, 69
Crec'hqueraultii, 62

dictydiola, 58
discoidea, 70
laeterubra, 69
Botrytis, 22
Boudiera, 11, 39, **51**
areolata, 52, **54**
canina, 79
Claussenii, 79
echinulata, 52
marginata, 53
microscopica, 79
Bulgaria, 35, 42, **194**, 198
bicolor, 233
carbonaria, 71
deligata, 107
globosa, 5, **195**, 196
inquinans, 195
melastoma, 195, **197**
platydiscus, 196
rufa, 195, **196**
spongiosa, 48
Burcardia, 194
globosa, 195
Burckhardia globosa, 195
Caloscypha, 47
fulgens, 50
Calycina melastoma, 197
Catinella, 140
Cenangium, 198
Craterium, 199
Cheilymenia, 157
coprinaria, 171
dalmeniensis, 170
fimetaria, 170
pulcherrima, 172
stercorea, 169
subhirsuta, 170
theleboloides, 170
Chlorosplenium, 91
epimyces, 91
striisporum, 90
Chorioactis, 198
Geaster, 200
Ciboria, 214
Ciliaria, 156
asperior, 44
confusa, 46
Lusatiae, 162
paludosa, 174
scutellata, 159
setosa, 166
trechispora, 44
umbrorum, 161
Cookeina, 17–19, 42, **187**
Afzelii, 188
Colensoi, 18, 19, 187, **190**
Hindsii, 188
sulcipes, 187, **188**
tetraspora, 187, **189**
Tricholoma, 187
Coprobia, 111
granulata, 116
Craterium microcrater, 199
Crouania, 34, 54
asperella, 62
carbonaria, 67
cinnabarina, 58
humosa, 69
miniata, 56
Cubonia, 39, 80
Curreyella, 54
foveata, 73
trachycarpa, 71
Cyathipodia, 201
Corium, 208
Dupainii, 207
longipes, 203
platypodia, 203
Dasyscypha allantospora, 154
Dermea Craterium, 199
Detonia, 34, 54
Constellatio, 69
convexella, 68
foveata, 73
fulgens, 50
laeterubra, 69
leiocarpa, 73
miniata, 57
modesta, 63
nigrans, 72
polytrichina, 75
trachycarpa, 71
Discina, 43, **216**
abietina, 228
ancilis, 216
apiculata, 216, **217**
clypeata, 233
convoluta, 216, **218**

helvetica, 216
leucoxantha, 216, **218**
melaleuca, 225
perlata, 216
repanda, 231
reticulata, 229
trachycarpa, 71
venosa, 229
Warnei, 216
Disciotis, 219
venosa, 229
Durandiomyces, 236, **242**
Phillipsii, 242
Elvela, 237, **245**, 255
acaulis, 215
Acetabulum, 202
adhaerens, 245, **250**
albida, 175
albipes, 246, **253**
atra, 245, **248**
auricula, 185
brevissima, 246, **252**
brunnea, 251, 254
californica, 246, **250**
capucinoides, 249
caroliniana, 246, **253**
ciliata, 159
coccinea, 97, 191
cochleata, 221, 231
costata, 253
crispa, 245, **247**
elastica, 245, **249**
ephippium, 205
equina, 155
esculenta, 251
Faulknerae, 207
fistulosa, 249
fuliginosa, 249
fusca, 247
Gigas, 253
gracilis, 249
grisea, 246
Hegani, 253
hemisphaerica, 49
hispida, 205
hybrida, 241
inflata, 215
infula, 27, 28, 246, **251**, 252
Klotzschiana, 245, 249
lacunosa, 246, 248
leucophaea, 247
lutea, 169
macropus, 205
macropus brevis, 206
Mitra, 245, **246**, 251
nigra, 208
nigricans, 246, 248
nivea, 247
pallescens, 247
pallida, 247
palustris, 245, **247**
pileata, 255
rhodopus, 251
sphaerospora, 246, **254**, 255
Stevensii, 249
sulcata, 246
umbraculiformis, 246, **251**
Underwoodii, 246, **254**
Fleischhakia, 105
rhizinoides, 106
Fuckelina, 201
Corium, 208
pubida, 210
Schweinitzii, 211
semitosta, 209
villosa, 206
Galactinia, 219
badia, 221
Boltoni, 226
brunneoatra, 222
ionella, 138
proteana, 227
proteana sparassoides, 242
pustulata, 224
succosa, 223
Geopyxis, 43, **211**
aluticolor, 190
amplispora, 231
brachypus, 206
bronca, 212, **213**
carbonaria, 212
Catinus, 212, **213**
cinerascens, 214
coccinea, 191
Craterium, 199
cupularis, 212
elata, 188
floccosa, 192

hesperidea, 193
Hicksii, 215
nebulosa, 214
nebulosoides, 214
occidentalis, 193
pallidula, 231
pulchra, 214
sordescens, 99
suburceolata, 212
verruculosa, 214
vulcanalis, 212, **214**
Geoscypha Schröteri, 141
Gyromitra, 245
brunnea, 254
caroliniana, 253
curtipes, 253
esculenta, 27, 28, 251, 252
esculenta crispa, 251
Gigas, 242, 253
infula, 251
Phillipsii, 242
sphaerospora, 254
Helotium
axillaris, 124
flammeum, 154
Pedrottii, 141
pithyum, 76
purpuratum, 182
testaceum, 125
thujinum, 78
Helvella, 4, 245
Heteroplegma, 219
caeruleum, 225
crenatum, 228
Humaria, 121, 156
adusta, 167
aggregata, 136
alpina, 169
aquatica, 108
araneosa, 130, 151
axillaris, 124
bella, 140
bicucullata, 99
callichroa, 180
cestrica, 98
Charteri, 103
clausa, 125
coccinea, 137
convexula, 128
Cookeina, 137
Crec'hqueraultii, 62
cremoricolor, 117
cupressina, 78
deerrata, 141
deligata, 132
depressa, 106
diversispora, 155
echinosperma, 62
fimeti, 232
flavoaurantiaca, 168
fuscocarpa, 140
fusispora, 136
gallinacea, 118
Gerardi, 138
glumarum, 133
granulata, 116
gregaria, 176
groenlandica, 140
haemastigma, 68
hemisphaerica, 175
hepatica, 139
hirtella, 180
humosa, 129
hydrophila, 106
ionella, 138
ithacaensis, 124
lacteocinerea, 234
lechithina, 108
leucoloma, 129
macrospora, 222
macrocystis, 167
melaloma, 167
melalomoides, 167
microspora, 141
miniata, 103
muralis, 137
ochroleuca, 131
olivatra, 140
ollaris, 135
omphalodes, 109
Oocardii, 108
orthotricha, 127
Peckii, 133
permuda, 134
Persoonii amethystina, 67
phycophila, 127
phyllogena, 134
pityina, 76
pulcherrima, 172

rhodoleuca, 163
rubens, 127
rubricosa, 131
rutilans, 100
salmonicolor, 140
scatigena, 140
Schröteri, 141
scutellata, 159
secreta, 140
semiimmersa, 130
setosa, 166
silvosa, 134
spissa, 222
stercorea, 169
subcrenulata, 140
subgranulata, 141
subhirsuta, 171
testacea, 125
tetraspora, 134
theleboloides, 170
trachyderma, 139
umbrorum, 161
violacea, 226
vitigena, 140
wisconsinensis, 132
Wrightii, 66
xanthomela americana, 141
Humarina, 16, 22, 41, **121**, 141
aggregata, 123, **136**
araneosa, 123, **130**
axillaris, 22, 122, **124**
clausa, 122, **125**
Clementsii, 127
coccinea, 123, **137**
convexula, 122, **128**
Cookeina, 123, **137**
deligata, 123, **132**
fusispora, 123, **136**
Gerardi, 123, **138**
glumarum, 123, **133**
hepatica, 123, **139**
ithacaensis, 22, 122, **124**
leucoloma, 122, **129**, 140
lilacina, 122, **128**
nectrioides, 123, **132**
ochroleuca, 123, **131**
ollaris, 123, **135**
orthotricha, 122, **127**
pallens, 122, **124**
Peckii, 123, **133**
permuda, 123, **134**
phycophila, 122
phyllogena, 123, **134**
purpurea, 123, **138**
rubens, 122, **127**
rufa, 123, **131**
semiimmersa, 123, **130**
testacea, 122, **125**
testacea macrospora, 126
tetraspora, 123, **134**
trachyderma, 123, **139**
wisconsinensis, 123, **132**
Humariella, 157
melaloma, 167
scutellata, 159
stercorea, 169
Lachnea, 35, 156
abundans, 177
alpina, 169
amphidoxa, 176
arenicola, 149
arenosa, 151
aspera, 210
asperrima, 104
aurantia, 152
aurantiopsis, 198
badioberbis, 162
barbata, 104
bryophila, 178
chaetoloma, 175
chrysotricha, 178
Clementsi, 162
coccinea, 191
confusa, 46
coprinaria, 171
Corium, 208
cretea, 177
crispata, 179
crucipila, 168
cubensis, 160
dalmeniensis, 170
diplotricha, 170
dispora, 161
Erinaceus, 178
fissilis, 180
fusicarpa, 210
Geaster, 149
gigantea, 209

gilva, 166
gregaria, 176
gregaria pseudogregaria, 174
grisea, 151
Hainesii, 209
hemispherica, 175
heterospora, 161
heterothrix, 149
hinnulea, 46
hirta, 180
intermixta, 167
irregularis, 177
lanuginosa, 149
livida, 180
Lojkaeana, 170
macropus, 205
maculosa, 172
melaloma, 167
melastoma, 197
miniata, 103
mirabilis, 194
nigrella, 49
paludosa, 174
pediseta, 171
pellita, 152
piliseta, 160
pseudocrenulata, 178
pseudogregaria, 174
pulcherrima, 172
punicea, 165
pygmaea, 179
rubropurpurea, 161
scubalonta, 171
scutellata, 159
sepulta, 149
Sequoiae, 167
setosa, 166
stercorea, 169
stictica, 160
subcrinita, 166
Sumneriana, 149
texensis, 160
theleboloides, 170
trechispora, 43
umbrorum, 161
vitellina, 179
Woolhopeia, 178
Lachnella, 154
cinnabarina, 154
Cupressi, 78
flammea, 154
incarnescens, 154
maculincola, 154
rhoina, 154
Lachnellula chrysophthalma, 78
Lamprospora, 15, 16, 23, 31, 32, 34, 35, 39, **54**
amethystina, 55, **67**
annulata, 55, **60**
areolata, 55, **59**
areolata australis, 32, **59**
ascoboloides, 55, **60**
brevispinosa, 55, **63**
carbonaria, 56, **67**
carbonicola, 67, 68
Constellatio, 56, **69**
Crec'hqueraultii, 11, 45, 55, **62**, 64
Crec'hqueraultii macrantha, 63
Crouani, 54, **56**
Detonia, 56, **72**
dictydiola, 55, **58**
discoidea, 56, **70**
exasperata, 56, **75**
gemmea, 56, **70**
haemastigma, 56, **68**
laetirubra, 54, **58**
leiocarpa, 56, **73**
lobata, 56, **74**
macrantha, 55, **63**
Maireana, 32, 55, **65**
miniata, 56
nigrans, 56, **72**
Planchonis, 32, 56, **74**
polytrichina, 56, **75**
salmonicolor, 56, **71**
spinulosa, 55, **61**
trachycarpa, 56, **71**, 72
tuberculata, 32, 55, **64**, 65
tuberculatella, 32, 55, **66**
wisconsinensis, 56, **69**
Wrightii, 55, **66**
Lasiobolus, 41, **155**
brachyascus, 81
diversisporus, 155
equinus, 155
pilosus, 155
pulcherrimus, 172
raripilus, 173

ruber, 155, **156**
stercoreus, 169
Leotia conica, 243
Leptopodia, 245
atra, 248
Corium, 208
elastica, 246
Klotzschiana, 249
Leucoloma, 121
araneosa, 130
asperior, 43
axillaris, 124
coccinea, 137
Constellatio, 69
convexula, 128
fusispora, 136
Hedwigii, 129
rubricosa, 131
rutilans, 100
tetraspora, 134
Leucoscypha excipulata, 129, 130
Macropodia, 201
Corium, 208
fusicarpa, 210
macropus, 205
platypodia, 203
pubida, 210
Schweinitzii, 211
semitosta, 209
subclavipes, 206
urceolata, 206
Macroscyphus, 202
acetabuliforme, 202
coccineus, 191
macropus, 205
Melachroia, 141
xanthomela americana, 141
Melascypha, 47
melaena, 48
Melastiza, 40, **103**
asperrima, 103, **104**
Charteri, 103
miniata, 103
pennsylvanica, 103, **104**
Microcrater, 198, 200
Microstoma, 190
hiemale, 193
Midotis, 181
gigantea, 181
macrotis, 181
Mitrophora, 253
caroliniana, 253
hybrida, 241
patula, 241
rimosipes, 241
semilibera, 241
Mitrula irregularis, 256
Mollisia, 62
Crec'hqueraultii, 62
Morchella, 29, 236, **237**
angusticeps, 237, **240**
bispora, 244
bispora truncata, 244
bohemica, 244
bohemica bispora, 244
caroliniana, 253
conica, 237, **239**
conica deliciosa, 240
continua, 238, 239
crassipes, 237
deliciosa, 237, **240**
elata, 240
esculenta, 27, 237, **238**
esculenta conica, 239
esculenta fulva, 238
esculenta longipes, 238
esculenta rotunda, 238
esculenta vulgaris, 238
gigaspora, 244
hybrida, 237, **241**
hybrida rimosipes, 241
Mitra, 241
patula, 241
punctipes, 241
Rete, 241
rimosipes, 241
rotunda, 238
vulgaris, 238
Morilla, 237
bohemica, 244
conica, 239
elata, 240
esculenta, 238
gigaspora, 244
semilibera, 241
Neottiella, 156, 157
albotecta, 180
callichroa, 180

corallina, 317
crucipila, 168
Hetieri, 164
leucoloma, 129
macrospora, 163
ollaris, 135
ovilla, 163
ovilla flavodisca, 163
Polytrichi, 163
sericeovillosa, 180
Spraguei, 180
Neottiopezis, 157
macrospora, 163, 180
Octospora, 156
fasciculata, 175
haemastigma, 68
leucoloma, 122, 129
pustulata, 224
rhizophora, 215
scutellata, 159
varia, 230
villosa, 205
Ombrophila, 141
Otidea, 97, 184, 219, 221
abietina, 228
aurantia, 97
auricula, 185
dochmia, 184
domingensis, 182
grandis, 186
Harperiana, 186
hirneoloides, 184
leporina, 185
neglecta, 186
obtecta, 187
ochracea, 185
onotica, 185
succosa, 223
Otidella, 47
fulgens, 50
fuscocana, 48
nigrella, 49
Pachyella, 219
Barleana, 233
Patella, 21, 35, 41, **156**
abundans, 21, 22, 158, **177**
albida, 158, **175**, 176
albocincta, 157, **163**
albospadicea, 158, **178**
ciliata, 159
coprinaria, 158, **171**
crucipila, 158, **168**
cubensis, 157, **160**
Erinaceus, 158, **178**, 180
fimetaria, 158, **173**
gilva, 158, **166**
gregaria, 158, **176**
Hetieri, 158, **164**
irregularis, 158, **177**
Lusatiae, 157, **162**
lutea, 169
maculosa, 158, **172**
melaloma, 158, **167**
miniata, 157, **162**
ovilla, 158, **163**
paludosa, 158, **174**
piliseta, 157, **160**
pulcherrima, 158, **172**
punicea, 158, **165**
pygmaea, 158, **179**
raripila, 158, 173
ricciophila, 158, **165**
scutellata, 35, 157, **159**, 180
Sequoiae, 158, **167**
setosa, 158, **166**, 179
stercorea, 158, **169**
theleboloides, 158, **170**
tuberculata, 158, **174**
umbrorum, 157, **161**, 180
Paxina, 42, **201**
Acetabulum, 201, **202**
Barlae, 201, **205**
Corium, 201, **208**, 211
Dupainii, 201, **207**
fusicarpa, 202, **210**
hispida, 201, **205**
leucomelas, 201, **204**
macropus, 201, **203**
nigrella, 201, **208**
olivacea, 202, **211**
platypodia, 201, **203**
semitosta, 202, **209**
subclavipes, 201
sulcata, 4, 201, **204**
Pelodiscus, 157
miniatus, 162
pilisetus, 160
Peltidium, 105

Cookei, 108
lignarium, 108
Oocardii, 108
Perrotia, 41, **154**
flammea, 154
Plectania adusta, 199
Peziza, 15, 43, 121, 148, 156, **219**
abietina, 220, **228**
abundans, 177
Acetabulum, 201, **202**
Adae, 230
adusta, 167, 199
Afzelii, 188
aggregata, 136
albocincta, 163
albospadicea, 178
albotecta, 180
aluticolor, 190
ampelina, 226
amplispora, 231
amplissima, 235
ancilis, 216
apiculata, 217
aquatica, 108
araneosa, 130
arenicola, 149
arenosa, 150, **151**
asperior, 43
assimilata, 224
atrorufa, 197
atrospora, 71
atrovinosa, 101
aurantia, 96, **97**
aurantiaca, 159
aurantiopsis, 198
auricula, 185
auriflava, 62
Aurora, 115
axillaris, 124
Babingtonii, 106
badia, 32, 220, **221**, 235
badioberbis, 162
Barleana, 233
bella, 140
bicucullata, 99
Boltoni, 226
brachypus, 206
bronca, 213
brunnea, 46
brunneoatra, 220, **222**
brunneovinosa, 234
Burcardia, 195
caerulea, 225
caeruleomaculata, 234
caligans, 235
carbonaria, 212
Catinus, 213
cerea, 233
cervina, 155
cestrica, 98
Charteri, 103
chlamydospora, 101
chlorophysa, 232
chrisophthalma, 128
chrysopela, 234
ciliata, 157, 159, 169
cinerea, 118
cinerella, 120
cinnabarina, 154
Clissoni, 235
clypeata, 221, **233**
coccinea, 97, 191
cochleata, 219, 221
Colensoi, 190
Commonsii, 98
concentrica, 220, **228**
confluens, 109
confusa, 46
Constellatio, 69
convexella, 68
convexula, 128
convoluta, 229, 235
coprinaria, 171
corallina, 137
cordovensis, 182
Corium, 207, 208
coronaria, 235
Craterium, 199
cremoricolor, 117
crenata, 212, 228
crenulata, 197
cretea, 177
crispata, 179
Crouani, 56
cruciata, 193
crucipila, 168
cubensis, 160
Cupressi, 78

cupressina, 78
cupularis, 212
cyanoderma, 50
dalmeniensis, 170
deerrata, 141
deligata, 107, 132
depressa, 106
difformis, 115
diversicolor, 155
dochmia, 184
domiciliana, 220, **230**
domingensis, 182
Dudleyi, 191
echinosperma, 62, 63, 224
elachroa, 141
elaeodes, 217, 218
Emileia, 220, **224**
epidendra, 191
Erinaceus, 178
exasperata, 75
fimetaria, 170, 232
fimeti, 221, **232**
flammea, 154
flavovirens, 44
floccosa, 192
fulgens, 50
fuliginea, 76
fuscocana, 48
fuscocarpa, 140
fusicarpa, 210
fusispora, 136
fusispora permuda, 134
gallinacea, 118
Geaster, 149, 235
gemmea, 70
Gerardi, 138
gigantea, 235
gilva, 166
globifera, 76
glumarum, 133
grandis, 186
granulata, 116
granulosa, 116
griseorosea, 220, **227**
Hainesii, 209
Harmoge, 182
hemisphaerica, 49, 175
hepatica, 139
hesperidea, 193
Hindsii, 188
hinnulea, 46
hirneoloides, 184
hirta, 180
hirtipes, 197
hispida, 175
humicola, 213
humosa, 129
humosoides, 115
hyalinella, 145
hybrida, 152
hydrophila, 106
Hystrix, 187
inquinans, 195
intermixta, 167
ionella, 138
irregularis, 235
irrorata, 235
labellum, 175
laeticolor, 159
laetirubra, 58
lanuginosa, 149
lanuginosa Sumneri, 149
lechithina, 108
Leineri, 76
leiocarpa, 73
leporina, 184, 185
leucobasis, 107
leucoloma, 75, 129
leucomelas, 204
leucoxantha, 218
livida, 180
lobata, 74
Lusatiae, 162
macrocalyx, 235
macropus, 205
macrospora, 222
macrotis, 181
maculincola, 154
maculosa, 172
majalis, 214
melaena, 48
melaleuca, 220, **225**
melaleucoides, 220, **225**
melaloma, 167
melania, 48
melastoma, 197
Meleagris, 154
micropus, 231

microspora, 141
miniata, 103
mirabilis, 193
modesta, 62, 63
monilifera, 47
Morgani, 210
muralis, 137
nana, 222
nebulosa, 214
nigrans, 72
nigrella, 48, 49
nummularia, 106
obtecta, 187
occidentalis, 193
odorata, 230
olivacea, 140
olivatra, 140
ollaris, 135
omphalodes, 109
onotica, 185
onotica ochracea, 185
Oocardii, 108
Oocardii lignaria, 108
orbicularis, 233
orthotricha, 127
ovilla, 163
pallida, 231
papillata, 155
paraphysata, 233
pellita, 152, 197
perlata, 216
permuda, 134
pertenuis, 139
Petersii, 224
pithya, 76, 78
pithyna, 76
poculiformis, 191
polyspora, 146
Polytrichi, 75, 163
prostrata, 193
proteana, 220, **227**, 242
proteana sparassoides, 242
proximella, 176
pubida, 210
pulcherrima, 172, 191
pulchra, 214
pustulata, 220, **224**, 234
radiculata, 99
repanda, 21, 220, 230, **231**, 233, 235
repanda amplispora, 231
repandoides, 235
reticulata, 229
retiderma, 101
rhizomorpha, 197
rhizophora, 215
rhizopus, 197
rivularis, 106
roseolilacina, 235
rubra, 130
rubricosa, 131
rubrofusca, 228
rufa, 131
rufescens, 141
rutilans, 100
salmonicolor, 140
sanguinaria, 67
scabra, 116
scabrosa, 71
scatigena, 140
schizospora, 46
Schröteri, 141
scubalonta, 171
scutellata, 159, 180
scutelloides, 46
secreta, 140
semiimmersa, 130
semitosta, 209
sepulta, 149
Sequoiae, 167
setosa, 166
sicula, 235
sordescens, 99
sphaeroplea, 46
spissa, 140, 220, **222**
splendens, 99
spongiosa, 48
Spraguei, 180
stercoraria, 81, 82
stercorea, 169
Stevensoniana, 231
stictica, 160
stipitata, 205
striispora, 187
stygia, 49
subclavipes, 206
subfusca, 111
subgranulata, 141
suburceolata, 212, 213

succosa, 220, **223**
sulcata, 204
sulcipes, 188
sylvestris, 221, **233**
tapeina, 139
testacea, 125
tetraonalis, 119
tetraspora, 134
texensis, 160
theleboloides, 170
tomentosa, 211
trachycarpa, 71
trachyderma, 139
trechispora, 43
Tricholoma, 187
truncicomes, 231
umbrina, 224
umbrorum, 161
umbrosa, 161
unicisa, 185
varia lignicola, 231
varia terrestris, 233
varia typica, 230
Venezuelae, 190
venosa, 218, 220, **229**
verruculosa, 215
vesiculosa, 21, 221, **231**
violacea, 138, 220, **226**, 228
vitellina, 179
vogesiaca, 48
vulcanalis, 214
Warnei, 216
Woolhopeia, 178
Wrightii, 66
Pezizula, 144
Cesatii, 121
crustacea, 145
Phacidium terrestre, 200
Phaeopezia, 54, 101
apiculata, 217
atrospora, 71
elaeodes, 44
Nuttallii, 87, 88
retiderma, 101
scabrosa, 71
vinacea, 222
Phalloboletus, 237
angusticeps, 240
bisporus, 244
bohemicus, 244
conicus, 239, 243
crassipes, 237
deliciosus, 240
elatus, 240
esculentus, 238
gigasporus, 244
Rete, 241
rimosipes, 241
Phallus, 215
acaulis, 215
acuminatus, 239
conicus, 243
crassipes, 237
crispus, 247
esculentus, 237, 238
Monacellus, 251
Rete, 241
Phialea cupressina, 78
Phillipsia, 17–19, 42, **182**, 198, 235
Chardoniana, 18, 182, **183**
dochmia, 182, **184**
domingensis, 182
gigantea, 182, **183**
kermesina, 182
subpurpurea, 182
Venezuelae, 190
Phleboscyphus, 201
Acetabulum, 202
macropus, 203
olivaceus, 211
radicatus, 214
Physomitra, 245
esculenta, 251
infula, 251
Pilobolus, 26
Pilocratera, 187
Engleriana, 188
Hindsii, 188
Tricholoma, 187
Pithya, 39, **76**
Cupressi, 22, 76–**78**
cupressina, 78
lacunosa, 76, **77**
pithya, 76
suecica, 78
thujina, 78
vulgaris, 76
Plectania, 42, 141, **190**

coccinea, 191
floccosa, **191**, 192
hiemalis, 17, 19, 191, 193
hirtipes, 197
melastoma, 197
nigrella, 49
occidentalis, 191, **193**
rhizomorpha, 197
rimosa, 211
Plicaria, 54, 219
badia, 221
brunneoatra, 222
caeruleomaculata, 234
carbonaria, 71
ferruginea, 71
flavovirens, 44
foveata, 73
Fuckelii caerulescens, 225
muralis, 137
Planchonis, 74
pustulata, 224
repanda, 231
repandoides, 235
reticulata, 229
rubrofusca, 228
succosa, 223
tracheia, 52
Plicariella, 34, 54
Constellatio, 69
fulgens, 50
leiocarpa, 73
miniata, 57
modesta, 62
Polytrichi, 75
trachycarpa, 71
Podophacidium terrestre, 200
Pseudombrophila, 41, **141**
deerata, 132, **141**
Pedrottii, 141
Pseudopeziza, 92
Pseudopithyella, 41, **153**
minuscula, 153
Pseudoplectania, 39, **47**, 215
fulgens, 48, **50**
melaena, 48
nigrella, 48, **49**
stygia, 49
vogesiaca, **48**, 208, 209
Pseudotis, 219
abietina, 228
Psilopezia, 40, **105**, 110, 223
aquatica, 105, **108**
deligata, 107
flavida, 105, **108**
hydrophila, 105, **106**
juruensis, 233
mirabilis, 109
myrothecioides, 105, 106
nummularia, 105, **106**, 234
Oocardii, 108
orbicularis, 233
trachyspora, 105, **107**
Pulparia spongiosa, 48
Pulvinula, 54
carbonaria, 67
Constellatio, 69
haemastigma, 68
Pustularia, 211, 219
Catinus, 213
cupularis, 212
gigantea, 235
Stevensoniana, 231
vesiculosa, 231
Pyronema, 3, 6–8, 11, 23–25, 27, 40, **109**
aurantiorubrum, 109
carneum, 115
confluens, 109
confluens inigneum, 110
haemastigma, 68
leucobasis, 107, 110
Marianum, 109
melalomum, 167
omphalodes, 6, 12, 27, **109**
subhirsutum, 167, 170
Pyronemella monilifera, 47
Rhizina, 43, **215**
Babingtonii, 106
helvetica, 216
inflata, 215
laevigata, 215
myrothecioides, 106
spongiosa, 216
undulata, 215
Ruhlandiella, 51
berolinensis, 51
hesperia, 51
Ryparobius, 41, **144**

brunneus, 144
Cookei, 145
Crouani, 142
crustaceus, 144, **145**, 147
hyalinellus, 144, **145**
monascus, 144, **147**
niveus, 147
pachyascus, 146
Pelletieri, 148
polysporus, 144, **146**
sexdecimsporus, 144
woolhopensis, 143
Saccobolus, 40, **92**, 96
depauperatus, 93, **95**
globuliferellus, 93, **95**
Kerverni, 93
neglectus, 94
obscurus, 96
portoricensis, 93, **94**
violascens, 93, **94**
Sarcoscypha, 190
albovillosa, 100
alpina, 193, 194
coccinea, 191, 192
Colensoi, 190
Corium, 208
cruciata, 193
dawsonensis, 214
floccosa, 192
melastoma, 197
minuscula, 153
occidentalis, 193
prostrata, 194
radiculata, 99
rhenana, 99
roseotincta, 194
striispora, 187
Sarcosoma, 194
globosa, 195
Sarcosphaera, 43, **235**
arenicola, 149
arenosa, 151
coronaria, 235
Geaster, 149
macrocalyx, 235
Sumneriana, 149
Sclerotinia, 13, 21, 215
baccata, 193
Sclerotium, 133
Scodellina, 42, **184**
aurantiaca, 97
auricula, 185
badia, 221
grandis, 185, **186**
leporina, 185
onotica, 185
pustulata, 224
vesiculosa, 231
Scutellinia, 156, 157
abundans, 177
adusta, 199
albospadicea, 178
alpina, 169
arenicola, 149
arenosa, 151
aurantiopsis, 198
badeoberbis, 162
brunnea, 46
chaetoloma, 175
coprinaria, 171
cretea, 177
dalmeniensis, 170
dispora, 161
Erinaceus, 178
fusicarpa, 210
Geaster, 149
gilva, 166
gregaria, 176
Hainesii, 209
hemisphaerica, 175
heterospora, 161
hirtipes, 197
hybrida, 152
irregularis, 177
laeticolor, 159
lanuginosa, 149
Lusatiae, 162
melaloma, 167
melastoma, 197
miniata, 103
pellita, 152
pulcherrima, 172
rhizomorpha, 197
scubalonta, 171
scutellata, 159
sepulta, 149
Sequoiae, 167
setosa, 166

stercorea, 169
stictica, 160
Sumneriana, 149
texensis, 160
theleboloides, 170
umbrarum, 161
Woolhopeia, 178
Sepultaria, 41, **148**
albida, 175
arenicola, **149**, 150
arenosa, 149–**151**, 180
aspera, 210
aurantia, 152
aurantiaca, 149
bryophila, 178
Geaster, 149, 180
gigantea, 209
grisea, 151
heterothrix, 149
Longii, 149, **150**
pediseta, 171
pellita, 149, **152**
pseudocrenulata, 178
punicea, 165
pygmaea, 179
rubropurpurea, 161
semiimmersa, 130
semitosta, 209
sepulta, 149, 150
Sumneri, 149
tenuis, 180
Sphaeridiobolus, 79
hyperboreus, 79
hyperboreus niveus, 79
Sphaerosoma, 39, **51**
alveolatum, 53
echinulatum, 11, 52
fuscescens, 51
hesperium, 51
Sphaerospora, 39, **43**
asperior, 44
brunnea, 43, **46**
confusa, 46
Durandi, 44
flavovirens, 43, **44**
hinnulea, 43, **46**
monilifera, 43, **47**
nigrella, 49
perplexa, 43, **45**
scutelloides, 46
trechispora, 43
Sphaerosporula, 43
asperior, 44
confusa, 46
flavovirens, 44
scutelloides, 46
trechispora, 44
Sphaerozosma, 51
Spragueola, 256
Streptotheca, 16, 41, **142**
Boudieri, 142
Crouani, 142
obscura, 142, **143**
woolhopensis, 142, **143**
Tapesia omphalodes, 109
Tarzetta cinerascens, 214
Thecotheus, 41, **147**
Pelletieri, 148
Thelebolus, 144, 147
lignicola, 147
monascus, 147
nanus, 147
stercoreus, 147
Tricharia, 157
gilva, 166
Trichophaea, 157
abundans, 177
albospadicea, 178
gregaria, 176
paludosa, 174
pseudogregaria, 174
Trichoscypha, 187
Afzelii, 188
Hindsii, 188
sulcipes, 188
Tricholoma, 187
Underwoodia, 237, **255**
columnaris, 255
Urnula, 42, **198**
Craterium, 199
Geaster, 199, **200**
melastoma, 197
terrestris, 200
Verpa, 237, **243**
atroalba, 245
bohemica, 243, **244**

chicoensis, 243
conica, 243, **245**
digitaliformis, 243
helvelloides, 243
Krombholzii, 243
perpusilla, 245
Wynnea, 4, 17–19, 42, **180**
americana, 19, 180, **181**
gigantea, 180, **181**
macrotis, 181
Wynnella, 184
auricula, 186